

## 65. Continued

$$\begin{aligned}
 \text{(a)} &= 2 \text{ (Area of triangle } ORT) \\
 &= 2 \cdot \frac{1}{2} \text{ (} x\text{-intercept of line } RT) \text{ (} y\text{-intercept of line } RT) \\
 &= 2 \cdot \frac{1}{2} \left( \frac{mx_0 - a}{m} \right) (a - mx_0) \\
 &= -m \left( \frac{mx_0 - a}{m} \right) \left( \frac{mx_0 - a}{m} \right) \\
 &= - \left( \frac{mx_0 - a}{m} \right)^2 \\
 &= -m \left( x_0 - \frac{a}{m} \right)^2
 \end{aligned}$$

Substituting  $x$  for  $x_0$ ,  $f'(x)$  for  $m$ , and  $f(x)$  for  $a$ , we

$$\text{have } A(x) = -f'(x) \left[ x - \frac{f(x)}{f'(x)} \right]^2.$$


(b) The domain is the open interval  $(0, 10)$ .

$$\text{To graph, let } y_1 = f(x) = 5 + 5\sqrt{1 - \frac{x^2}{100}},$$

$$y_2 = f'(x) = \text{NDER}(y_1), \text{ and}$$

$$y_3 = A(x) = -y_2 \left( x - \frac{y_1}{y_2} \right)^2.$$

The graph of the area function  $y_3 = A(x)$  is shown below.



$[0, 10]$  by  $[-100, 1000]$

The vertical asymptotes at  $x = 0$  and  $x = 10$  correspond to horizontal or vertical tangent lines, which do not form triangles.

(c) Using our expression for the  $y$ -intercept of the tangent line, the height of the triangle is

$$\begin{aligned}
 a - mx &= f(x) - f'(x) \cdot x \\
 &= 5 + \frac{1}{2}\sqrt{100 - x^2} - \frac{-x}{2\sqrt{100 - x^2}}x \\
 &= 5 + \frac{1}{2}\sqrt{100 - x^2} + \frac{x^2}{2\sqrt{100 - x^2}}
 \end{aligned}$$

We may use graphing methods or the analytic method in part (d) to find that the minimum value of  $A(x)$  occurs at  $x \approx 8.66$ . Substituting this value into the expression above, the height of the triangle is 15. This is 3 times the  $y$ -coordinate of the center of the ellipse.

(d) Part (a) remains unchanged. The domain is  $(0, C)$ . To graph, note that

$$f(x) = B + B\sqrt{1 - \frac{x^2}{C^2}} = B + \frac{B}{C}\sqrt{C^2 - x^2} \text{ and}$$

$$f'(x) = \frac{B}{C} \frac{1}{2\sqrt{C^2 - x^2}} (-2x) = \frac{-Bx}{C\sqrt{C^2 - x^2}}.$$

Therefore, we have

$$\begin{aligned}
 A(x) &= -f'(x) \left[ x - \frac{f(x)}{f'(x)} \right]^2 \\
 &= \frac{Bx}{C\sqrt{C^2 - x^2}} \left[ x - \frac{B + \frac{B}{C}\sqrt{C^2 - x^2}}{\frac{-Bx}{C\sqrt{C^2 - x^2}}} \right]^2 \\
 &= \frac{Bx}{C\sqrt{C^2 - x^2}} \left[ x - \frac{(BC + B\sqrt{C^2 - x^2})\sqrt{C^2 - x^2}}{-Bx} \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[ Bx^2 + (BC + B\sqrt{C^2 - x^2})(\sqrt{C^2 - x^2}) \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[ Bx^2 + BC\sqrt{C^2 - x^2} + B(C^2 - x^2) \right]^2 \\
 &= \frac{1}{BCx\sqrt{C^2 - x^2}} \left[ BC(C + \sqrt{C^2 - x^2}) \right]^2 \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})^2}{x\sqrt{C^2 - x^2}}
 \end{aligned}$$

$$\begin{aligned}
 A'(x) &= BC \cdot \frac{(x\sqrt{C^2 - x^2})(2)(C + \sqrt{C^2 - x^2}) \left( \frac{-x}{\sqrt{C^2 - x^2}} \right) - (C + \sqrt{C^2 - x^2})^2 \left( x \frac{-x}{\sqrt{C^2 - x^2}} + \sqrt{C^2 - x^2} (1) \right)}{x^2(C^2 - x^2)} \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)} \left[ \frac{-2x^2 - (C + \sqrt{C^2 - x^2})}{\left( \frac{-x^2}{\sqrt{C^2 - x^2}} + \sqrt{C^2 - x^2} \right)} \right] \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2\sqrt{C^2 - x^2}} \left[ \frac{-2x^2 + \frac{Cx^2}{\sqrt{C^2 - x^2}}}{-C\sqrt{C^2 - x^2} + x^2 - (C^2 - x^2)} \right] \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)} \left( \frac{Cx^2}{\sqrt{C^2 - x^2}} - C\sqrt{C^2 - x^2} - C^2 \right) \\
 &= \frac{BC(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)^{3/2}} \left[ Cx^2 - C(C^2 - x^2) - C^2\sqrt{C^2 - x^2} \right] \\
 &= \frac{BC^2(C + \sqrt{C^2 - x^2})}{x^2(C^2 - x^2)^{3/2}} (2x^2 - C^2 - C\sqrt{C^2 - x^2})
 \end{aligned}$$

To find the critical points for  $0 < x < C$ , we solve:

$$\begin{aligned}
 2x^2 - C^2 &= C\sqrt{C^2 - x^2} \\
 4x^4 - 4C^2x^2 + C^4 &= C^4 = C^2x^2 \\
 4x^4 - 3C^2x^2 &= 0 \\
 x^2(4x^2 - 3C^2) &= 0
 \end{aligned}$$

The minimum value of  $A(x)$  for  $0 < x < C$  occurs at the critical point  $x = \frac{C\sqrt{3}}{2}$ , or  $x^2 = \frac{3C^2}{4}$ . The corresponding triangle height is


65. Continued

$$\begin{aligned}
 a - mx &= f(x) - f'(x) \cdot x \\
 &= B + \frac{B}{C} \sqrt{C^2 - x^2} + \frac{Bx^2}{C\sqrt{C^2 - x^2}} \\
 &= B + \frac{B}{C} \sqrt{C^2 - \frac{3C^2}{4}} + \frac{B \left( \frac{3C^2}{4} \right)}{C\sqrt{C^2 - \frac{3C^2}{4}}} \\
 &= B + \frac{B \left( \frac{C}{2} \right)}{\frac{C^2}{2}} + \frac{3BC^2}{C^2} \\
 &= B + \frac{B}{2} + \frac{3B}{2} \\
 &= 3B
 \end{aligned}$$

This shows that the triangle has minimum area when its height is  $3B$ .

**Section 4.5** Linerization and Newton's Method (pp. 233–245)

**Exploration 1** Appreciating Local Linearity



$$y = (x^2 + 0.0001)^{1/4} + 0.9$$

The function appears to come to a point.

$$\begin{aligned}
 2. \quad f'(a) &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \\
 &= \lim_{x \rightarrow a} \frac{(x^2 + 0.0001)^{1/4} + 0.9 - ((0 + 0.0001)^{1/4} + 0.9)}{x - 0} \\
 &= \lim_{x \rightarrow a} \frac{(x^2 + 0.0001)^{1/4} - 0.1}{x} = 0
 \end{aligned}$$

$f(x)$  is differentiable at  $x = 0$ , and the equation of the tangent line is  $y = 1$ .

3. The graph of the function at that point seems to become the graph of a straight line with repeated zooming.

4. The graph will eventually look like the tangent line.



**Exploration 2** Using Newton's Method on Your Calculator

See text page 237.

**Quick Review 4.5**

1.  $\frac{dy}{dx} = \cos(x^2 + 1) \cdot \frac{d}{dx}(x^2 + 1) = 2x \cos(x^2 + 1)$

2. 
$$\begin{aligned}
 \frac{dy}{dx} &= \frac{(x+1)(1-\sin x) - (x+\cos x)(1)}{(x+1)^2} \\
 &= \frac{x - x \sin x + 1 - \sin x - x - \cos x}{(x+1)^2} \\
 &= \frac{1 - \cos x - (x+1) \sin x}{(x+1)^2}
 \end{aligned}$$



5.  $f'(x) = (x)(-e^{-x}) + (e^{-x})(1) = e^{-x} - xe^{-x}$   
 $f'(0) = 1$   
 The line passes through  $(0, 1)$  and has slope 1. Its equation is  $y = x + 1$ .

6.  $f'(x) = (x)(-e^{-x}) + (e^{-x})(1) = e^{-x} - xe^{-x}$   
 $f'(-1) = e^1 - (-e^1) = 2e$   
 The line passes through  $(-1, -e + 1)$  and has slope  $2e$ . Its equation is  $y = 2e(x + 1) + (-e + 1)$ , or  $y = 2ex + e + 1$ .

7. (a)  $x + 1 = 0$   
 $x = -1$   
 (b)  $2ex + e + 1 = 0$   
 $2ex = -(e + 1)$   
 $x = -\frac{e + 1}{2e} \approx -0.684$

8.  $f'(x) = 3x^2 - 4$   
 $f'(1) = 3(1)^2 - 4 = -1$   
 Since  $f(1) = -2$  and  $f'(1) = -1$ , the graph of  $g(x)$  passes through  $(1, -2)$  and has slope  $-1$ . Its equation is  $g(x) = -1(x - 1) + (-2)$ , or  $g(x) = -x - 1$ .


$x$	$f(x)$	$g(x)$
0.7	-1.457	-1.7
0.8	-1.688	-1.8
0.9	-1.871	-1.9
1.0	-2	-2
1.1	-2.069	-2.1
1.2	-2.072	-2.2
1.3	-2.003	-2.3

9.  $f'(x) = \cos x$   
 $f'(1.5) = \cos 1.5$   
 Since  $f(1.5) = \sin 1.5$  and  $f'(1.5) = \cos 1.5$ , the tangent line passes through  $(1.5, \sin 1.5)$  and has slope  $\cos 1.5$ . Its equation is  $y = (\cos 1.5)(x - 1.5) + \sin 1.5$ , or approximately  $y = 0.071x + 0.891$



$[0, \pi]$  by  $[-0.2, 1.3]$

10. For  $x > 3$ ,  $f'(x) = \frac{1}{2\sqrt{x-3}}$ , and so  $f'(4) = \frac{1}{2}$ . Since  $f(4) = 1$  and  $f'(4) = \frac{1}{2}$ , the tangent line passes through  $(4, 1)$  and has slope  $\frac{1}{2}$ . Its equation is  $y = \frac{1}{2}(x - 4) + 1$ , or  $y = \frac{1}{2}x - 1$ .



$[-1, 7]$  by  $[-2, 2]$

### Section 4.5 Exercises

1. (a)  $f'(x) = 3x^2 - 2$   
 We have  $f(2) = 7$  and  $f'(2) = 10$ .  
 $L(x) = f(2) + f'(2)(x - 2)$   
 $= 7 + 10(x - 2)$   
 $= 10x - 13$

- (b) Since  $f(2.1) = 8.061$  and  $L(2.1) = 8$ , the approximation differs from the true value in absolute value by less than  $10^{-1}$ .

2. (a)  $f'(x) = \frac{1}{2\sqrt{x^2+9}}(2x) = \frac{x}{\sqrt{x^2+9}}$

We have  $f(-4) = 5$  and  $f'(-4) = -\frac{4}{5}$ .

$$\begin{aligned} L(x) &= f(-4) + f'(-4)(x - (-4)) \\ &= 5 - \frac{4}{5}(x + 4) \\ &= -\frac{4}{5}x + \frac{9}{5} \end{aligned}$$

- (b) Since  $f(-3.9) \approx 4.9204$  and  $L(-3.9) = 4.92$ , the approximation differs from the true value by less than  $10^{-3}$ .

3. (a)  $f'(x) = 1 - x^{-2}$   
 We have  $f(1) = 2$  and  $f'(1) = 0$ .  
 $L(x) = f(1) + f'(1)(x - 1)$   
 $= 2 + 0(x - 1)$   
 $= 2$

- (b) Since  $f(1.1) \approx 2.009$  and  $L(1.1) = 2$ , the approximation differs from the true value by less than  $10^{-2}$ .

4. (a)  $f'(x) = \frac{1}{x+1}$   
 We have  $f(0) = 0$  and  $f'(0) = 1$ .  
 $L(x) = f(0) + f'(0)(x - 0)$   
 $= 0 + 1x$   
 $= x$

- (b) Since  $f(0.1) \approx 0.0953$  and  $L(0.1) = 0.1$ , the approximation differs from the true value by less than  $10^{-2}$ .

5. (a)  $f'(x) = \sec^2 x$   
 We have  $f(\pi) = 0$  and  $f'(\pi) = 1$ .  
 $L(x) = f(\pi) + f'(\pi)(x - \pi)$   
 $= 0 + 1(x - \pi)$   
 $= x - \pi$

- (b) Since  $f(\pi + 0.1) \approx 0.10033$  and  $L(\pi + 0.1) = 0.1$ , the approximation differs from the true value in absolute value by less than  $10^{-3}$ .

6. (a)  $f'(x) = -\frac{1}{\sqrt{1-x^2}}$

We have  $f(0) = \frac{\pi}{2}$  and  $f'(0) = -1$ .

$$\begin{aligned} L(x) &= f(0) + f'(0)(x - 0) \\ &= \frac{\pi}{2} + (-1)(x - 0) \\ &= -x + \frac{\pi}{2} \end{aligned}$$

## 6. Continued

(b) Since  $f(0.1) \approx 1.47063$  and  $L(0.1) \approx 1.47080$ , the approximation differs from the true value in absolute value by less than  $10^{-3}$ .

$$7. f'(x) = k(1+x)^{k-1}$$

We have  $f(0) = 1$  and  $f'(0) = k$ .

$$\begin{aligned} L(x) &= f(0) + f'(0)(x-0) \\ &= 1 + k(x-0) \\ &= 1 + kx \end{aligned}$$

$$8. (a) (1.002)^{100} = (1+0.002)^{100} \approx 1 + (100)(0.002) = 1.2;$$

$$|1.002^{100} - 1.2| \approx 0.021 < 10^{-1}$$

$$(b) \sqrt[3]{1.009} = (1+0.009)^{1/3} \approx 1 + \frac{1}{3}(0.009) = 1.003;$$

$$|\sqrt[3]{1.009} - 1.003| \approx 9 \times 10^{-6} < 10^{-5}$$

$$9. (a) f(x) = (1-x)^6 = [1+(-x)]^6 \approx 1 + 6(-x) = 1 - 6x$$

$$(b) f(x) = \frac{2}{1-x} = 2[1+(-x)]^{-1} \approx 2[1+(-1)(-x)] = 2 + 2x$$

$$(c) f(x) = (1+x)^{-1/2} \approx 1 + \left(-\frac{1}{2}\right)x = 1 - \frac{x}{2}$$

$$10. (a) f(x) = (4+3x)^{1/3} = 4^{1/3} \left(1 + \frac{3x}{4}\right)^{1/3}$$

$$\approx 4^{1/3} \left(1 + \frac{1}{3} \left(\frac{3x}{4}\right)\right) = 4^{1/3} \left(1 + \frac{x}{4}\right)$$

$$(b) f(x) = \sqrt{2+x^2} = \sqrt{2} \left(1 + \frac{x^2}{2}\right)^{1/2}$$

$$\approx \sqrt{2} \left(1 + \frac{1}{2} \left(\frac{x^2}{2}\right)\right) = \sqrt{2} \left(1 + \frac{x^2}{4}\right)$$

$$(c) f(x) = \left(1 - \frac{1}{2+x}\right)^{2/3} = \left[1 + \left(-\frac{1}{2+x}\right)\right]^{2/3}$$

$$\approx 1 + \frac{2}{3} \left(-\frac{1}{2+x}\right) = 1 - \frac{2}{6+3x}$$

$$11. x = 100$$

$$f'(100) = \frac{1}{2}(100)^{-1/2} = 0.05$$

$$f(100) = 10 + 0.05(101 - 100) = 10.05$$

$$12. x = 27$$

$$f'(27) = \frac{1}{3}(27)^{-2/3} = \frac{1}{27}$$

$$f(27) = 3 + (1/27)(26 - 27)$$

$$y = 3 - \frac{1}{27} \approx 2.962$$

$$13. x = 1000$$

$$f'(1000) = \frac{1}{3}(1000)^{-2/3} = \frac{1}{300}$$

$$y = 10 + (1/300)(x - 1000)$$

$$y = 10 - \frac{1}{150} = 9.99\bar{3}$$

$$14. x = 81$$

$$f'(81) = \frac{1}{2}(81)^{-1/2} = \frac{1}{18}$$

$$y = 9 + \frac{1}{18}(80 - 81)$$

$$y = 9 - \frac{1}{18} = 8.9\bar{4}$$

15. Let  $f(x) = x^3 + x - 1$ . Then  $f'(x) = 3x^2 + 1$  and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^3 + x_n - 1}{3x_n^2 + 1}$$

Note that  $f$  is cubic and  $f'$  is always positive, so there is exactly one solution. We choose  $x_1 = 0$ .

$$x_1 = 0$$

$$x_2 = 1$$

$$x_3 = 0.75$$

$$x_4 \approx 0.6860465$$

$$x_5 \approx 0.6823396$$

$$x_6 \approx 0.6823278$$


$$x_7 \approx 0.6823278$$

Solution:  $x \approx 0.6823278$ .

16. Let  $f(x) = x^4 + x - 3$ . Then  $f'(x) = 4x^3 + 1$  and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^4 + x_n - 3}{4x_n^3 + 1}$$

The graph of  $y = f(x)$  shows that  $f(x) = 0$  has two solutions.



$[-3, 3]$  by  $[-4, 4]$

$$x_1 = -1.5$$

$$x_2 = -1.455$$

$$x_3 \approx -1.4526332$$

$$x_4 \approx -1.4526269$$

$$x_5 \approx -1.4526269$$

$$x_1 = 1.2$$

$$x_2 \approx 1.6541962$$

$$x_3 \approx 1.1640373$$

$$x_4 \approx 1.1640351$$

$$x_5 \approx 1.1640351$$

Solution:  $x \approx -1.452627, 1.164035$


17. Let  $f(x) = x^2 - 2x + 1 - \sin x$ .

Then  $f'(x) = 2x - 2\cos x$  and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^2 - 2x_n + 1 - \sin x_n}{2x_n - 2\cos x_n}$$

## 17. Continued

The graph of  $y = f(x)$  shows that  $f(x) = 0$  has two solutions



$[-4, 4]$  by  $[-3, 3]$

$$\begin{aligned}x_1 &= 0.3 & x_1 &= 2 \\x_2 &\approx 0.3825699 & x_2 &\approx 1.9624598 \\x_3 &\approx 0.3862295 & x_3 &\approx 1.9615695 \\x_4 &\approx 0.3862369 & x_4 &\approx 1.9615690 \\x_5 &\approx 0.3862369 & x_5 &\approx 1.9615690\end{aligned}$$

Solutions:  $x \approx 0.386237, 1.961569$

18. Let  $f(x) = x^4 - 2$ . Then  $f'(x) = 4x^3$  and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^4 - 2}{4x_n^3}$$

Note that  $f(x) = 0$  clearly has two solutions, namely

$x = \pm\sqrt[4]{2}$ . We use Newton's method to find the decimal equivalents.

$$\begin{aligned}x_1 &= 1.5 \\x_2 &\approx 1.2731481 \\x_3 &\approx 1.1971498 \\x_4 &\approx 1.1892858 \\x_5 &\approx 1.1892071 \\x_6 &\approx 1.1892071\end{aligned}$$

Solutions:  $x \approx \pm 1.189207$

19. (a) Since  $\frac{dy}{dx} = 3x^2 - 3$ ,  $dy = (3x^2 - 3)dx$ .

(b) At the given values,  
 $dy = (3 \cdot 2^2 - 3)(0.05) = 9(0.05) = 0.45$ .

20. (a) Since  $\frac{dy}{dx} = \frac{(1+x^2)(2) - (2x)(2x)}{(1+x^2)^2} = \frac{2-2x^2}{(1+x^2)^2}$ ,

$$dy = \frac{2-2x^2}{(1+x^2)^2} dx.$$

(b) At the given values,

$$\begin{aligned}dy &= \frac{2-2(-2)^2}{[1+(-2)^2]^2}(0.1) = \frac{2-8}{5^2}(0.1) \\&= -0.024.\end{aligned}$$

21. (a) Since  $\frac{dy}{dx} = (x^2)\left(\frac{1}{x}\right) + (\ln x)(2x) = 2x \ln x + x$ ,

$$dy = (2x \ln x + x)dx.$$

(b) At the given values,

$$dy = [2(1) \ln(1) + 1](0.01) = 1(0.01) = 0.01$$

22. (a) Since  $\frac{dy}{dx} = (x)\left(\frac{1}{2\sqrt{1-x^2}}\right)(-2x) + (\sqrt{1-x^2})(1)$

$$= \frac{-x^2}{\sqrt{1-x^2}} + \sqrt{1-x^2} = \frac{-x^2 + (1-x^2)}{\sqrt{1-x^2}} = \frac{1-2x^2}{\sqrt{1-x^2}},$$

$$dy = \frac{1-2x^2}{\sqrt{1-x^2}} dx.$$

(b) At the given values,  $dy = \frac{1-2(0)^2}{\sqrt{1-(0)^2}}(-0.2) = -0.2$ .

23. (a) Since  $\frac{dy}{dx} = e^{\sin x} \cos x$ ,  $dy = (\cos x)e^{\sin x} dx$ .

(b) At the given values,

$$dy = (\cos \pi)(e^{\sin \pi})(-0.1) = (-1)(1)(-0.1) = 0.1.$$

24. (a) Since  $\frac{dy}{dx} = -3 \csc\left(1-\frac{x}{3}\right) \cot\left(1-\frac{x}{3}\right)\left(-\frac{1}{3}\right)$

$$= \csc\left(1-\frac{x}{3}\right) \cot\left(1-\frac{x}{3}\right),$$

$$dy = \csc\left(1-\frac{x}{3}\right) \cot\left(1-\frac{x}{3}\right) dx.$$

(b) At the given values,

$$\begin{aligned}dy &= \csc\left(1-\frac{1}{3}\right) \cot\left(1-\frac{1}{3}\right)(0.1) \\&= 0.1 \csc \frac{2}{3} \cot \frac{2}{3} \approx 0.205525\end{aligned}$$

25. (a)  $y + xy - x = 0$

$$y(1+x) = x$$

$$y = \frac{x}{x+1}$$

$$\text{Since } \frac{dy}{dx} = \frac{(x+1)(1) - (x)(1)}{(x+1)^2} = \frac{1}{(x+1)^2},$$

$$dy = \frac{dx}{(x+1)^2}.$$

(b) At the given values,

$$dy = \frac{0.01}{(0+1)^2} = 0.01.$$

26. (a)  $2y = x^2 - xy$

$$2dy = 2x dx - x dy - y dx$$

$$dy(2+x) = (2x-y)dx$$

$$dy = \left(\frac{2x-y}{2+x}\right) dx$$

(b) At the given values, and  $y = 1$  from the original

$$\text{equation, } dy = \left(\frac{2(2)-1}{2+2}\right)(-0.05) = -0.0375$$

$$27. \frac{dy}{dx} = \sqrt{1-x^2}$$

$$dy = \left( -\frac{2x}{2\sqrt{1-x^2}} \right) dx$$

$$dy = -\frac{x}{\sqrt{1-x^2}} dx$$

$$28. \frac{dy}{dx} = e^{5x} + x^5$$

$$dy = (5e^{5x} + 5x^4) dx$$

$$29. \frac{dy}{dx} = \tan^{-1} 4x$$

$$\frac{d}{dx} \tan^{-1} u = \frac{1}{1+u^2} \frac{du}{dx}$$

$$u = 4x$$

$$\frac{du}{dx} = 4$$

$$dy = \left( \frac{4}{1+16x^2} \right) dx$$

$$30. \frac{dy}{dx} = (8^x + x^8)$$

$$\frac{d}{dx} a^x = (\ln a) a^x$$

$$dy = (8^x \ln 8 + 8x^7) dx$$

$$31. (a) \Delta f = f(0.1) - f(0) = 0.21 - 0 = 0.21$$

$$(b) \text{ Since } f'(x) = 2x + 2, f'(0) = 2.$$

$$\text{Therefore, } df = 2 dx = 2(0.1) = 0.2.$$

$$(c) |\Delta f - df| = |0.21 - 0.2| = 0.01$$

$$32. (a) \Delta f = f(1.1) - f(1) = 0.231 - 0 = 0.231$$

$$(b) \text{ Since } f'(x) = 3x^2 - 1, f'(1) = 2.$$

$$\text{Therefore, } df = 2 dx = 2(0.1) = 0.2.$$

$$(c) |\Delta f - df| = |0.231 - 0.2| = 0.031$$

$$33. (a) \Delta f = f(0.55) - f(0.5) = \frac{20}{11} - 2 = -\frac{2}{11}$$

$$(b) \text{ Since } f'(x) = -x^{-2}, f'(0.5) = -4.$$

$$\text{Therefore, } df = -4 dx = -4(0.05) = -0.2 = -\frac{1}{5}$$

$$(c) |\Delta f - df| = \left| -\frac{2}{11} + \frac{1}{5} \right| = \frac{1}{55}$$

$$34. (a) \Delta f = f(1.01) - f(1) = 1.04060401 - 1 = 0.04060401$$

$$(b) \text{ Since } f'(x) = 4x^3, f'(1) = 4.$$

$$\text{Therefore, } df = 4 dx = 4(0.01) = 0.04.$$

$$(c) |\Delta f - df| = |0.04060401 - 0.04| = 0.00060401$$

$$35. \text{ Note that } \frac{dV}{dr} = 4\pi r^2, dV = 4\pi r^2 dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in volume is approximately } 4\pi a^2 dr.$$

$$36. \text{ Note that } \frac{dS}{dr} = 8\pi r, \text{ so } dS = 8\pi r dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in surface area is approximately } 8\pi a dr.$$

$$37. \text{ Note that } \frac{dV}{dx} = 3x^2, \text{ so } dV = 3x^2 dx. \text{ When } x \text{ changes from } a \text{ to } a + dx, \text{ the change in volume is approximately } 3a^2 dx.$$

$$38. \text{ Note that } \frac{dS}{dx} = 12x, \text{ so } dS = 12x dx. \text{ When } x \text{ changes from } a \text{ to } a + dx, \text{ the change in surface area is approximately } 12a dx.$$

$$39. \text{ Note that } \frac{dV}{dr} = 2\pi rh, \text{ so } dV = 2\pi rh dr. \text{ When } r \text{ changes from } a \text{ to } a + dr, \text{ the change in volume is approximately}$$

$$40. \text{ Note that } \frac{dS}{dh} = 2\pi r, \text{ so } dS = 2\pi r dh. \text{ When } h \text{ changes from } a \text{ to } a + dh, \text{ the change in lateral surface area is approximately } 2\pi r dh.$$

$$41. A = \pi r^2$$

$$dA = 2\pi r dr$$

$$dA = 2\pi(10)(0.1) = 6.3 \text{ in}^2$$

$$42. v = \frac{4}{3}\pi r^3$$

$$dV = 4\pi r^2 dr$$

$$dV = 4\pi(8)^2(0.3) = 241 \text{ in}^2$$

$$43. v = s^3$$

$$dV = 3s^2 ds$$

$$dV = 3(15)^2(0.2) = 135 \text{ cm}^2$$

$$44. A = \frac{\sqrt{3}}{4}s^2$$

$$dA = \frac{\sqrt{3}}{2}s ds$$

$$dA = \frac{\sqrt{3}}{2}(20)(0.5) = 8.7 \text{ cm}^2$$

$$45. (a) \text{ Note that } f'(0) = \cos 0 = 1.$$

$$L(x) = f(0) + f'(0)(x-0) = 1 + 1x = x + 1$$

$$(b) f(0.1) = L(0.1) = 1.1$$

45. Continued

(c) The actual value is less than 1.1. This is because the derivative is decreasing over the interval  $[0, 0.1]$ , which means that the graph of  $f(x)$  is concave down and lies below its linearization in this interval.

46. (a) Note that  $A = \pi r^2$  and  $\frac{dA}{dr} = 2\pi r$ , so  $dA = 2\pi r dr$ .

When  $r$  changes from  $a$  to  $a + dr$ , the change in area is approximately  $2\pi a dr$ . Substituting 2 for  $a$  and 0.02 for  $dr$ , the change in area is approximately  $2\pi(2)(0.02) = 0.08\pi \approx 0.2513$

(b)  $\frac{dA}{A} = \frac{0.08\pi}{4\pi} = 0.02 = 2\%$

47. Let  $A$  = cross section area,  $C$  = circumference, and

$D$  = diameter. Then  $D = \frac{C}{\pi}$ , so  $\frac{dD}{dC} = \frac{1}{\pi}$

and  $dD = \frac{1}{\pi}dC$ . Also,  $A = \pi\left(\frac{D}{2}\right)^2 = \pi\left(\frac{C}{2\pi}\right)^2 = \frac{C^2}{4\pi}$ ,

so  $\frac{dA}{dC} = \frac{C}{2\pi}$  and  $dA = \frac{C}{2\pi}dC$ . When  $C$  increases from  $10\pi$  in. to  $10\pi + 2$  in. the diameter increases by

$dD = \frac{1}{\pi}(2) = \frac{2}{\pi} \approx 0.6366$  in. and the area increases by

approximately  $dA = \frac{10\pi}{2\pi}(2) = 10$  in<sup>2</sup>.

48. Let  $x$  = edge length and  $V$  = volume. Then  $V = x^3$ , and so  $dV = 3x^2 dx$ . With  $x = 10$  cm and  $dx = 0.01x = 0.1$  cm,

we have  $V = 10^3 = 1000$  cm<sup>3</sup> and

$dV = 3(10)^2(0.1) = 30$  cm<sup>3</sup>, so the percentage error in the volume measurement is approximately

$\frac{dV}{V} = \frac{30}{1000} = 0.03 = 3\%$ .

49. Let  $x$  = side length and  $A$  = area. Then  $A = x^2$  and

$\frac{dA}{dx} = 2x$ , so  $dA = 2x dx$ . We want  $|dA| \leq 0.02A$ , which

gives  $|2x dx| \leq 0.02x^2$ , or  $|dx| \leq 0.01x$ . The side length should be measured with an error of no more than 1%.

For  $\theta = 75^\circ = \frac{5\pi}{12}$  radians, we have

$|\delta\theta| < 0.04 \sin \frac{5\pi}{12} \cos \frac{5\pi}{12} = 0.01$  radian. The angle should be

measured with an error of less than 0.01 radian (or approximately 0.57 degrees), which is a percentage error of approximately 0.76%.

50. (a) Note that  $V = \pi r^2 h = 10\pi r^2 = 2.5\pi D^2$ , where  $D$  is the

interior diameter of the tank. Then  $\frac{dV}{dD} = 5\pi D$ ,

so  $dV = 5\pi D dD$ . We want  $|dV| \leq 0.01V$ , which

gives  $|5\pi D dD| \leq 0.01(2.5\pi D^2)$ , or  $|dD| \leq 0.005D$ . The

interior diameter should be measured with an error of no more than 0.5%.

(b) Now we let  $D$  represent the exterior diameter of the tank, and we assume that the paint coverage rate (number of square feet covered per gallon of paint) is known precisely. Then, to determine the amount of paint within 5%, we need to calculate the lateral surface area  $S$  with an error of no more than 5%. Note that

$S = 2\pi r h = 10\pi D$ , so  $\frac{dS}{dD} = 10\pi$  and  $dS = 10\pi dD$ . We

want  $|dS| \leq 0.05S$ , which gives  $|10\pi dD| \leq 0.05(10\pi D)$ ,

or  $dD \leq 0.05D$ . The exterior diameter should be measured with an error of no more than 5%.

51. Note that  $V = \pi r^2 h$ , where  $h$  is constant. Then  $\frac{dV}{dr} = 2\pi r h$ .

The percent change is given by

$\frac{dV}{V} = \frac{2\pi r h dr}{\pi r^2 h} = 2 \frac{dr}{r} = 2 \frac{0.1\%r}{r} = 0.2\%$ .

52. Note that  $\frac{dV}{dh} = 3\pi h^2$ , so  $dV = 3\pi h^2 dh$ . We want

$|dV| \leq 0.01V$ , which gives  $|3\pi h^2 dh| \leq 0.01(\pi h^3)$ ,

or  $|dh| \leq \frac{0.01h}{3}$ . The height should be measured with an

error of no more than  $\frac{1}{3}\%$ .

53. Since  $V = \frac{4}{3}\pi r^3$ , we have

$dV = 4\pi r^2 dr = 4\pi r^2 \left(\frac{1}{16\pi}\right) = \frac{r^2}{4}$ . The volume error in

each case is simply  $\frac{r^2}{4}$  in<sup>3</sup>.

Sphere Type	True Radius	Tape error	Radius Error	Volume Error
Orange	2 in.	1/8 in.	1/16π in.	1 in. <sup>3</sup>
Melon	4 in.	1/8 in.	1/16π in.	4 in. <sup>3</sup>
Beach Ball	7 in.	1/8 in.	1/16π in.	12.25 in. <sup>3</sup>

54. Since  $A = 4\pi r^2$ , we have  $dA = 8\pi r dr = 8\pi r \left(\frac{1}{16\pi}\right) = \frac{r}{2}$ .

The surface area error in each case is simply  $\frac{r}{2} \text{ in}^2$ .

Sphere Type	True Radius	Tape Error	Radius Error	Volume Error
Orange	2 in.	1/8 in.	1/16π in.	1 in. <sup>2</sup>
Melon	4 in.	1/8 in.	1/16π in.	2 in. <sup>2</sup>
Beach Ball	7 in.	1/8 in.	1/16π in.	3.5 in. <sup>2</sup>

55. We have  $\frac{dW}{dg} = -bg^{-2}$ , so  $dW = -bg^{-2}dg$ .

Then  $\frac{dW_{\text{moon}}}{dW_{\text{earth}}} = \frac{-b(5.2)^{-2}dg}{-b(32)^{-2}dg} = \frac{32^2}{5.2^2} \approx 37.87$ . The ratio is about 37.87 to 1.

56. (a) Note that  $T = 2\pi L^{1/2}g^{-1/2}$ , so  $\frac{dT}{dg} = -\pi L^{1/2}g^{-3/2}$  and

$$dT = -\pi L^{1/2}g^{-3/2}dg.$$

(b) Note that  $dT$  and  $dg$  have opposite signs. Thus, if  $g$  increases,  $T$  decreases and the clock speeds up.

(c) 
$$-\pi L^{1/2}g^{-3/2}dg = dT$$

$$-\pi(100)^{1/2}(980)^{-3/2}dg = 0.001$$

$$dg \approx -0.9765$$

Since  $dg \approx -0.9765$ ,  $g = 980 - 0.9765 = 979.0235$ .

57. True. A look at the graph reveals the problem. The graph decreases after  $x = 1$  toward a horizontal asymptote of  $x = 0$ , so the  $x$ -intercepts of the tangent lines keep getting bigger without approaching a zero.

58. False. By the product rule,  $d(uv) = udv + vdu$ .

59. B.  $f(x) = e^x$   
 $f'(x) = e^x$   
 $L(x) = e^1 + e^1(x-1)$   
 $L(x) = ex$

60. A.  $y = \tan x$   
 $dy = (\sec^2 x)dx = (\sec^2 \pi)0.5$   
 $dy = -0.25$

61. D.  $f(x) = x - x^3 + 2$   
 $f'(x) = 1 - 3x^2$   

$$x_{n+1} = x_n - \frac{x_n x_n^3 + 2}{1 - 3x_n^2}$$

$$x_2 = 1 - \frac{1 - (1)^3 + 2}{1 - 3(1)^2} = 2$$

$$x_3 = 2 - \frac{2 - (2)^3 + 2}{1 - 3(2)^2} = \frac{18}{11}$$

62. A.  $f(x) = \sqrt[3]{x}$ ,  $x = 64$

$$f'(64) = \frac{1}{3}(64)^{-2/3} = \frac{1}{48}$$

$$\sqrt[3]{66} = 4 + \frac{1}{48}(66 - 64)$$

$$\sqrt[3]{66} = 4.042$$

The calculator returns 4.041, or a 0.01% difference.


63. If  $f'(x) \neq 0$ , we have  $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = x_1 - \frac{0}{f'(x_1)} = x_1$ .

Therefore,  $x_2 = x_1$ , and all later approximations are also equal to  $x_1$ .

64. If  $x_1 = h$ , then  $f'(x_1) = \frac{1}{2h^{1/2}}$  and

$$x_2 = h - \frac{h^{1/2}}{\frac{1}{2h^{1/2}}} = h - 2h = -h. \text{ If } x_1 = -h, \text{ then}$$

$$f'(x_1) = -\frac{1}{2\sqrt{h}} \text{ and } x_2 = -h - \frac{h^{1/2}}{-\frac{1}{2\sqrt{h}}} = -h + 2h = h$$




[-3, 3] by [-0.5, 2]

65. Note that  $f'(x) = \frac{1}{3}x^{-2/3}$  and so

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^{1/3}}{\frac{x_n^{-2/3}}{3}} = x_n - 3x_n = -2x_n. \text{ For}$$

$x_1 = 1$ , we have  $x_2 = -2$ ,  $x_3 = 4$ ,  $x_4 = -8$ , and

$x_5 = 16$ ;  $|x_n| = 2^{n-1}$ .



[-10, 10] by [-3, 3]

66. (a) i.  $Q(a) = f(a)$  implies that  $b_0 = f(a)$ .

ii. Since  $Q'(x) = b_1 + 2b_2(x-a)$ ,  $Q'(a) = f'(a)$  implies that  $b_1 = f'(a)$ .



## 66. Continued

iii. Since  $Q''(x) = 2b_2$ ,  $Q''(a) = f''(a)$  implies that

$$b_2 = \frac{f''(a)}{2}$$

In summary,  $b_0 = f(a)$ ,  $b_1 = f'(a)$ , and  $b_2 = \frac{f''(a)}{2}$ .

(b)  $f(x) = (1-x)^{-1}$   
 $f'(x) = -1(1-x)^{-2}(-1) = (1-x)^{-2}$   
 $f''(x) = -2(1-x)^{-3}(-1) = 2(1-x)^{-3}$

Since  $f(0) = 1$ ,  $f'(0) = 1$ , and  $f''(0) = 2$ , the coefficients are

$$b_0 = 1, b_1 = 1, \text{ and } b_2 = \frac{2}{2} = 1. \text{ The quadratic approximation}$$

is  $Q(x) = 1 + x + x^2$ .



$[-2.35, 2.35]$  by  $[-1.25, 3.25]$

As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(d)  $g(x) = x^{-1}$   
 $g'(x) = -x^{-2}$   
 $g''(x) = 2x^{-3}$   
 Since  $g(1) = 1$ ,  $g'(1) = -1$ , and  $g''(1) = 2$ , the coefficients are  $b_0 = 1$ ,  $b_1 = -1$ , and  $b_2 = \frac{2}{2} = 1$ . The

quadratic approximation is  $Q(x) = 1 - (x-1) + (x-1)^2$ .



$[-1.35, 3.35]$  by  $[-1.25, 3.25]$

As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(e)  $h(x) = (1+x)^{1/2}$

$$h'(x) = \frac{1}{2}(1+x)^{-1/2}$$

$$h''(x) = -\frac{1}{4}(1+x)^{-3/2}$$

Since  $h(0) = 1$ ,  $h'(0) = \frac{1}{2}$ , and  $h''(0) = -\frac{1}{4}$ , the

coefficients are  $b_0 = 1$ ,  $b_1 = \frac{1}{2}$ , and  $b_2 = \frac{-\frac{1}{4}}{2} = -\frac{1}{8}$ .

The quadratic approximation is  $Q(x) = 1 + \frac{x}{2} - \frac{x^2}{8}$ .



$[-1.35, 3.35]$  by  $[-1.25, 3.25]$

As one zooms in, the two graphs quickly become indistinguishable. They appear to be identical.

(f) The linearization of any differentiable function  $u(x)$  at  $x = a$  is  $L(x) = u(a) + u'(a)(x-a) = b_0 + b_1(x-a)$ , where  $b_0$  and  $b_1$  are the coefficients of the constant and linear terms of the quadratic approximation. Thus, the linearization for  $f(x)$  at  $x = 0$  is  $1 + x$ ; the linearization for  $g(x)$  at  $x = 1$  is  $1 - (x-1)$  or  $2 - x$ ; and the linearization for  $h(x)$  at  $x = 0$  is  $1 + \frac{x}{2}$ .

67. Finding a zero of  $\sin x$  by Newton's method would use the

recursive formula  $x_{n+1} = x_n - \frac{\sin(x_n)}{\cos(x_n)} = x_n - \tan x_n$ , and that

is exactly what the calculator would be doing. Any zero of  $\sin x$  would be a multiple of  $\pi$ .

68. Just multiply the corresponding derivative formulas by  $dx$ .

(a) Since  $\frac{d}{dx}(c) = 0$ ,  $d(c) = 0$ .

(b) Since  $\frac{d}{dx}(cu) = c \frac{du}{dx}$ ,  $d(cu) = c du$ .

(c) Since  $\frac{d}{dx}(u+v) = \frac{du}{dx} + \frac{dv}{dx}$ ,  $d(u+v) = du + dv$

(d) Since  $\frac{d}{dx}(u \cdot v) = u \frac{dv}{dx} + v \frac{du}{dx}$ ,  $d(u \cdot v) = u dv + v du$ .

(e) Since  $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$ ,  $d\left(\frac{u}{v}\right) = \frac{v du - u dv}{v^2}$ .

(f) Since  $\frac{d}{dx}u^n = nu^{n-1} \frac{du}{dx}$ ,  $d(u^n) = nu^{n-1} du$ .

69.  $\lim_{x \rightarrow 0} \frac{\tan x}{x} = \lim_{x \rightarrow 0} \frac{\sin x / \cos x}{x}$   
 $= \lim_{x \rightarrow 0} \left( \frac{1}{\cos x} \cdot \frac{\sin x}{x} \right)$   
 $= \left( \lim_{x \rightarrow 0} \frac{1}{\cos x} \right) \left( \lim_{x \rightarrow 0} \frac{\sin x}{x} \right)$   
 $= (1)(1) = 1.$

- 70.
- $g(a) = c$
- , so if
- $E(a) = 0$
- , then
- $g(a) = f(a)$
- and
- $c = f(a)$
- .

Then  $E(x) = f(x) - g(x) = f(x) - f(a) - m(x - a)$ .

Thus, 
$$\frac{E(x)}{x-a} = \frac{f(x) - f(a)}{x-a} - m.$$

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x-a} = f'(a), \text{ so } \lim_{x \rightarrow a} \frac{E(x)}{x-a} = f'(a) - m.$$

Therefore, if the limit of  $\frac{E(x)}{x-a}$  is zero, then  $m = f'(a)$  and  $g(x) = L(x)$ .

71. 
$$f'(x) = \frac{1}{2\sqrt{x+1}} + \cos x$$

We have  $f(0) = 1$  and  $f'(0) = \frac{3}{2}$

$$\begin{aligned} L(x) &= f(0) + f'(0)(x-0) \\ &= 1 + \frac{3}{2}x \end{aligned}$$


The linearization is the sum of the two individual

linearizations, which are  $x$  for  $\sin x$  and  $1 + \frac{1}{2}x$  for  $\sqrt{x+1}$ .**Section 4.6 Related Rates (pp. 246–255)****Exploration 1 Sliding Ladder**

1. Here the  $x$ -axis represents the ground and the  $y$ -axis represents the wall. The curve  $(x_1, y_1)$  gives the position of the bottom of the ladder (distance from the wall) at any time  $t$  in  $0 \leq t \leq 5$ . The curve  $(x_2, y_2)$  gives the position of the top of the ladder at any time in  $0 \leq t \leq 5$ .

2.  $0 \leq t \leq 5$

4. This is a snapshot at
- $t \approx 3$
- . 1. The top of the ladder is moving down the
- $y$
- axis and the bottom of the ladder is moving to the right on the
- $x$
- axis. The end of the ladder is accelerating. Both axes are hidden from view.



[-1, 15] by [-1, 15]

6. 
$$\frac{dy}{dt} = \frac{-4T}{\sqrt{10^2 - (2T)^2}}$$

- 7.
- $y'(3) \approx -4.24 \text{ ft/sec}^2$
- . The negative number means the ladder is falling.

8. Since
- $\lim_{t \rightarrow (13/3)^-} y'(t) = -\infty$
- , the speed of the top of the ladder is infinite as it hits the ground.

**Quick Review 4.6**

1.  $D = \sqrt{(7-0)^2 + (0-5)^2} = \sqrt{49+25} = \sqrt{74}$

2.  $D = \sqrt{(b-0)^2 + (0-a)^2} = \sqrt{a^2 + b^2}$

3. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx}(2xy + y^2) &= \frac{d}{dx}(x+y) \\ 2x \frac{dy}{dx} + 2y(1) + 2y \frac{dy}{dx} &= (1) + \frac{dy}{dx} \\ (2x + 2y - 1) \frac{dy}{dx} &= 1 - 2y \\ \frac{dy}{dx} &= \frac{1 - 2y}{2x + 2y - 1} \end{aligned}$$

4. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx}(x \sin y) &= \frac{d}{dx}(1 - xy) \\ (x)(\cos y) \frac{dy}{dx} + (\sin y)(1) &= -x \frac{dy}{dx} - y(1) \\ (x + x \cos y) \frac{dy}{dx} &= -y - \sin y \\ \frac{dy}{dx} &= \frac{-y - \sin y}{x + x \cos y} \\ \frac{dy}{dx} &= -\frac{y + \sin y}{x + x \cos y} \end{aligned}$$

5. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx} x^2 &= \frac{d}{dx} \tan y \\ 2x &= \sec^2 y \frac{dy}{dx} \\ \frac{dy}{dx} &= \frac{2x}{\sec^2 y} \\ \frac{dy}{dx} &= 2x \cos^2 y \end{aligned}$$

6. Use implicit differentiation.

$$\begin{aligned} \frac{d}{dx} \ln(x+y) &= \frac{d}{dx}(2x) \\ \frac{1}{x+y} \left( 1 + \frac{dy}{dx} \right) &= 2 \\ 1 + \frac{dy}{dx} &= 2(x+y) \\ \frac{dy}{dx} &= 2x + 2y - 1 \end{aligned}$$

7. Using
- $A(-2, 1)$
- we create the parametric equations
- $x = -2 + at$
- and
- $y = 1 + bt$
- , which determine a line passing through
- $A$
- at
- $t = 0$
- . We determine
- $a$
- and
- $b$
- so that the line passes through
- $B(4, -3)$
- at
- $t = 1$
- . Since
- $4 = -2 + a$
- , we have
- $a = 6$
- , and since
- $-3 = 1 + b$
- , we have
- $b = -4$
- . Thus, one parametrization for the line segment is
- $x = -2 + 6t$
- ,
- $y = 1 - 4t$
- ,
- $0 \leq t \leq 1$
- . (Other answers are possible.)

8. Using  $A(0, -4)$ , we create the parametric equations  $x = 0 + at$  and  $y = -4 + bt$ , which determine a line passing through  $A$  at  $t = 0$ . We now determine  $a$  and  $b$  so that the line passes through  $B(5, 0)$  at  $t = 1$ . Since  $5 = 0 + a$ , we have  $a = 5$ , and since  $0 = -4 + b$ , we have  $b = 4$ . Thus, one parametrization for the line segment is  $x = 5t$ ,  $y = -4 + 4t$ ,  $0 \leq t \leq 1$ . (Other answers are possible.)

9. One possible answer:  $\frac{\pi}{2} \leq t \leq \frac{3\pi}{2}$

10. One possible answer:  $\frac{3\pi}{2} \leq t \leq 2\pi$

### Section 4.6 Exercises

1. Since  $\frac{dA}{dt} = \frac{dA}{dr} \frac{dr}{dt}$ , we have  $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$ ,

2. Since  $\frac{dS}{dt} = \frac{dS}{dr} \frac{dr}{dt}$ , we have  $\frac{dS}{dt} = 8\pi r \frac{dr}{dt}$ .

3. (a) Since  $\frac{dV}{dt} = \frac{dV}{dh} \frac{dh}{dt}$ , we have  $\frac{dV}{dt} = \pi r^2 \frac{dh}{dt}$ .

(b) Since  $\frac{dV}{dt} = \frac{dV}{dr} \frac{dr}{dt}$ , we have  $\frac{dV}{dt} = 2\pi r h \frac{dr}{dt}$ .

(c)  $\frac{dV}{dt} = \frac{d}{dt} \pi r^2 h = \pi \frac{d}{dt} (r^2 h)$

$$\frac{dV}{dt} = \pi \left( r^2 \frac{dh}{dt} + h(2r) \frac{dr}{dt} \right)$$

$$\frac{dV}{dt} = \pi r^2 \frac{dh}{dt} + 2\pi r h \frac{dr}{dt}$$

4. (a)  $\frac{dP}{dt} = \frac{d}{dt} (RI^2)$

$$\frac{dP}{dt} = R \frac{d}{dt} I^2 + I^2 \frac{dR}{dt}$$

$$\frac{dP}{dt} = R \left( 2I \frac{dI}{dt} \right) + I^2 \frac{dR}{dt}$$

$$\frac{dP}{dt} = 2RI \frac{dI}{dt} + I^2 \frac{dR}{dt}$$

(b) If  $P$  is constant, we have  $\frac{dP}{dt} = 0$ , which means

$$2RI \frac{dI}{dt} + I^2 \frac{dR}{dt} = 0, \text{ or } \frac{dR}{dt} = -\frac{2R}{I} \frac{dI}{dt} = -\frac{2P}{I^3} \frac{dI}{dt}$$

5.  $\frac{ds}{dt} = \frac{d}{dt} \sqrt{x^2 + y^2 + z^2}$

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2 + z^2}} \frac{d}{dt} (x^2 + y^2 + z^2)$$

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2 + z^2}} \left( 2x \frac{dx}{dt} + 2y \frac{dy}{dt} + 2z \frac{dz}{dt} \right)$$

$$\frac{ds}{dt} = \frac{x \frac{dx}{dt} + y \frac{dy}{dt} + z \frac{dz}{dt}}{\sqrt{x^2 + y^2 + z^2}}$$

6.  $\frac{dA}{dt} = \frac{d}{dt} \left( \frac{1}{2} ab \sin \theta \right)$

$$\frac{dA}{dt} = \frac{1}{2} \left( \frac{da}{dt} \cdot b \cdot \sin \theta + a \cdot \frac{db}{dt} \cdot \sin \theta + ab \cdot \frac{d}{dt} \sin \theta \right)$$

$$\frac{dA}{dt} = \frac{1}{2} \left( b \sin \theta \frac{da}{dt} + a \sin \theta \frac{db}{dt} + ab \cos \theta \frac{d\theta}{dt} \right)$$

7. (a) Since  $V$  is increasing at the rate of 1 volt/sec,

$$\frac{dV}{dt} = 1 \text{ volt/sec.}$$

(b) Since  $I$  is decreasing at the rate of

$$\frac{1}{3} \text{ amp/sec, } \frac{dI}{dt} = -\frac{1}{3} \text{ amp/sec.}$$

(c) Differentiating both sides of  $V = IR$ , we have

$$\frac{dV}{dt} = I \frac{dR}{dt} + R \frac{dI}{dt}$$

(d) Note that  $V = IR$  gives  $12 = 2R$ , so  $R = 6$  ohms. Now substitute the known values into the equation in (c).

$$1 = 2 \frac{dR}{dt} + 6 \left( -\frac{1}{3} \right)$$

$$3 = 2 \frac{dR}{dt}$$

$$\frac{dR}{dt} = \frac{3}{2} \text{ ohms/sec}$$

$R$  is changing at the rate of  $\frac{3}{2}$  ohms/sec. Since this value is positive,  $R$  is increasing.

8. Step 1:

$r$  = radius of plate

$A$  = area of plate

Step 2:

At the instant in question,  $\frac{dr}{dt} = 0.01$  cm/sec,  $r = 50$  cm.

Step 3:

We want to find  $\frac{dA}{dt}$ .

Step 4:

$$A = \pi r^2$$

Step 5:

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

Step 6:

$$\frac{dA}{dt} = 2\pi(50)(0.01) = \pi \text{ cm}^2 / \text{sec}$$

At the instant in question, the area is increasing at the rate of  $\pi \text{ cm}^2 / \text{sec}$ .

## 9. Step 1:

 $l$  = length of rectangle $w$  = width of rectangle $A$  = area of rectangle $P$  = perimeter of rectangle $D$  = length of a diagonal of the rectangle

## Step 2:

At the instant in question,

$$\frac{dl}{dt} = -2 \text{ cm/sec}, \quad \frac{dw}{dt} = 2 \text{ cm/sec}, \quad l = 12 \text{ cm}, \quad \text{and } w = 5 \text{ cm}.$$

## Step 3:

We want to find  $\frac{dA}{dt}$ ,  $\frac{dP}{dt}$ , and  $\frac{dD}{dt}$ .

Steps 4, 5, and 6:

(a)  $A = lw$

$$\frac{dA}{dt} = l \frac{dw}{dt} + w \frac{dl}{dt}$$

$$\frac{dA}{dt} = (12)(2) + (5)(-2) = 14 \text{ cm}^2/\text{sec}$$

The rate of change of the area is  $14 \text{ cm}^2/\text{sec}$ .

(b)  $P = 2l + 2w$

$$\frac{dP}{dt} = 2 \frac{dl}{dt} + 2 \frac{dw}{dt}$$

$$\frac{dP}{dt} = 2(-2) + 2(2) = 0 \text{ cm/sec}$$

The rate of change of the perimeter is  $0 \text{ cm/sec}$ .

(c)  $D = \sqrt{l^2 + w^2}$

$$\frac{dD}{dt} = \frac{1}{2\sqrt{l^2 + w^2}} \left( 2l \frac{dl}{dt} + 2w \frac{dw}{dt} \right) = \frac{l \frac{dl}{dt} + w \frac{dw}{dt}}{\sqrt{l^2 + w^2}}$$

$$\frac{dD}{dt} = \frac{(12)(-2) + (5)(2)}{\sqrt{12^2 + 5^2}} = -\frac{14}{13} \text{ cm/sec}$$

The rate of change of the length of the diameter is

$$-\frac{14}{13} \text{ cm/sec}.$$

(d) The area is increasing, because its derivative is positive.

The perimeter is not changing, because its derivative is zero. The diagonal length is decreasing, because its derivative is negative.

## 10. Step 1:

 $x, y, z$  = edge lengths of the box $V$  = volume of the box $S$  = surface area of the box $s$  = diagonal length of the box

## Step 2:

At the instant in question,

$$\frac{dx}{dt} = 1 \text{ m/sec}, \quad \frac{dy}{dt} = -2 \text{ m/sec}, \quad \frac{dz}{dt} = 1 \text{ m/sec}, \quad x = 4 \text{ m}, \\ y = 3 \text{ m}, \quad \text{and } z = 2 \text{ m}.$$

## Step 3:

We want to find  $\frac{dV}{dt}$ ,  $\frac{dS}{dt}$ , and  $\frac{ds}{dt}$ .

Steps 4, 5, and 6:

(a)  $V = xyz$

$$\frac{dV}{dt} = xy \frac{dz}{dt} + xz \frac{dy}{dt} + yz \frac{dx}{dt}$$

$$\frac{dV}{dt} = (4)(3)(1) + (4)(2)(-2) + (3)(2)(1) = 2 \text{ m}^3/\text{sec}$$

The rate of change of the volume is  $2 \text{ m}^3/\text{sec}$ .

(b)  $S = 2(xy + xz + yz)$

$$\frac{dS}{dt} = 2 \left( x \frac{dy}{dt} + y \frac{dx}{dt} + x \frac{dz}{dt} + z \frac{dx}{dt} + y \frac{dz}{dt} + z \frac{dy}{dt} \right)$$

$$\frac{dS}{dt} = 2[(4)(-2) + (3)(1) + (4)(1) + (2)(1) \\ + (2)(1) + (3)(1) + (2)(-2)] = 0 \text{ m}^2/\text{sec}$$

The rate of change of the surface area is  $0 \text{ m}^2/\text{sec}$ .

(c)  $s = \sqrt{x^2 + y^2 + z^2}$

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2 + z^2}} \left( 2x \frac{dx}{dt} + 2y \frac{dy}{dt} + 2z \frac{dz}{dt} \right)$$

$$= \frac{x \frac{dx}{dt} + y \frac{dy}{dt} + z \frac{dz}{dt}}{\sqrt{x^2 + y^2 + z^2}}$$

$$\frac{ds}{dt} = \frac{(4)(1) + (3)(-2) + (2)(1)}{\sqrt{4^2 + 3^2 + 2^2}} = \frac{0}{\sqrt{29}} = 0 \text{ m/sec}$$

The rate of change of the diagonal length is  $0 \text{ m/sec}$ .

## 11. Step 1:

 $r$  = radius of spherical balloon $S$  = surface area of spherical balloon $V$  = volume of spherical balloon

## Step 2:

At the instant in question,  $\frac{dV}{dt} = 100\pi \text{ ft}^3/\text{min}$  and  $r = 5 \text{ ft}$ .

## Step 3:

We want to find the values of  $\frac{dr}{dt}$  and  $\frac{dS}{dt}$ .

Steps 4, 5, and 6:

(a)  $V = \frac{4}{3}\pi r^3$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$100\pi = 4\pi(5)^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = 1 \text{ ft/min}$$

The radius is increasing at the rate of  $1 \text{ ft/min}$ .

## 11. Continued

(b)  $S = 4\pi r^2$

$$\frac{dS}{dt} = 8\pi r \frac{dr}{dt}$$

$$\frac{dS}{dt} = 8\pi(5)(1)$$

$$\frac{dS}{dt} = 40\pi \text{ ft}^2 / \text{min}$$

The surface area is increasing at the rate of  $40\pi$   $\text{ft}^2/\text{min}$ .

## 12. Step 1:

$r$  = radius of spherical droplet

$S$  = surface area of spherical droplet

$V$  = volume of spherical droplet

## Step 2:

No numerical information is given.

## Step 3:

We want to show that  $\frac{dr}{dt}$  is constant.

## Step 4:

$$S = 4\pi r^2, V = \frac{4}{3}\pi r^3, \frac{dV}{dt} = kS \text{ for some constant } k$$

Steps 5 and 6:

$$\text{Differentiating } V = \frac{4}{3}\pi r^3, \text{ we have } \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}.$$

Substituting  $kS$  for  $\frac{dV}{dt}$  and  $S$  for  $4\pi r^2$ , we

$$\text{have } kS = S \frac{dr}{dt}, \text{ or } S \frac{dr}{dt} = k.$$

## 13. Step 1:

$s$  = (diagonal) distance from antenna to airplane

$x$  = horizontal distance from antenna to airplane

## Step 2:

At the instant in question,

$$s = 10 \text{ mi and } \frac{ds}{dt} = 300 \text{ mph.}$$

## Step 3:

We want to find  $\frac{dx}{dt}$ .

## Step 4:

$$x^2 + 49 = s^2 \text{ or } x = \sqrt{s^2 - 49}$$

## Step 5:


$$\frac{dx}{dt} = \frac{1}{2\sqrt{s^2 - 49}} \left( 2s \frac{ds}{dt} \right) = \frac{s}{\sqrt{s^2 - 49}} \frac{ds}{dt}$$

## Step 6:

$$\frac{dx}{dt} = \frac{10}{\sqrt{10^2 - 49}} (300) = \frac{3000}{\sqrt{51}} \text{ mph} \approx 420.08 \text{ mph}$$

The speed of the airplane is about 420.08 mph.

## 14. Step 1:



$s$  = length of kite string

$x$  = horizontal distance from Inge to kite

## Step 2:

At the instant in question,  $\frac{dx}{dt} = 25$  ft/sec and  $s = 500$  ft

## Step 3:

We want to find  $\frac{ds}{dt}$ .

## Step 4:

$$x^2 + 300^2 = s^2$$

## Step 5:

$$2x \frac{dx}{dt} = 2s \frac{ds}{dt} \text{ or } x \frac{dx}{dt} = s \frac{ds}{dt}$$


## Step 6:

At the instant in question, since  $x^2 + 300^2 = s^2$ , we have

$$x = \sqrt{s^2 - 300^2} = \sqrt{500^2 - 300^2} = 400.$$

Thus  $(400)(25) = (500) \frac{ds}{dt}$ , so  $\frac{ds}{dt} = 20$  ft/sec. Inge must let the string out at the rate of 20 ft/sec.

## 15. Step 1:



The cylinder shown represents the shape of the hole.

$r$  = radius of cylinder

$V$  = volume of cylinder

## Step 2:

$$\text{At the instant in question, } \frac{dr}{dt} = \frac{0.001 \text{ in.}}{3 \text{ min}} = \frac{1}{3000} \text{ in./min}$$

and (since the diameter is 3.800 in.),  $r = 1.900$  in.

## Step 3:

We want to find  $\frac{dV}{dt}$ .

## Step 4:

$$V = \pi r^2(6) = 6\pi r^2$$

## Step 5:

$$\frac{dV}{dt} = 12\pi r \frac{dr}{dt}$$

15. Continued


Step 6:

$$\frac{dV}{dt} = 12\pi(1.900)\left(\frac{1}{3000}\right) = \frac{19\pi}{2500} = 0.0076\pi$$

$$\approx 0.0239 \text{ in}^3/\text{min}.$$

The volume is increasing at the rate of approximately  $0.0239 \text{ in}^3/\text{min}$ .

16. Step 1:



$r$  = base radius of cone  
 $h$  = height of cone  
 $V$  = volume of cone

Step 2:

At the instant in question,  $h = 4 \text{ m}$  and  $\frac{dV}{dt} = 10 \text{ m}^3/\text{min}$ .

Step 3:

We want to find  $\frac{dh}{dt}$  and  $\frac{dr}{dt}$ .

Step 4:

Since the height is  $\frac{3}{8}$  of the base diameter, we have

$$h = \frac{3}{8}(2r) \text{ or } r = \frac{4}{3}h.$$

We also have  $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi\left(\frac{4}{3}h\right)^2 h = \frac{16\pi h^3}{27}$ . We will

use the equations  $V = \frac{16\pi h^3}{27}$  and  $r = \frac{4}{3}h$ .

Step 5 and 6:

(a) 
$$\frac{dV}{dt} = \frac{16\pi h^2}{9} \frac{dh}{dt}$$

$$10 = \frac{16\pi(4)^2}{9} \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{45}{128\pi} \text{ m/min} = \frac{1125}{32\pi} \text{ cm/min}$$

The height is changing at the rate of

$$\frac{1125}{32\pi} \approx 11.19 \text{ cm/min}.$$


(b) Using the results from Step 4 and part (a), we have

$$\frac{dr}{dt} = \frac{4}{3} \frac{dh}{dt} = \frac{4}{3} \left( \frac{1125}{32\pi} \right) = \frac{375}{8\pi} \text{ cm/min}.$$

The radius is changing at the rate of

$$\frac{375}{8\pi} \approx 14.92 \text{ cm/min}.$$

17. Step 1:



$r$  = radius of top surface of water  
 $h$  = depth of water in reservoir  
 $V$  = volume of water in reservoir

Step 2:

At the instant in question,  $\frac{dV}{dt} = -50 \text{ m}^3/\text{min}$  and  $h = 5 \text{ m}$ .

Step 3:

We want to find  $-\frac{dh}{dt}$  and  $\frac{dr}{dt}$ .

Step 4:

Note that  $\frac{h}{r} = \frac{6}{45}$  by similar cones, so  $r = 7.5h$ .

Then  $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi(7.5h)^2 h = 18.75\pi h^3$

Steps 5 and 6:

(a) Since  $V = 18.75\pi h^3$ ,  $\frac{dV}{dt} = 56.25\pi h^2 \frac{dh}{dt}$ .

Thus  $-50 = 56.25\pi(5^2) \frac{dh}{dt}$ , and

$$\text{so } \frac{dh}{dt} = -\frac{8}{225\pi} \text{ m/min} = -\frac{32}{9\pi} \text{ cm/min}.$$

The water level is falling by  $\frac{32}{9\pi} \approx 1.13 \text{ cm/min}$ .

(Since  $\frac{dh}{dt} < 0$ , the rate at which the water level is falling is positive.)

(b) Since  $r = 7.5h$ ,  $\frac{dr}{dt} = 7.5 \frac{dh}{dt} = -\frac{80}{3\pi} \text{ cm/min}$ . The rate of

change of the radius of the water's surface is

$$-\frac{80}{3\pi} \approx -8.49 \text{ cm/min}.$$

18. (a) Step 1:

$y$  = depth of water in bowl  
 $V$  = volume of water in bowl

Step 2:

At the instant in question,  $\frac{dV}{dt} = -6 \text{ m}^3/\text{min}$  and

$y = 8 \text{ m}$ .

Step 3:

We want to find the value of  $\frac{dy}{dt}$ .

Step 4:

$$V = \frac{\pi}{3} y^2 (39 - y) \text{ or } V = 13\pi y^2 - \frac{\pi}{3} y^3$$

## 18. Continued

Step 5:

$$\frac{dV}{dt} = (26\pi y - \pi y^2) \frac{dy}{dt}$$

Step 6:

$$-6 = [26\pi(8) - \pi(8^2)] \frac{dy}{dt}$$

$$-6 = 144\pi \frac{dy}{dt}$$

$$\frac{dy}{dt} = -\frac{1}{24\pi} \approx -0.01326 \text{ m/min}$$

$$\text{or } -\frac{25}{6\pi} \approx -1.326 \text{ cm/min}$$

(b) Since  $r^2 + (13 - y)^2 = 13^2$ ,

$$r = \sqrt{169 - (13 - y)^2} = \sqrt{26y - y^2}.$$

(c) Step 1:

 $y$  = depth of water $r$  = radius of water surface $V$  = volume of water in bowl

Step 2:

At the instant in question,  $\frac{dV}{dt} = -6 \text{ m}^3/\text{min}$ ,  $y = 8 \text{ m}$ ,and therefore (from part (a))  $\frac{dy}{dt} = -\frac{1}{24\pi} \text{ m/min}$ .

Step 3:

We want to find the value of  $\frac{dr}{dt}$ .

Step 4:

From part (b),  $r = \sqrt{26y - y^2}$ .

Step 5:

$$\frac{dr}{dt} = \frac{1}{2\sqrt{26y - y^2}} (26 - 2y) \frac{dy}{dt} = \frac{13 - y}{\sqrt{26y - y^2}} \frac{dy}{dt}$$

Step 6:

$$\frac{dr}{dt} = \frac{13 - 8}{\sqrt{26(8) - 8^2}} \left( -\frac{1}{24\pi} \right) = \frac{5}{12} \left( -\frac{1}{24\pi} \right)$$

$$= -\frac{5}{288\pi} \approx -0.00553 \text{ m/min}$$

$$\text{or } -\frac{125}{72\pi} \approx -0.553 \text{ cm/min}$$

## 19. Step 1:

 $x$  = distance from wall to base of ladder $y$  = height of top of ladder $A$  = area of triangle formed by the ladder, wall, and ground $\theta$  = angle between the ladder and the ground

Step 2:

At the instant in question,  $x = 12 \text{ ft}$  and  $\frac{dx}{dt} = 5 \text{ ft/sec}$ .

Step 3:

We want to find  $-\frac{dy}{dt}$ ,  $\frac{dA}{dt}$ , and  $\frac{d\theta}{dt}$ .

Steps 4, 5, and 6:

(a)  $x^2 + y^2 = 169$ 

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

To evaluate, note that, at the instant in question,

$$y = \sqrt{169 - x^2} = \sqrt{169 - 12^2} = 5.$$

$$\text{Then } 2(12)(5) + 2(5) \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = -12 \text{ ft/sec} \left( \text{or } -\frac{dy}{dt} = 12 \text{ ft/sec} \right)$$

The top of the ladder is sliding down the wall at the rate of 12 ft/sec. (Note that the *downward* rate of motion is positive.)(b)  $A = \frac{1}{2}xy$ 

$$\frac{dA}{dt} = \frac{1}{2} \left( x \frac{dy}{dt} + y \frac{dx}{dt} \right)$$

Using the results from step 2 and from part (a), we have

$$\frac{dA}{dt} = \frac{1}{2} [(12)(-12) + (5)(5)] = -\frac{119}{2} \text{ ft}^2/\text{sec}.$$

The area of the triangle is changing at the rate of  $-59.5 \text{ ft}^2/\text{sec}$ .(c)  $\tan \theta = \frac{y}{x}$ 

$$\sec^2 \theta \frac{d\theta}{dt} = \frac{x \frac{dy}{dt} - y \frac{dx}{dt}}{x^2}$$

Since  $\tan \theta = \frac{5}{12}$ , we have

$$\left( \text{for } 0 \leq \theta < \frac{\pi}{2} \right) \cos \theta = \frac{12}{13} \text{ and so } \sec^2 \theta \frac{1}{\left( \frac{12}{13} \right)^2} = \frac{169}{144}.$$

Combining this result with the results from step 2 and from part (a), we have  $\frac{169}{144} \frac{d\theta}{dt} = \frac{(12)(-12) - (5)(5)}{12^2}$ , so
$$\frac{d\theta}{dt} = -1 \text{ radian/sec}.$$

The angle is changing at the rate of  $-1 \text{ radian/sec}$ .

## 20. Step 1:

 $h$  = height (or depth) of the water in the trough $V$  = volume of water in the trough

Step 2:

At the instant in question,  $\frac{dV}{dt} = 2.5 \text{ ft}^3/\text{min}$  and  $h = 2 \text{ ft}$ .

## 20. Continued

Step 3:

We want to find  $\frac{dh}{dt}$ .

Step 4:

The width of the top surface of the water is  $\frac{4}{3}h$ , so wehave  $V = \frac{1}{2}(h)\left(\frac{4}{3}h\right)(15)$ , or  $V = 10h^2$ 

Step 5:

$$\frac{dV}{dt} = 20h \frac{dh}{dt}$$

Step 6:

$$2.5 = 20(2) \frac{dh}{dt}$$

$$\frac{dh}{dt} = 0.0625 = \frac{1}{16} \text{ ft/min}$$

The water level is increasing at the rate of  $\frac{1}{16}$  ft/min.

## 21. Step 1:

 $l$  = length of rope $x$  = horizontal distance from boat to dock $\theta$  = angle between the rope and a vertical line

Step 2:

At the instant in question,  $\frac{dl}{dt} = -2$  ft/sec and  $l = 10$  ft.

Step 3:

We want to find the values of  $-\frac{dx}{dt}$  and  $\frac{d\theta}{dt}$ .

Steps 4, 5, and 6:

(a)  $x = \sqrt{l^2 - 36}$

$$\frac{dx}{dt} = \frac{l}{\sqrt{l^2 - 36}} \frac{dl}{dt}$$

$$\frac{dx}{dt} = \frac{10}{\sqrt{10^2 - 36}} (-2) = -2.5 \text{ ft/sec}$$

The boat is approaching the dock at the rate of 2.5 ft/sec.

(b)  $\theta = \cos^{-1} \frac{6}{l}$

$$\frac{d\theta}{dt} = -\frac{1}{\sqrt{1 - \left(\frac{6}{l}\right)^2}} \left(-\frac{6}{l^2}\right) \frac{dl}{dt}$$

$$\frac{d\theta}{dt} = -\frac{1}{\sqrt{1 - 0.6^2}} \left(-\frac{6}{10^2}\right) (-2) = -\frac{3}{20} \text{ radian/sec}$$

The rate of change of angle  $\theta$  is  $-\frac{3}{20}$  radian/sec.

## 22. Step 1:

 $x$  = distance from origin to bicycle $y$  = height of balloon (distance from origin to balloon) $s$  = distance from balloon to bicycle

Step 2:

We know that  $\frac{dy}{dt}$  is a constant 1 ft/sec and  $\frac{dx}{dt}$  is aconstant 17 ft/sec. Three seconds before the instant in question, the values of  $x$  and  $y$  are  $x = 0$  ft and  $y = 65$  ft.Therefore, at the instant in question  $x = 51$  ft and  $y = 68$  ft.

Step 3:

We want to find the value of  $\frac{ds}{dt}$  at the instant in question.

Step 4:

$$s = \sqrt{x^2 + y^2}$$

Step 5:

$$\frac{ds}{dt} = \frac{1}{2\sqrt{x^2 + y^2}} \left( 2x \frac{dx}{dt} + 2y \frac{dy}{dt} \right) = \frac{x \frac{dx}{dt} + y \frac{dy}{dt}}{\sqrt{x^2 + y^2}}$$

Step 6:

$$\frac{ds}{dt} = \frac{(51)(17) + (68)(1)}{\sqrt{51^2 + 68^2}} = 11 \text{ ft/sec}$$

The distance between the balloon and the bicycle is increasing at the rate of 11 ft/sec.

23.  $\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt} = -10(1+x^2)^{-2} (2x) \frac{dx}{dt} = -\frac{20x}{(1+x^2)} \frac{dx}{dt}$

Since  $\frac{dx}{dt} = 3$  cm/sec, we have

$$\frac{dy}{dt} = -\frac{60x}{(1+x^2)^2} \text{ cm/sec.}$$

(a)  $\frac{dy}{dt} = -\frac{60(-2)}{[1+(-2)^2]^2} = \frac{120}{5^2} = \frac{24}{5} \text{ cm/sec}$

(b)  $\frac{dy}{dt} = -\frac{60(0)}{(1+0^2)^2} = 0 \text{ cm/sec}$

(c)  $\frac{dy}{dt} = -\frac{60(20)}{(1+20^2)^2} \approx -0.00746 \text{ cm/sec}$

24.  $\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt} = (3x^2 - 4) \frac{dx}{dt}$

Since  $\frac{dx}{dt} = -2$  cm/sec, we have  $\frac{dy}{dt} = 8 - 6x^2$  cm/sec.


(a)  $\frac{dy}{dt} = 8 - 6(-3)^2 = -46 \text{ cm/sec}$

(b)  $\frac{dy}{dt} = 8 - 6(1)^2 = 2 \text{ cm/sec}$

(c)  $\frac{dy}{dt} = 8 - 6(4)^2 = -88 \text{ cm/sec}$



25. Step 1:



$x$  =  $x$ -coordinate of particle's location  
 $y$  =  $y$ -coordinate of particle's location  
 $\theta$  = angle of inclination of line joining the particle to the origin.

Step 2:

At the instant in question,

$$\frac{dx}{dt} = 10 \text{ m/sec and } x = 3 \text{ m.}$$

Step 3:

We want to find  $\frac{d\theta}{dt}$ .

Step 4:

Since  $y = x^2$ , we have  $\tan \theta = \frac{y}{x} = \frac{x^2}{x} = x$  and so,

for  $x > 0$ ,

$$\theta = \tan^{-1} x.$$

Step 5:


$$\frac{d\theta}{dt} = \frac{1}{1+x^2} \frac{dx}{dt}$$

Step 6:

$$\frac{d\theta}{dt} = \frac{1}{1+3^2} (10) = 1 \text{ radian/sec}$$

The angle of inclination is increasing at the rate of 1 radian/sec.

26. Step 1:



$x$  =  $x$ -coordinate of particle's location  
 $y$  =  $y$ -coordinate of particle's location  
 $\theta$  = angle of inclination of line joining the particle to the origin

Step 2:

At the instant in question,  $\frac{dx}{dt} = -8 \text{ m/sec}$  and  $x = -4 \text{ m}$ .

Step 3:

We want to find  $\frac{d\theta}{dt}$ ,

Step 4:

Since  $y = \sqrt{-x}$ , we have  $\tan \theta = \frac{y}{x} = \frac{\sqrt{-x}}{x} = (-x)^{-1/2}$ ,

and so, for  $x < 0$ ,

$$\theta = \pi + \tan^{-1} [(-x)^{1/2}] = \pi - \tan^{-1} (-x)^{-1/2}.$$

Step 5:

$$\begin{aligned} \frac{d\theta}{dt} &= -\frac{1}{1+[(x)^{-1/2}]^2} \left( -\frac{1}{2}(-x)^{-3/2}(-1) \right) \frac{dx}{dt} \\ &= -\frac{1}{1-\left(\frac{1}{x}\right)} \frac{1}{2(-x)^{3/2}} \frac{dx}{dt} \\ &= \frac{1}{2\sqrt{-x}(x-1)} \frac{dx}{dt} \end{aligned}$$

Step 6:

$$\frac{d\theta}{dt} = \frac{1}{2\sqrt{4}(-4-1)} (-8) = \frac{2}{5} \text{ radian/sec}$$

The angle of inclination is increasing at the rate of

$$\frac{2}{5} \text{ radian/sec.}$$

27. Step 1:

 $r$  = radius of balls plus ice $S$  = surface area of ball plus ice $V$  = volume of ball plus ice

Step 2:

At the instant in question,

$$\frac{dV}{dt} = -8 \text{ mL/min} = -8 \text{ cm}^3/\text{min} \text{ and } r = \frac{1}{2}(20) = 10 \text{ cm.}$$

Step 3:

We want to find  $-\frac{dS}{dt}$ .

Step 4:

We have  $V = \frac{4}{3}\pi r^3$  and  $S = 4\pi r^2$ . These equations can be

combined by noting that  $r = \left(\frac{3V}{4\pi}\right)^{1/3}$ , so  $S = 4\pi \left(\frac{3V}{4\pi}\right)^{2/3}$

Step 5:

$$\frac{dS}{dt} = 4\pi \left(\frac{2}{3}\right) \left(\frac{3V}{4\pi}\right)^{-1/3} \left(\frac{3}{4\pi}\right) \frac{dV}{dt} = 2 \left(\frac{3V}{4\pi}\right)^{-1/3} \frac{dV}{dt}$$

Step 6:

$$\text{Note that } V = \frac{4}{3}\pi(10)^3 = \frac{4000\pi}{3}.$$

$$\frac{dS}{dt} = 2 \left(\frac{3}{4\pi} \cdot \frac{4000\pi}{3}\right)^{-1/3} (-8) = \frac{-16}{\sqrt[3]{1000}} = -1.6 \text{ cm}^2/\text{min}$$

Since  $\frac{dS}{dt} < 0$ , the rate of decrease is positive. The surface area is decreasing at the rate of  $1.6 \text{ cm}^2/\text{min}$ .

## 28. Step 1:

 $x$  =  $x$ -coordinate of particle $y$  =  $y$ -coordinate of particle $D$  = distance from origin to particle

## Step 2:

At the instant in question,  $x = 5$  m,  $y = 12$  m,

$$\frac{dx}{dt} = -1 \text{ m/sec, and } \frac{dy}{dt} = -5 \text{ m/sec.}$$

## Step 3:

We want to find  $\frac{dD}{dt}$ .

## Step 4:

$$D = \sqrt{x^2 + y^2}$$

## Step 5:


$$\frac{dD}{dt} = \frac{1}{2\sqrt{x^2 + y^2}} \left( 2x \frac{dx}{dt} + 2y \frac{dy}{dt} \right) = \frac{x \frac{dx}{dt} + y \frac{dy}{dt}}{\sqrt{x^2 + y^2}}$$

## Step 6:

$$\frac{dD}{dt} = \frac{(5)(-1) + (12)(-5)}{\sqrt{5^2 + 12^2}} = -5 \text{ m/sec}$$

The particle's distance from the origin is changing at the rate of  $-5$  m/sec.

## 29. Step 1:

 $x$  = distance from streetlight base to man $s$  = length of shadow

## Step 2:

At the instant in question,  $\frac{dx}{dt} = -5$  ft/sec and  $x = 10$  ft.

## Step 3:

We want to find  $\frac{ds}{dt}$ .

## Step 4:

By similar triangles,  $\frac{s}{6} = \frac{s+x}{16}$ . This is equivalent to

$$16s = 6s + 6x, \text{ or } s = \frac{3}{5}x.$$

## Step 5:

$$\frac{ds}{dt} = \frac{3}{5} \frac{dx}{dt}$$

## Step 6:

$$\frac{ds}{dt} = \frac{3}{5}(-5) = -3 \text{ ft/sec}$$

The shadow length is changing at the rate of  $-3$  ft/sec.

## 30. Step 1:

 $s$  = distance ball has fallen $x$  = distance from bottom of pole to shadow

## Step 2:

At the instant in question,  $s = 16 \left( \frac{1}{2} \right)^2 = 4$  ft and

$$\frac{ds}{dt} = 32 \left( \frac{1}{2} \right) = 16 \text{ ft/sec.}$$

## Step 3:

We want to find  $\frac{dx}{dt}$ .

## Step 4:

By similar triangles,  $\frac{x-30}{50-s} = \frac{x}{50}$ . This is equivalent to

$$50x - 1500 = 50x - sx, \text{ or } sx = 1500. \text{ We will use}$$

$$x = 1500s^{-1}.$$

## Step 5:

$$\frac{dx}{dt} = -500s^{-2} \frac{ds}{dt}$$

## Step 6:

$$\frac{dx}{dt} = -1500(4)^{-2}(16) = -1500 \text{ ft/sec}$$

The shadow is moving at a velocity of  $-1500$  ft/sec.

## 31. Step 1:

 $x$  = position of car ( $x = 0$  when car is right in front of you) $\theta$  = camera angle. (We assume  $\theta$  is negative until the car passes in front of you, and then positive.)

## Step 2:

At the first instant in question,  $x = 0$  ft and  $\frac{dx}{dt} = 264$  ft/sec.A half second later,  $x = \frac{1}{2}(264) = 132$  ft and  $\frac{dx}{dt} = 264$  ft/sec.

## Step 3:

We want to find  $\frac{d\theta}{dt}$  at each of the two instants.

## Step 4:

$$\theta = \tan^{-1} \left( \frac{x}{132} \right)$$

## Step 5:

$$\frac{d\theta}{dt} = \frac{1}{1 + \left( \frac{x}{132} \right)^2} \cdot \frac{1}{132} \frac{dx}{dt}$$

## 31. Continued

Step 6:

$$\text{When } x = 0: \frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{0}{132}\right)^2} \left(\frac{1}{132}\right) (264) = 2 \text{ radians/sec}$$

$$\text{When } x = 132: \frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{132}{132}\right)^2} \left(\frac{1}{132}\right) (264) = 1 \text{ radians/sec}$$

## 32. Step 1:

 $p = x$ -coordinate of plane's position $x = x$ -coordinate of car's position $s =$  distance from plane to car (line-of-sight)

Step 2:

At the instant in question,

$$p = 0, \frac{dp}{dt} = 120 \text{ mph}, s = 5 \text{ mi}, \text{ and } \frac{ds}{dt} = -160 \text{ mph.}$$

Step 3:

$$\text{We want to find } -\frac{dx}{dt}.$$

Step 4:

$$(x - p)^2 + 3^2 = s^2$$

Step 5:

$$2(x - p) \left( \frac{dx}{dt} - \frac{dp}{dt} \right) = 2s \frac{ds}{dt}$$

Step 6:

Note that, at the instant in question,

$$x = \sqrt{5^2 - 3^2} = 4 \text{ mi.}$$

$$2(4 - 0) \left( \frac{dx}{dt} - 120 \right) = 2(5)(-160)$$

$$8 \left( \frac{dx}{dt} - 120 \right) = -1600$$

$$\frac{dx}{dt} - 120 = -200$$

$$\frac{dx}{dt} = -80 \text{ mph}$$

The car's speed is 80 mph.

## 33. Step 1:

 $s =$  shadow length $\theta =$  sun's angle of elevation

Step 2:

At the instant in question,

$$s = 60 \text{ ft and } \frac{d\theta}{dt} = 0.27^\circ / \text{min} = 0.0015\pi \text{ radian/min.}$$

Step 3:

$$\text{We want to find } -\frac{ds}{dt}.$$

Step 4:

$$\tan \theta = \frac{80}{s} \text{ or } s = 80 \cot \theta$$

Step 5:

$$\frac{ds}{dt} = -80 \csc^2 \theta \frac{d\theta}{dt}$$

Step 6:

Note that, at the moment in question, since  $\tan$ 

$$\theta = \frac{80}{60} \text{ and } 0 < \theta < \frac{\pi}{2}, \text{ we have } \sin \theta = \frac{4}{5} \text{ and so}$$

$$\csc \theta = \frac{5}{4}.$$

$$\frac{ds}{dt} = -80 \left( \frac{5}{4} \right)^2 (0.0015\pi)$$

$$= -0.1875\pi \frac{\text{ft}}{\text{min}} \cdot \frac{12 \text{ in}}{1 \text{ ft}}$$

$$= -2.25\pi \text{ in./min}$$

$$\approx -7.1 \text{ in./min}$$

Since  $\frac{ds}{dt} < 0$ , the rate at which the shadow length is*decreasing* is positive. The shadow length is decreasing at the rate of approximately 7.1 in./min.

## 34. Step 1:

 $a =$  distance from origin to  $A$  $b =$  distance from origin to  $B$  $\theta =$  angle shown in problem statement

Step 2:

$$\text{At the instant in question, } \frac{da}{dt} = -2 \text{ m/sec, } \frac{db}{dt} = 1 \text{ m/sec,}$$

$$a = 10 \text{ m, and } b = 20 \text{ m.}$$

Step 3:

$$\text{We want to find } \frac{d\theta}{dt}.$$

Step 4:

$$\tan \theta = \frac{a}{b} \text{ or } \theta = \tan^{-1} \left( \frac{a}{b} \right)$$

Step 5:

$$\frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{a}{b}\right)^2} \frac{b \frac{da}{dt} - a \frac{db}{dt}}{b^2} = \frac{b \frac{da}{dt} - a \frac{db}{dt}}{a^2 + b^2}$$


Step 6:

$$\frac{d\theta}{dt} = \frac{(20)(-2) - (10)(1)}{10^2 + 20^2} = -0.1 \text{ radian/sec}$$

$$\approx -5.73 \text{ degrees/sec}$$

To the nearest degree, the angle is changing at the rate of  $-6$  degrees per second.

35. Step 1:

 $a$  = distance from  $O$  to  $A$  $b$  = distance from  $O$  to  $B$  $c$  = distance from  $A$  to  $B$ 

Step 2:

At the instant in question,  $a = 5$  nautical miles,  $b = 3$ nautical miles,  $\frac{da}{dt} = 14$  knots, and  $\frac{db}{dt} = 21$  knots.

Step 3:

We want to find  $\frac{dc}{dt}$ ,

Step 4:

Law of Cosines:  $c^2 = a^2 + b^2 - 2ab \cos 120^\circ$ 

$$c^2 = a^2 + b^2 + ab$$

Step 5:

$$2c \frac{dc}{dt} = 2a \frac{da}{dt} + 2b \frac{db}{dt} + a \frac{db}{dt} + b \frac{da}{dt}$$

Step 6:

Note that, at the instant in question,

$$c = \sqrt{a^2 + b^2 + ab} = \sqrt{(5)^2 + (3)^2 + (5)(3)} = \sqrt{49} = 7$$

$$2(7) \frac{dc}{dt} = 2(5)(14) + 2(3)(21) + (5)(21) + (3)(14)$$

$$14 \frac{dc}{dt} = 413$$

$$\frac{dc}{dt} = 29.5 \text{ knots}$$

The ships are moving apart at a rate of 29.5 knots.

36. True. Since  $\frac{dC}{dt} = 2\pi \frac{dr}{dt}$ , a constant $\frac{dr}{dt}$  results in a constant  $\frac{dC}{dt}$ .37. False. Since  $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$ , the value of  $\frac{dA}{dt}$  depends on  $r$ .38. A.  $v = s^3$ 

$$dv = 3s^2 ds$$

$$24 = 3s^2(2)$$

$$s = 2 \text{ in}$$

39. E.  $sA = 6s^2$ 

$$dsA = 12s ds$$

$$12 = 12s ds$$

$$ds = \frac{1}{s}$$

$$V = s^3$$

$$dV = 3s^2 ds = 3s^2 \frac{1}{s}$$

$$24 = 3s$$

$$s = 8 \text{ in}$$

40. C.  $\frac{x}{y} \frac{dx}{dt} = \frac{dy}{dt}$ 

$$\frac{0.6}{0.8} 3 = \frac{dy}{dt}$$

$$\frac{dy}{dt} = 2.25, \text{ but it is negative because } y \text{ is decreasing.}$$

$$\frac{dy}{dt} = -2.25.$$

41. B.  $v = \pi r^2 h$ 

$$sA = 2\pi rh$$

$$dv = \pi r^2 dh$$

$$dsA = 2\pi h dr$$

$$dv = dsA$$

$$\pi r^2 dh = 2\pi h dr$$

$$\frac{dh}{h} = 2 \frac{dr}{r^2}$$

$$\frac{2}{100} = 2 \frac{dr}{(1)^2}$$

$$dr = .01 \frac{\text{cm}}{\text{sec}}$$

42. (a)  $\frac{dc}{dt} = \frac{d}{dt}(x^3 - 6x^2 + 15x)$ 

$$= (3x^2 - 12x + 15) \frac{dx}{dt}$$

$$= [3(2)^2 - 12(2) + 15](0.1)$$

$$= 0.3$$

$$\frac{dr}{dt} = \frac{d}{dt}(9x) = 9 \frac{dx}{dt} = 9(0.1) = 0.9$$

$$\frac{dp}{dt} = \frac{dr}{dt} - \frac{dc}{dt} = 0.9 - 0.3 = 0.6$$

(b)  $\frac{dc}{dt} = \frac{d}{dt}\left(x^3 - 6x^2 + \frac{45}{x}\right)$ 

$$= \left(3x^2 - 12x - \frac{45}{x^2}\right) \frac{dx}{dt}$$

$$= \left[3(1.5)^2 - 12(1.5) - \frac{45}{1.5^2}\right](0.05)$$

$$= -1.5625$$

## 42. Continued

$$(b) \frac{dr}{dt} = \frac{d}{dt}(70x) = 70 \frac{dx}{dt} = 70(0.05) = 3.5$$

$$\frac{dp}{dt} = \frac{dr}{dt} - \frac{dc}{dt} = 3.5 - (-1.5625) = 5.0625$$

43. (a) Note that the level of the coffee in the cone is not needed until part (b).

Step 1:

$V_1$  = volume of coffee in pot

$y$  = depth of coffee in pot

Step 2:

$$\frac{dV_1}{dt} = 10 \text{ in}^3 / \text{min}$$

Step 3:

We want to find the value of  $\frac{dy}{dt}$ .

Step 4:

$$V_1 = 9\pi y$$

Step 5:

$$\frac{dV_1}{dt} = 9\pi \frac{dy}{dt}$$

Step 6:

$$10 = 9\pi \frac{dy}{dt}$$

$$\frac{dy}{dt} = \frac{10}{9\pi} \approx 0.354 \text{ in./min}$$

The level in the pot is increasing at the rate of approximately 0.354 in./min.

(b) Step 1:

$V_2$  = volume of coffee in filter

$r$  = radius of surface of coffee in filter

$h$  = depth of coffee in filter

Step 2:

At the instant in question,  $\frac{dV_2}{dt} = -10 \text{ in}^3 / \text{min}$  and

$h = 5 \text{ in.}$

Step 3:

We want to find  $-\frac{dh}{dt}$ .

Step 4:

Note that  $\frac{r}{h} = \frac{3}{6}$ , so  $r = \frac{h}{2}$ .

$$\text{Then } V_2 = \frac{1}{3} \pi r^2 h = \frac{\pi h^3}{12}.$$

Step 5:

$$\frac{dV_2}{dt} = \frac{\pi h^2}{4} \frac{dh}{dt}$$

Step 6:

$$-10 = \frac{\pi(5)^2}{4} \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{8}{5\pi} \text{ in./min}$$

Note that  $\frac{dh}{dt} < 0$ , so the rate at which the level is

falling is positive. The level in the cone is falling at the

rate of  $\frac{8}{5\pi} \approx 0.509 \text{ in./min.}$

44. Step 1:

$Q$  = rate of  $\text{CO}_2$  exhalation (mL/min)

$D$  = difference between  $\text{CO}_2$  concentration in blood

pumped to the lungs and  $\text{CO}_2$  concentration in blood

returning from the lungs (mL/L)

$y$  = cardiac output

Step 2:

At the instant in question,  $Q = 233 \text{ mL/min}$ ,  $D = 41 \text{ mL/L}$ ,

$$\frac{dD}{dt} = -2 \text{ (mL/L)/min, and } \frac{dQ}{dt} = 0 \text{ mL/min}^2.$$

Step 3:

We want to find the value of  $\frac{dy}{dt}$ .

Step 4:

$$y = \frac{Q}{D}$$

Step 5:

$$\frac{dy}{dt} = \frac{D \frac{dQ}{dt} - Q \frac{dD}{dt}}{D^2}$$

Step 6:

$$\frac{dy}{dt} = \frac{(41)(0) - (233)(-2)}{(41)^2} = \frac{466}{1681} \approx 0.277 \text{ L/min}^2$$

The cardiac output is increasing at the rate of approximately  $0.277 \text{ L/min}^{-2}$ .

45. (a) The point being plotted would correspond to a point on the edge of the wheel as the wheel turns.

(b) One possible answer is  $\theta = 16\pi t$ , where  $t$  is in seconds. (An arbitrary constant may be added to this expression, and we have assumed counterclockwise motion.)

## 45. Continued

(c) In general, assuming counterclockwise motion:

$$\frac{dx}{dt} = -2 \sin \theta \frac{d\theta}{dt} = -2(\sin \theta)(16\pi) = -32\pi \sin \theta$$

$$\frac{dy}{dt} = 2 \cos \theta \frac{d\theta}{dt} = 2(\cos \theta)(16\pi) = 32\pi \cos \theta$$

$$\text{At } \theta = \frac{\pi}{4}:$$

$$\frac{dx}{dt} = -32\pi \sin \frac{\pi}{4} = -16\pi(\sqrt{2}) \approx -71.086 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \frac{\pi}{4} = 16\pi(\sqrt{2}) \approx 71.086 \text{ ft/sec}$$

$$\text{At } \theta = \frac{\pi}{2}:$$

$$\frac{dx}{dt} = -32\pi \sin \frac{\pi}{2} = -32\pi \approx -100.531 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \frac{\pi}{2} = 0 \text{ ft/sec}$$

$$\text{At } \theta = \pi:$$

$$\frac{dx}{dt} = -32\pi \sin \pi = 0 \text{ ft/sec}$$

$$\frac{dy}{dt} = 32\pi \cos \pi = -32\pi \approx -100.531 \text{ ft/sec}$$

46. (a) One possible answer:

$$x = 30 \cos \theta, y = 40 + 30 \sin \theta$$

(b) Since the ferris wheel makes one revolution every 10 sec, we may let  $\theta = 0.2\pi t$  and we may write  $x = 30 \cos 0.2\pi t$ ,  $y = 40 + 30 \sin 0.2\pi t$ . (This assumes that the ferris wheel revolves counterclockwise.)

In general:

$$\frac{dx}{dt} = -30(\sin 0.2\pi t)(0.2\pi) = -6\pi \sin 0.2\pi t$$

$$\frac{dy}{dt} = 30(\cos 0.2\pi t)(0.2\pi) = 6\pi \cos 0.2\pi t$$

At  $t = 5$ :

$$\frac{dx}{dt} = -6\pi \sin \pi = 0 \text{ ft/sec}$$

$$\frac{dy}{dt} = 6\pi \cos \pi = 6\pi(-1) \approx -18.850 \text{ ft/sec}$$

At  $t = 8$ :

$$\frac{dx}{dt} = -6\pi \sin 1.6\pi \approx 17.927 \text{ ft/sec}$$

$$\frac{dy}{dt} = 6\pi \cos 1.6\pi \approx 5.825 \text{ ft/sec}$$

$$\begin{aligned} 47. (a) \quad \frac{dy}{dt} &= \frac{d}{dt}(uv) = u \frac{dv}{dt} + v \frac{du}{dt} \\ &= u(0.05v) + v(0.04u) \\ &= 0.09uv \\ &= 0.09y \end{aligned}$$

Since  $\frac{dy}{dt} = 0.09y$ , the rate of growth of total production is 9% per year.

$$\begin{aligned} (b) \quad \frac{dy}{dt} &= \frac{d}{dt}(uv) = u \frac{dv}{dt} + v \frac{du}{dt} \\ &= u(0.03v) + v(-0.02u) \\ &= 0.01uv \\ &= 0.01y \end{aligned}$$

The total production is increasing at the rate of 1% per year.

## Quick Quiz Sections 4.4–4.6

$$1. B. \quad x_{n+1} = x_n - \frac{f(x)}{f'(x)}$$

$$f(x) = x^3 + 2x - 1$$

$$f'(x) = 3x^2 + 2$$

$$x_2 = 1 - \frac{(1)^3 + 2(1) - 1}{3(1)^2 + 2} = \frac{3}{5}$$

$$x_3 = \frac{3}{5} - \frac{\left(\frac{3}{5}\right)^3 + 2\left(\frac{3}{5}\right) - 1}{3\left(\frac{3}{5}\right)^2 + 2} = 0.465$$

$$2. B. \quad z^2 = x^2 + y^2$$

$$z = \sqrt{4^2 + 3^2} = 5$$

$$2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$5 = 4 \left( 3 \frac{dy}{dt} \right) + 3 \frac{dy}{dt}$$

$$\frac{dy}{dt} = \frac{1}{3}$$

$$\frac{dx}{dt} = 3 \frac{dy}{dt} = 3 \left( \frac{1}{3} \right) = 1$$

$$3. A. \quad x(t) = 70$$

$$y(t) = 60t$$

$$z(t) = ((60t)^2 + 70^2)^{1/2}$$

$$\frac{dz}{dt} = \frac{1}{2}(3600t^2 + 4900)^{-1/2} (7200t)$$

$$\frac{dz}{dt} = \frac{7200(4)}{2(3600(4)^2 + 4900)^{1/2}}$$

$$\frac{dz}{dt} = 57.6$$

$$4. (a) \quad f(x) = \sqrt{x}$$

$$x = 25$$

$$f'(25) = \frac{1}{2}(25)^{-1/2} = \frac{1}{10}$$

$$\sqrt{26} = 5 + \frac{1}{10}(26 - 25) = 5.1$$

## 4. Continued

$$(b) x_{n+1} = x_n - \frac{f(x)}{f'(x)}, f(x) = x^2 - 26 = 0$$

$$x_2 = 5 - \frac{(5)^2 - 26}{2(5)} = 5.1$$

$$(c) f(x) = \sqrt[3]{x}$$

$$x = 3$$

$$f'(27) = \frac{1}{3}(27)^{-2/3} = \frac{1}{27}$$

$$\sqrt{26} = 3 + \frac{1}{27}(26 - 27)$$

$$\sqrt{26} = 2.963$$

## Chapter 4 Review (pp. 256–260)

$$1. y = x\sqrt{2-x}$$

$$\begin{aligned} y' &= x \left( \frac{1}{2\sqrt{2-x}} \right) (-1) + (\sqrt{2-x})(1) \\ &= \frac{-x + 2(2-x)}{2\sqrt{2-x}} \\ &= \frac{4-3x}{2\sqrt{2-x}} \end{aligned}$$

The first derivative has a zero at  $\frac{4}{3}$ .

$$\text{Critical point value: } x = \frac{4}{3} \quad y = \frac{4\sqrt{6}}{9} \approx 1.09$$

$$\text{Endpoint values: } \begin{aligned} x = -2 \quad y = -4 \\ x = 2 \quad y = 0 \end{aligned}$$

The global maximum value is  $\frac{4\sqrt{6}}{9}$  at  $x = \frac{4}{3}$ , and the global minimum value is  $-4$  at  $x = -2$ .

2. Since  $y$  is a cubic function with a positive leading coefficient, we have  $\lim_{x \rightarrow -\infty} y = -\infty$  and  $\lim_{x \rightarrow \infty} y = \infty$ . There are no global extrema.


$$\begin{aligned} 3. y' &= (x^2)(e^{1/x^2})(-2x^{-3}) + (e^{1/x^2})(2x) \\ &= 2e^{1/x^2} \left( -\frac{1}{x} + x \right) \\ &= \frac{2e^{1/x^2}(x-1)(x+1)}{x} \end{aligned}$$

Intervals	$x < -1$	$-1 < x < 0$	$0 < x < 1$	$x > 1$
Sign of $y'$	-	+	-	+
Behavior of $y$	Decreasing	Increasing	Decreasing	Increasing

$$\begin{aligned} y'' &= \frac{d}{dx} [2e^{1/x^2}(-x^{-1} + x)] \\ &= (2e^{1/x^2})(x^{-2} + 1) + (-x^{-1} + x)(2e^{1/x^2})(-2x^{-3}) \\ &= (2e^{1/x^2})(x^{-2} + 1 + 2x^{-4} - 2x^{-2}) \\ &= \frac{2e^{1/x^2}(x^4 - x^2 + 2)}{x^4} \\ &= \frac{2e^{1/x^2}[(x^2 - 0.5)^2 + 1.75]}{x^4} \end{aligned}$$

The second derivative is always positive (where defined), so the function is concave up for all  $x \neq 0$ .

Graphical support:



$[-4, 4]$  by  $[-1, 5]$

(a)  $[-1, 0)$  and  $(1, \infty)$

(b)  $(-\infty, -1]$  and  $(0, 1]$

(c)  $(-\infty, 0)$  and  $(0, \infty)$

(d) None

(e) Local (and absolute) minima at  $(1, e)$  and  $(-1, e)$

(f) None

4. Note that the domain of the function is  $[-2, 2]$ .

$$\begin{aligned} y' &= x \left( \frac{1}{2\sqrt{4-x^2}} \right) (-2x) + (\sqrt{4-x^2})(1) \\ &= \frac{-x^2 + (4-x^2)}{\sqrt{4-x^2}} \\ &= \frac{4-2x^2}{\sqrt{4-x^2}} \end{aligned}$$

Intervals	$-2 < x < -\sqrt{2}$	$-\sqrt{2} < x < \sqrt{2}$	$\sqrt{2} < x < 2$
Sign of $y'$	-	+	-
Behavior of $y$	Decreasing	Increasing	Decreasing


$$\begin{aligned} y'' &= \frac{(\sqrt{4-x^2})(-4x) - (4-2x^2) \left( \frac{1}{2\sqrt{4-x^2}} \right) (-2x)}{4-x^2} \\ &= \frac{2x(x^2-6)}{(4-x^2)^{3/2}} \end{aligned}$$

Note that the values  $x = \pm\sqrt{6}$  are not zeros of  $y''$  because they fall outside of the domain.

Intervals	$-2 < x < 0$	$0 < x < 2$
Sign of $y''$	+	-
Behavior of $y$	Concave up	Concave down

4. Continued

Graphical support:



$[-2.35, 2.35]$  by  $[-3.5, 3.5]$

- (a)  $[-\sqrt{2}, \sqrt{2}]$
- (b)  $[-2, -\sqrt{2}]$  and  $[\sqrt{2}, 2]$
- (c)  $(-2, 0)$
- (d)  $(0, 2)$
- (e) Local maxima:  $(-2, 0), (\sqrt{2}, 2)$

Local minima:  $(2, 0), (-\sqrt{2}, -2)$

Note that the extrema at  $x = \pm\sqrt{2}$  are also absolute extrema.

- (f)  $(0, 0)$

5.  $y' = 1 - 2x - 4x^3$


Using grapher techniques, the zero of  $y'$  is  $x \approx 0.385$ .

Intervals	$x < 0.385$	$0.385 < x$
Sign of $y'$	+	-
Behavior of $y$	Increasing	Decreasing

$y'' = -2 - 12x^2 = -2(1 + 6x^2)$

The second derivative is always negative so the function is concave down for all  $x$ .

Graphical support:



$[-4, 4]$  by  $[-4, 2]$

- (a) Approximately  $(-\infty, 0.385]$
- (b) Approximately  $[0.385, \infty)$
- (c) None
- (d)  $(-\infty, \infty)$
- (e) Local (and absolute) maximum at  $\approx (0.385, 1.215)$
- (f) None


6.  $y' = e^{x-1} - 1$

Intervals	$x < 1$	$1 < x$
Sign of $y'$	-	+
Behavior of $y$	Decreasing	Increasing

$y'' = e^{x-1}$

The second derivative is always positive, so the function is concave up for all  $x$ .

Graphical support:



$[-4, 4]$  by  $[-2, 4]$

- (a)  $[1, \infty)$
- (b)  $(-\infty, 1]$
- (c)  $(-\infty, \infty)$
- (d) None
- (e) Local (and absolute) minimum at  $(1, 0)$
- (f) None

7. Note that the domain is  $(-1, 1)$ .

$y = (1 - x^2)^{-1/4}$

$y' = -\frac{1}{4}(1 - x^2)^{-5/4}(-2x) = \frac{x}{2(1 - x^2)^{5/4}}$

Intervals	$-1 < x < 0$	$0 < x < 1$
Sign of $y'$	-	+
Behavior of $y$	Decreasing	Increasing


$$y'' = \frac{2(1-x^2)^{5/4} - (x)(2) \left(\frac{5}{4}\right)(1-x^2)^{1/4}(-2x)}{4(1-x^2)^{5/2}}$$

$$= \frac{(1-x^2)^{1/4}[2-2x^2+5x^2]}{4(1-x^2)^{5/2}}$$

$$= \frac{3x^2+2}{4(1-x^2)^{9/4}}$$

The second derivative is always positive, so the function is concave up on its domain  $(-1, 1)$ .

Graphical support:



$[-1.3, 1.3]$  by  $[-1, 3]$

- (a)  $[0, 1)$
- (b)  $(-1, 0]$
- (c)  $(-1, 1)$
- (d) None
- (e) Local minimum at  $(0, 1)$
- (f) None



$$8. y' = \frac{(x^3 - 1)(1) - (x)(3x^2)}{(x^3 - 1)^2} = \frac{2x^3 + 1}{(x^3 - 1)^2}$$

Intervals	$x < -2^{-1/3}$	$-2^{-1/3} < x < 1$	$1 < x$
Sign of $y'$	+	-	-
Behavior of $y$	Increasing	Decreasing	Decreasing

$$y'' = -\frac{(x^3 - 1)^2(6x^2) - (2x^3 + 1)(2)(x^3 - 1)(3x^2)}{(x^3 - 1)^4}$$

$$= -\frac{(x^3 - 1)(6x^2) - (2x^3 + 1)(6x^2)}{(x^3 - 1)^3}$$

$$= \frac{6x^2(x^3 + 2)}{(x^3 - 1)^3}$$

Intervals	$x < -2^{1/3}$	$-2^{1/3} < x < 0$	$0 < x < 1$	$1 < x$
Sign of $y''$	+	-	-	+
Behavior of $y$	Concave up	Concave down	Concave down	Concave up

Graphical support:



[-4.7, 4.7] by [-3.1, 3.1]

- (a)  $(-\infty, -2^{-1/3}] = (-\infty, -0.794]$
- (b)  $[-2^{-1/3}, 1) = [-0.794, 1)$  and  $(1, \infty)$
- (c)  $(-\infty, -2^{1/3}) = (-\infty, -1.260)$  and  $(1, \infty)$
- (d)  $(-2^{1/3}, 1) = (-1.260, 1)$
- (e) Local minimum at  $(-2^{-1/3}, \frac{2}{3} \cdot 2^{-1/3}) = (-0.794, 0.529)$
- (f)  $(-2^{1/3}, \frac{1}{3} \cdot 2^{1/3}) = (-1.260, 0.420)$

9. Note that the domain is  $[-1, 1]$ .

$$y' = -\frac{1}{\sqrt{1-x^2}}$$


Since  $y'$  is negative on  $(-1, 1)$  and  $y$  is continuous,  $y$  is decreasing on its domain  $[-1, 1]$ .

$$y'' = \frac{d}{dx} [-(1-x^2)^{-1/2}]$$

$$= \frac{1}{2}(1-x^2)^{-3/2}(-2x) = -\frac{x}{(1-x^2)^{3/2}}$$

Intervals	$-1 < x < 0$	$0 < x < 1$
Sign of $y''$	+	-
Behavior of $y$	Concave up	Concave down


Graphical support:




[-1.175, 1.175] by  $[-\frac{\pi}{4}, \frac{5\pi}{4}]$

- (a) None
- (b)  $[-1, 1]$
- (c)  $(-1, 0)$
- (d)  $(0, 1)$
- (e) Local (and absolute) maximum at  $(-1, \pi)$ ;  
local (and absolute) minimum at  $(1, 0)$
- (f)  $(0, \frac{\pi}{2})$


10. This problem can be solved graphically by using NDER to obtain the graphs shown below.



[-4, 4] by [-1, 0.3]



[-4, 4] by [-0.4, 0.6]



[-4, 4] by [-0.7, 0.8]

An alternative approach using a combination of algebraic and graphical techniques follows. Note that the denominator of  $y$  is always positive because it is equivalent to  $(x+1)^2 + 2$ .

$$y' = \frac{(x^2 + 2x + 3)(1) - (x)(2x + 2)}{(x^2 + 2x + 3)^2}$$

$$= \frac{-x^2 + 3}{(x^2 + 2x + 3)^2}$$

Intervals	$x < -\sqrt{3}$	$-\sqrt{3} < x < \sqrt{3}$	$\sqrt{3} < x$
Sign of $y'$	-	+	-
Behavior of $y$	Decreasing	Increasing	Decreasing

$$y'' = \frac{(x^2 + 2x + 3)^2(-2x) - (-x^2 + 3)(2)(x^2 + 2x + 3)(2x + 2)}{(x^2 + 2x + 3)^4}$$

$$= \frac{(x^2 + 2x + 3)(-2x) - 2(2x + 2)(-x^2 + 3)}{(x^2 + 2x + 3)^3}$$

$$= \frac{2x^3 - 18x - 12}{(x^2 + 2x + 3)^3}$$

10. Continued

Using graphing techniques, the zeros of  $2x^3 - 18x - 12$  (and hence of  $y''$ ) are at  $x \approx -2.584$ ,  $x \approx -0.706$ , and  $x \approx 3.290$ .

Intervals	$(-\infty, -2.584)$	$(-2.584, -0.706)$	$(-0.706, 3.290)$	$(3.290, \infty)$
Sign of $y''$	-	+	-	+
Behavior of $y$	Concave down	Concave up	Concave down	Concave up

- (a)  $[-\sqrt{3}, \sqrt{3}]$
- (b)  $(-\infty, -\sqrt{3})$  and  $(\sqrt{3}, \infty)$
- (c) Approximately  $(-2.584, -0.706)$  and  $(3.290, \infty)$
- (d) Approximately  $(-\infty, -2.584)$  and  $(-0.706, 3.290)$
- (e) Local maximum at  $(\sqrt{3}, \frac{\sqrt{3}-1}{4}) \approx (1.732, 0.183)$ ;  
local minimum at  $(-\sqrt{3}, \frac{-\sqrt{3}-1}{4}) \approx (-1.732, -0.683)$
- (f)  $\approx (-2.584, -0.573)$ ,  $(-0.706, -0.338)$ , and  $(3.290, 0.161)$

11. For  $x > 0$ ,  $y' = \frac{d}{dx} \ln x = \frac{1}{x}$   
 For  $x < 0$ :  $y' = \frac{d}{dx} \ln(-x) = \frac{1}{-x}(-1) = \frac{1}{x}$


Thus  $y' = \frac{1}{x}$  for all  $x$  in the domain.

Intervals	$(-2, 0)$	$(0, 2)$
Sign of $y'$	-	+
Behavior of $y$	Decreasing	Increasing

$y'' = -x^{-2}$

The second derivative always negative, so the function is concave down on each open interval of its domain.

Graphical support:



$[-2.35, 2.35]$  by  $[-3, 1.5]$

- (a)  $(0, 2]$
- (b)  $[-2, 0)$
- (c) None
- (d)  $(-2, 0)$  and  $(0, 2)$

(e) Local (and absolute) maxima at  $(-2, \ln 2)$  and  $(2, \ln 2)$

(f) None

12.  $y' = 3 \cos 3x - 4 \sin 4x$

Using graphing techniques, the zeros of  $y'$  in the domain

$0 \leq x < 2\pi$  are  $x \approx 0.176$ ,  $x \approx 0.994$ ,  $x = \frac{\pi}{2} \approx 1.57$ ,  
 $x \approx 2.148$ , and  $x \approx 2.965$ ,  $x = 3.834$ ,  $x = \frac{3\pi}{2}$ ,  $x \approx 5.591$

Intervals	$0 < x < 0.176$	$0.176 < x < 0.994$	$0.994 < x < \frac{\pi}{2}$	$\frac{\pi}{2} < x < 2.148$	$2.148 < x < 2.965$
Sign of $y'$	+	-	+	-	+
Behavior of $y$	Increasing	Decreasing	Increasing	Decreasing	Increasing

Intervals	$2.965 < x < 3.834$	$3.834 < x < \frac{3\pi}{2}$	$\frac{3\pi}{2} < x < 5.591$	$5.591 < x < 2\pi$
Sign of $y'$	-	+	-	+
Behavior of $y$	Decreasing	Increasing	Decreasing	Increasing

$y'' = -9 \sin 3x - 16 \cos 4x$


Using graphing techniques, the zeros of  $y''$  in the domain

$0 \leq x < 2\pi$  are  $x \approx 0.542$ ,  $x \approx 1.266$ ,  $x \approx 1.876$ ,  
 $x \approx 2.600$ ,  $x \approx 3.425$ ,  $x \approx 4.281$ ,  $x \approx 5.144$  and  $x \approx 6.000$ .

Intervals	$0 < x < 0.542$	$0.542 < x < 1.266$	$1.266 < x < 1.876$	$1.876 < x < 2.600$	$2.600 < x < 3.425$
Sign of $y''$	-	+	-	+	-
Behavior of $y$	Concave down	Concave up	Concave down	Concave up	Concave down

Intervals	$3.425 < x < 4.281$	$4.281 < x < 5.144$	$5.144 < x < 6.000$	$6.00 < x < 2\pi$
Sign of $y''$	+	-	+	-
Behavior of $y$	Concave up	Concave down	Concave up	Concave down

Graphical support:



$[-\frac{\pi}{4}, \frac{9\pi}{4}]$  by  $[-2.5, 2.5]$

12. Continued

(a) Approximately  $[0, 0.176]$ ,

$$\left[0.994, \frac{\pi}{2}\right], [2.148, 2.965], \left[3.834, \frac{3\pi}{2}\right], \text{ and } [5.591, 2\pi]$$

(b) Approximately  $[0.176, 0.994]$ ,

$$\left[\frac{\pi}{2}, 2.148\right], [2.965, 3.834], \text{ and } \left[\frac{3\pi}{2}, 5.591\right]$$

(c) Approximately  $(0.542, 1.266)$ ,  $(1.876, 2.600)$ ,  $(3.425, 4.281)$ , and  $(5.144, 6.000)$

(d) Approximately  $(0, 0.542)$ ,  $(1.266, 1.876)$ ,  $(2.600, 3.425)$ ,  $(4.281, 5.144)$ , and  $(6.000, 2\pi)$

(e) Local maxima at  $\approx (0.176, 1.266)$ ,  $\left(\frac{\pi}{2}, 0\right)$

and  $(2.965, 1.266)$ ,  $\left(\frac{3\pi}{2}, 2\right)$ , and  $(2\pi, 1)$ ;

local minima at  $\approx (0, 1)$ ,  $(0.994, -0.513)$ ,  $(2.148, -0.513)$ ,  $(3.834, -1.806)$ , and  $(5.591, -1.806)$

Note that the local extrema at  $x = 3.834$ ,  $x = \frac{3\pi}{2}$ , and  $x = 5.591$  are also extrema.

(f)  $\approx (0.542, 0.437)$ ,  $(1.266, -0.267)$ ,  $(1.876, -0.267)$ ,  $(2.600, 0.437)$ ,  $(3.425, -0.329)$ ,  $(4.281, 0.120)$ ,  $(5.144, 0.120)$ , and  $(6.000, -0.329)$


13.  $y' = \begin{cases} -e^{-x}, & x < 0 \\ 4 - 3x^2, & x > 0 \end{cases}$

Intervals	$x < 0$	$0 < x < \frac{2}{\sqrt{3}}$	$\frac{2}{\sqrt{3}} < x$
Sign of $y'$	-	+	-
Behavior of $y$	Decreasing	Increasing	Decreasing

$$y'' = \begin{cases} e^{-x}, & x > 0 \\ -6x, & x < 0 \end{cases}$$

Intervals	$x < 0$	$0 < x$
Sign of $y''$	+	-
Behavior of $y$	Concave up	Concave down

Graphical support:



$[-4, 4]$  by  $[-2, 4]$

(a)  $\left[0, \frac{2}{\sqrt{3}}\right]$

(b)  $(-\infty, 0]$  and  $\left[\frac{2}{\sqrt{3}}, \infty\right)$

(c)  $(-\infty, 0)$

(d)  $(0, \infty)$

(e) Local maximum at  $\left(\frac{2}{\sqrt{3}}, \frac{16}{3\sqrt{3}}\right) \approx (1.155, 3.079)$

(f) None. Note that there is no point of inflection at  $x = 0$  because the derivative is undefined and no tangent line exists at this point.

14.  $y' = -5x^4 + 7x^2 + 10x + 4$

Using graphing techniques, the zeros of  $y'$  are  $x \approx -0.578$  and  $x \approx -1.692$ .

Intervals	$x < -0.578$	$-0.578 < x < 1.692$	$1.692 < x$
Sign of $y'$	-	+	-
Behavior of $y$	Decreasing	Increasing	Decreasing

$$y'' = -20x^3 + 14x + 10$$

Using graphing techniques, the zeros of  $y''$  is  $x = 1.079$ .

Intervals	$x < 1.079$	$1.079 < x$
Sign of $y''$	+	-
Behavior of $y$	Concave up	Concave down

Graphical support:



$[-4, 4]$  by  $[-10, 25]$

(a) Approximately  $[-0.578, 1.692]$

(b) Approximately  $(-\infty, -0.578]$  and  $[1.692, \infty)$

(c) Approximately  $(-\infty, 1.079)$

(d) Approximately  $(1.079, \infty)$

(e) Local maximum at  $\approx (1.692, 20.517)$ ; local minimum at  $\approx (-0.578, 0.972)$

(f)  $\approx (1.079, 13.601)$

15.  $y = 2x^{4/5} - x^{9/5}$


$$y' = \frac{8}{5}x^{-1/5} - \frac{9}{5}x^{4/5} = \frac{8-9x}{5\sqrt[5]{x}}$$

Intervals	$x < 0$	$0 < x < \frac{8}{9}$	$\frac{8}{9} < x$
Sign of $y'$	-	+	-
Behavior of $y$	Decreasing	Increasing	Decreasing

$$y'' = -\frac{8}{25}x^{-6/5} - \frac{36}{25}x^{-1/5} = \frac{4(2+9x)}{25x^{6/5}}$$

Intervals	$x < -\frac{2}{9}$	$-\frac{2}{9} < x < 0$	$0 < x$
Sign of $y''$	+	-	-
Behavior of $y$	Concave up	Concave down	Concave down

Graphical support:



[-4, 4] by [-3, 3]

(a)  $\left[0, \frac{8}{9}\right]$

(b)  $(-\infty, 0]$  and  $\left[\frac{8}{9}, \infty\right)$

(c)  $\left(-\infty, -\frac{2}{9}\right)$

(d)  $\left(-\frac{2}{9}, 0\right)$  and  $(0, \infty)$

(e) Local maximum


at  $\left(\frac{8}{9}, \frac{10}{9} \cdot \left(\frac{8}{9}\right)^{4/5}\right) \approx (0.889, 1.011)$ ; local minimum

at  $(0, 0)$

(f)  $\left(-\frac{2}{9}, \frac{20}{9} \cdot \left(-\frac{2}{9}\right)^{4/5}\right) \approx \left(-\frac{2}{9}, 0.667\right)$

16. We use a combination of analytic and grapher techniques to solve this problem. Depending on the viewing windows chosen, graphs obtained using NDER may exhibit strange behavior near  $x = 2$  because, for example, NDER  $(y, 2) \approx 5,000,000$  while  $y'$  is actually undefined at

$x = 2$ . The graph of  $y = \frac{5-4x+4x^2-x^3}{x-2}$  is shown below.




[-5.875, 5.875] by [-50, 30]

$$y' = \frac{(x-2)(-4+8x-3x^2) - (5-4x+4x^2-x^3)(1)}{(x-2)^2}$$

$$= \frac{-2x^3+10x^2-16x+3}{(x-2)^2}$$

The graph of  $y'$  is shown below.



[-5.875, 5.875] by [-50, 30]

The zero of  $y'$  is  $x \approx 0.215$ .


Intervals	$x < 0.215$	$0.215 < x < 2$	$2 < x$
Sign of $y'$	+	-	-
Behavior of $y$	Increasing	Decreasing	Decreasing

$$y'' = \frac{(x-2)^2(-6x^2+20x-16) - (-2x^3+10x^2-16x+3)(2)(x-2)}{(x-2)^4}$$

$$= \frac{(x-2)(-6x^2+20x-16) - 2(-2x^3+10x^2-16x+3)}{(x-2)^3}$$

$$= \frac{-2(x^3-6x^2+12x-13)}{(x-2)^3}$$

The graph of  $y''$  is shown below.



[-5.875, 5.875] by [-20, 20]

The zero of  $x^3 - 6x^2 + 12x - 13$  (and hence of  $y''$ ) is  $x \approx 3.710$ .

Intervals	$x < 2$	$2 < x < 3.710$	$3.710 < x$
Sign of $y''$	-	+	-
Behavior of $y$	Concave down	Concave up	Concave down

16. Continued

- (a) Approximately  $(-\infty, 0.215]$
- (b) Approximately  $[0.215, 2)$  and  $(2, \infty)$
- (c) Approximately  $(2, 3.710)$
- (d)  $(-\infty, 2)$  and approximately  $(3.710, \infty)$
- (e) Local maximum at  $\approx (0.215, -2.417)$
- (f)  $\approx (3.710, -3.420)$

17.  $y' = 6(x+1)(x-2)^2$

Intervals	$x < -1$	$-1 < x < 2$	$2 < x$
Sign of $y'$	-	+	+
Behavior of $y$	Decreasing	Increasing	Increasing

$$y'' = 6(x+1)(2)(x-2) + 6(x-2)^2(1)$$

$$= 6(x-2)[2(x+2) + (x-2)]$$

$$= 18x(x-2)$$

Intervals	$x < 0$	$0 < x < 2$	$2 < x$
Sign of $y''$	+	-	+
Behavior of $y$	Concave up	Concave down	Concave up

- (a) There are no local maxima.
- (b) There is a local (and absolute) minimum at  $x = -1$ .
- (c) There are points of inflection at  $x = 0$  and at  $x = 2$ .

18.  $y' = 6(x+1)(x-2)$

Intervals	$x < -1$	$-1 < x < 2$	$2 < x$
Sign of $y'$	+	-	+
Behavior of $y$	Increasing	Decreasing	Increasing

$$y'' = \frac{d}{dx} 6(x^2 - x - 2) = 6(2x - 1)$$

Intervals	$x < \frac{1}{2}$	$\frac{1}{2} < x$
Sign of $y''$	-	+
Behavior of $y$	Concave down	Concave up

- (a) There is a local maximum at  $x = -1$ .
- (b) There is a local maximum at  $x = 2$ .
- (c) There is a point of inflection at  $x = \frac{1}{2}$ .

19. Since  $\frac{d}{dx} \left( -\frac{1}{4}x^{-4} - e^{-x} \right) = x^{-5} + e^{-x}$ ,

$$f(x) = -\frac{1}{4}x^{-4} - e^{-x} + C.$$

20. Since  $\frac{d}{dx} \sec x = \sec x \tan x$ ,  $f(x) = \sec x + C$ .

21. Since  $\frac{d}{dx} \left( 2 \ln x + \frac{1}{3}x^3 + x \right) = \frac{2}{x} + x^2 + 1$ ,

$$f(x) = 2 \ln x + \frac{1}{3}x^3 + x + C.$$

22. Since  $\frac{d}{dx} \left( \frac{2}{3}x^{3/2} + 2x^{1/2} \right) = \sqrt{x} + \frac{1}{\sqrt{x}}$ ,

$$f(x) = \frac{2}{3}x^{3/2} + 2x^{1/2} + C.$$

23.  $f(x) = -\cos x + \sin x + C$

$$f(\pi) = 3$$

$$1 + 0 + C = 3$$

$$C = 2$$

$$f(x) = -\cos x + \sin x + 2$$

24.  $f(x) = \frac{3}{4}x^{4/3} + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + C$

$$f(1) = 0$$

$$\frac{3}{4} + \frac{1}{3} + \frac{1}{2} + 1 + C = 0$$

$$C = -\frac{31}{12}$$

$$f(x) = \frac{3}{4}x^{4/3} + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x - \frac{31}{12}$$

25.  $v(t) = s'(t) = 9.8t + 5$

$$s(t) = 4.9t^2 + 5t + C$$

$$s(0) = 10$$

$$C = 10$$

$$s(t) = 4.9t^2 + 5t + 10$$

26.  $a(t) = v'(t) = 32$

$$v(t) = 32t + C_1$$

$$v(0) = 20$$

$$C_1 = 20$$

$$v(t) = s'(t) = 32t + 20$$

$$s(t) = 16t^2 + 20t + C_2$$

$$s(0) = 5$$

$$C_2 = 5$$

$$s(t) = 16t^2 + 20t + 5$$

27.  $f(x) = \tan x$   
 $f'(x) = \sec^2 x$

$$\begin{aligned} L(x) &= f\left(-\frac{\pi}{4}\right) + f'\left(-\frac{\pi}{4}\right)\left[x - \left(-\frac{\pi}{4}\right)\right] \\ &= \tan\left(-\frac{\pi}{4}\right) + \sec^2\left(-\frac{\pi}{4}\right)\left(x + \frac{\pi}{4}\right) \\ &= -1 + 2\left(x + \frac{\pi}{4}\right) \\ &= 2x + \frac{\pi}{2} - 1 \end{aligned}$$

28.  $f(x) = \sec x$   
 $f'(x) = \sec x \tan x$

$$\begin{aligned} L(x) &= f\left(\frac{\pi}{4}\right) + f'\left(\frac{\pi}{4}\right)\left(x - \frac{\pi}{4}\right) \\ &= \sec\left(\frac{\pi}{4}\right) + \sec\left(\frac{\pi}{4}\right)\tan\left(\frac{\pi}{4}\right)\left(x - \frac{\pi}{4}\right) \\ &= \sqrt{2} + \sqrt{2}(1)\left(x - \frac{\pi}{4}\right) \\ &= \sqrt{2}x - \frac{\pi\sqrt{2}}{4} + \sqrt{2} \end{aligned}$$

29.  $f(x) = \frac{1}{1 + \tan x}$   
 $f'(x) = -(1 + \tan x)^{-2}(\sec^2 x)$   
 $= -\frac{1}{\cos^2 x(1 + \tan x)^2}$   
 $= -\frac{1}{(\cos x + \sin x)^2}$   
 $L(x) = f(0) + f'(0)(x - 0)$   
 $= 1 - 1(x - 0)$   
 $= -x + 1$

30.  $f(x) = e^x + \sin x$   
 $f'(x) = e^x + \cos x$   
 $L(x) = f(0) + f'(0)(x - 0)$   
 $= 1 + 2(x - 0)$   
 $= 2x + 1$

31. The global minimum value of  $\frac{1}{2}$  occurs at  $x = 2$ .

32. (a) The values of  $y'$  and  $y''$  are both negative where the graph is decreasing and concave down, at  $T$ .

(b) The value of  $y'$  is negative and the value of  $y''$  is positive where the graph is decreasing and concave up, at  $P$ .


33. (a) The function is increasing on the interval  $(0, 2]$ .

(b) The function is decreasing on the interval  $[-3, 0)$ .

(c) The local extreme values occur only at the endpoints of the domain. A local maximum value of 1 occurs at  $x = -13$ , and a local maximum value of 3 occurs at  $x = 2$ .

34. The 24th day


35.



36. (a) We know that  $f$  is decreasing on  $[0, 1]$  and increasing on  $[1, 3]$ , the absolute minimum value occurs at  $x = 1$  and the absolute maximum value occurs at an endpoint. Since  $f(0) = 0$ ,  $f(1) = -2$ , and  $f(3) = 3$ , the absolute minimum value is  $-2$  at  $x = 1$  and the absolute maximum value is  $3$  at  $x = 3$ .

(b) The concavity of the graph does not change. There are no points of inflection.

(c)



37. (a)  $f(x)$  is continuous on  $[0.5, 3]$  and differentiable on  $(0.5, 3)$ .

(b)  $f'(x) = (x)\left(\frac{1}{x}\right) + (\ln x)(1) = 1 + \ln x$

Using  $a = 0.5$  and  $b = 3$ , we solve as follows.

$$\begin{aligned} f'(c) &= \frac{f(3) - f(0.5)}{3 - 0.5} \\ 1 + \ln c &= \frac{3 \ln 3 - 0.5 \ln 0.5}{2.5} \\ \ln c &= \frac{\ln\left(\frac{3^3}{0.5^{0.5}}\right)}{2.5} - 1 \\ \ln c &= 0.4 \ln(27\sqrt{2}) - 1 \\ c &= e^{-1}(27\sqrt{2})^{0.4} \\ c &= e^{-1}\sqrt[3]{1458} \approx 1.579 \end{aligned}$$

(c) The slope of the line is

$$m = \frac{f(b) - f(a)}{b - a} = 0.4 \ln(27\sqrt{2}) - 0.2 \ln 1458, \text{ and the line passes through } (3, 3 \ln 3). \text{ Its equation is } y = 0.2(\ln 1458)(x - 3) + 3 \ln 3, \text{ or approximately } y = 1.457x - 1.075.$$

## 37. Continued

(d) The slope of the line is  $m = 0.2 \ln 1458$ , and the line passes through

$$(c, f(c)) = (e^{-1} \sqrt[5]{1458}, e^{-1} \sqrt[5]{1458}(-1 + 0.2 \ln 1458)) \\ \approx (1.579, 0.722).$$

Its equation is

$$y = 0.2(\ln 1458)(x - c) + f(c), \\ y = 0.2 \ln 1458(x - e^{-1} \sqrt[5]{1458}) \\ + e^{-1} \sqrt[5]{1458}(-1 + 0.2 \ln 1458),$$

$$y = 0.2(\ln 1458)x - e^{-1} \sqrt[5]{1458}, \\ \text{or approximately } y = 1.457x - 1.579.$$

38. (a)  $v(t) = s'(t) = 4 - 6t - 3t^2$

(b)  $a(t) = v'(t) = -6 - 6t$

(c) The particle starts at position 3 moving in the positive direction, but decelerating. At approximately  $t = 0.528$ , it reaches position 4.128 and changes direction, beginning to move in the negative direction. After that, it continues to accelerate while moving in the negative direction.

39. (a)  $L(x) = f(0) + f'(0)(x - 0) \\ = -1 + 0(x - 0) = -1$


(b)  $f(0.1) \approx L(0.1) = -1$

(c) Greater than the approximation in (b), since  $f''(x)$  is actually positive over the interval  $(0, 0.1)$  and the estimate is based on the derivative being 0.

40. (a) Since  $\frac{dy}{dx} = (x^2)(-e^{-x}) + (e^{-x})(2x) + (2x - x^2)e^{-x}$ ,  
 $dy = (2x - x^2)e^{-x} dx$ .

(b)  $dy = [2(1) - (1)^2](e^{-1})(0.01) \\ = 0.01e^{-1} \\ \approx 0.00368$

41. (a) With some rounding,  $y = \frac{1633001.59}{1 + 17.471e^{-0.06378t}}$



(c)  $y = \frac{1633001.59}{1 + 17.471e^{-0.06378(80)}} + 829,210 = 2,305,337$


(d) Using the Second Derivative, we find the maximum rate of growth about 1885. We find a point of inflection here, which shows the beginning of a decline in the rate of growth.

(e)  $y = \frac{1633001.59}{1 + 17.471e^{-0.06378(\infty)}} \approx 2,462,000$ , which is the approximate maximum population.

(f) There are many possible causes. Advances in transportation began drawing the population southward after 1920, and Tennessee was well-situated geographically to become a crossroads of river, railroad, and automobile routes. By the year 2000 there had been numerous other demographic changes. It should be pointed out that the census years in the data (1850–1910) include the years of the Civil War and Reconstruction, so the regression is based on unusual data.

42.  $f(x) = 2 \cos x - \sqrt{1+x}$   
 $f'(x) = -2 \sin x - \frac{1}{2\sqrt{1+x}}$   
 $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$   
 $= x_n - \frac{2 \cos x_n - \sqrt{1+x_n}}{-2 \sin x_n - \frac{1}{2\sqrt{1+x_n}}}$

The graph of  $y = f(x)$  shows that  $f(x) = 0$  has one solution, near  $x = 1$ .



$x_1 = 1$   
 $x_2 \approx 0.8361848$   
 $x_3 \approx 0.8283814$   
 $x_4 \approx 0.8283608$   
 $x_5 \approx 0.8283608$

Solution:  $x \approx 0.828361$

43. Let  $t$  represent time in seconds, where the rocket lifts off at  $t = 0$ . Since  $a(t) = v'(t) = 20$ , m/sec<sup>2</sup> and  $v(0) = 0$  m/sec, we have  $v(t) = 20t$ , and so  $v(60) = 1200$  m/sec. The speed after 1 minute (60 seconds) will be 1200 m/sec.

44. Let  $t$  represent time in seconds, where the rock is blasted upward at  $t = 0$ . Since  $a(t) = v'(t) = -3.72 \text{ m/sec}^2$  and  $v(0) = 93 \text{ m/sec}$ , we have  $v(t) = -3.72t + 93$ . Since  $s'(t) = -3.72t + 93$  and  $s(0) = 0$ , we have  $s(t) = -1.86t^2 + 93t$ . Solving  $v(t) = 0$ , we find that the rock attains its maximum height at  $t = 25$  sec and its height at that time is  $s(25) = 1162.5 \text{ m}$ .

45. Note that  $s = 100 - 2r$  and the sector area is given by


$$A = \pi r^2 \left( \frac{s}{2\pi r} \right) = \frac{1}{2}rs = \frac{1}{2}r(100 - 2r) = 50r - r^2. \text{ To find}$$

the domain of  $A(r) = 50r - r^2$ , note that  $r > 0$  and

$$0 < s < 2\pi r, \text{ which gives } 12.1 \approx \frac{50}{\pi + 1} < r < 50. \text{ Since}$$

$A'(r) = 50 - 2r$ , the critical point occurs at  $r = 25$ . This value is in the domain and corresponds to the maximum area because  $A''(r) = -2$ , which is negative for all  $r$ . The greatest area is attained when  $r = 25$  ft and  $s = 50$  ft.

46.



For  $0 < x < \sqrt{27}$ , the triangle with vertices at  $(0, 0)$  and  $(\pm x, 27 - x^2)$  has an area given by

$$A(x) = \frac{1}{2}(2x)(27 - x^2) = 27x - x^3. \text{ Since}$$

$A' = 27 - 3x^2 = 3(3 - x)(3 + x)$  and  $A'' = -6x$ , the critical point in the interval  $(0, \sqrt{27})$  occurs at  $x = 3$  and corresponds to the maximum area because  $A''(x)$  is negative in this interval. The largest possible area is  $A(3) = 54$  square units.

47. If the dimensions are  $x$  ft by  $x$  ft by  $h$  ft, then the total amount of steel used is  $x^2 + 4xh$  ft<sup>2</sup>. Therefore,

$$x^2 + 4xh = 108 \text{ and so } h = \frac{108 - x^2}{4x}. \text{ The volume is given}$$

$$\text{by } V(x) = x^2h = \frac{108x - x^3}{4} = 27x - 0.25x^3. \text{ Then}$$

$$V'(x) = 27 - 0.75x^2 = 0.75(6 + x)(6 - x) \text{ and}$$

$V''(x) = -1.5x$ . The critical point occurs at  $x = 6$ , and it corresponds to the maximum volume because  $V''(x) < 0$

for  $x > 0$ . The corresponding height is  $\frac{108 - 6^2}{4(6)} = 3$  ft. The

base measures 6 ft by 6 ft, and the height is 3 ft.

48. If the dimensions are  $x$  ft by  $x$  ft by  $h$  ft, then we have

$$x^2h = 32 \text{ and so } h = \frac{32}{x^2}. \text{ Neglecting the quarter-inch}$$

thickness of the steel, the area of the steel used is

$$A(x) = x^2 + 4xh = x^2 + \frac{128}{x}. \text{ We can minimize the weight}$$

of the vat by minimizing this quantity. Now

$$A'(x) = 2x - 128x^{-2} = \frac{2}{x^2}(x^3 - 4^3) \text{ and}$$

$A''(x) = 2 + 256x^{-3}$ . The critical point occurs at  $x = 4$  and corresponds to the minimum possible area because

$$A''(x) > 0 \text{ for } x > 0. \text{ The corresponding height is } \frac{32}{4^2} = 2 \text{ ft.}$$

The base should measure 4 ft by 4 ft, and the height should be 2 ft.

49. We have  $r^2 + \left(\frac{h}{2}\right)^2 = 3$ , so  $r^2 = 3 - \frac{h^2}{4}$ . We wish to

minimize the cylinder's volume

$$V = \pi r^2 h = \pi \left( 3 - \frac{h^2}{4} \right) h = 3\pi h - \frac{\pi h^3}{4} \text{ for } 0 < h < 2\sqrt{3}.$$

$$\text{Since } \frac{dV}{dh} = 3\pi - \frac{3\pi h^2}{4} = \frac{3\pi}{4}(2 + h)(2 - h) \text{ and}$$

$$\frac{d^2V}{dh^2} = -\frac{3\pi h}{2}, \text{ the critical point occurs at } h = 2 \text{ and it}$$

corresponds to the maximum value because  $\frac{d^2V}{dh^2} < 0$  for

$h > 0$ . The corresponding value of  $r$  is  $\sqrt{3 - \frac{2^2}{4}} = \sqrt{2}$ . The

largest possible cylinder has height 2 and radius  $\sqrt{2}$ .

50. Note that, from similar cones,  $\frac{r}{6} = \frac{12 - h}{12}$ , so  $h = 12 - 2r$ .

The volume of the smaller cone is given by

$$V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi r^2(12 - 2r) = 4\pi r^2 - \frac{2\pi}{3}r^3 \text{ for } 0 < r < 6.$$

Then  $\frac{dV}{dr} = 8\pi r - 2\pi r^2 = 2\pi r(4 - r)$ , so the critical point


occurs at  $r = 4$ . This critical point corresponds to the

maximum volume because  $\frac{dV}{dr} > 0$  for  $0 < r < 4$  and

$\frac{dV}{dr} < 0$  for  $4 < r < 6$ . The smaller cone has the largest


possible value when  $r = 4$  ft and  $h = 4$  ft.





(a)  $V(x) = x(15 - 2x)(5 - x)$

(b, c) Domain:  $0 < x < 5$



The maximum volume is approximately 66.019 and it occurs when  $x \approx 1.962$  in.

(d) Note that  $V(x) = 2x^3 - 25x^2 + 75x$ ,

so  $V'(x) = 6x^2 - 50x + 75$ .

Solving  $V'(x) = 0$ , we have


$$x = \frac{50 \pm \sqrt{(-50)^2 - 4(6)(75)}}{2(6)} = \frac{50 \pm \sqrt{700}}{12}$$

$$= \frac{50 \pm 10\sqrt{7}}{12} = \frac{25 \pm 5\sqrt{7}}{6}$$

These solutions are approximately  $x \approx 1.962$  and  $x = 6.371$ , so the critical point in the appropriate domain occurs at

$$x = \frac{25 - 5\sqrt{7}}{6}$$

52.



For  $0 < x < \frac{5\pi}{3}$ , the area of the rectangle is given by

$$A(x) = (2x)(8 \cos 0.3x) = 16x \cos 0.3x.$$

Then  $A'(x) = 16x(-0.3 \sin 0.3x) + 16(\cos 0.3x)(1)$

$$= 16(\cos 0.3x - 0.3x \sin 0.3x)$$

Solving  $A'(x) = 0$  graphically, we find that the critical point occurs at  $x \approx 2.868$  and the corresponding area is approximately 29.925 square units.

53. The cost (in thousands of dollars) is given by

$$C(x) = 40x + 30(20 - y) = 40x + 600 - 30\sqrt{x^2 - 144}.$$

$$\text{Then } C'(x) = 40 - \frac{30}{2\sqrt{x^2 - 144}}(2x) = 40 - \frac{30x}{\sqrt{x^2 - 144}}.$$

Solving  $C'(x) = 0$ , we have:

$$\frac{30x}{\sqrt{x^2 - 144}} = 40$$

$$3x = 4\sqrt{x^2 - 144}$$

$$9x^2 = 16x^2 - 2304$$

$$2304 = 7x^2$$

Choose the positive solution:

$$x = +\frac{48}{\sqrt{7}} \approx 18.142 \text{ mi}$$

$$y = \sqrt{x^2 - 12^2} = \frac{36}{\sqrt{7}} \approx 13.607 \text{ mi}$$

54. The length of the track is given by  $2x + 2\pi r$ , so we have  $2x + 2\pi r = 400$  and therefore  $x = 200 - \pi r$ . Then the area of the rectangle is

$$A(r) = 2rx$$

$$= 2r(200 - \pi r)$$

$$= 400r - 2\pi r^2, \text{ for } 0 < r < \frac{200}{\pi}.$$

Therefore,  $A'(r) = 400 - 4\pi r$  and  $A''(r) = -4\pi$ , so the

critical point occurs at  $r = \frac{100}{\pi}$  m and this point

corresponds to the maximum rectangle area because  $A''(r) < 0$  for all  $r$ .

The corresponding value of  $x$  is

$$x = 200 - \pi \left( \frac{100}{\pi} \right) = 100 \text{ m}.$$

The rectangle will have the largest possible area when

$$x = 100 \text{ m and } r = \frac{100}{\pi} \text{ m}.$$

55. Assume the profit is  $k$  dollars per hundred grade B tires and  $2k$  dollars per hundred grade A tires.

Then the profit is given by

$$P(x) = 2kx + k \cdot \frac{40 - 10x}{5 - x}$$

$$= 2k \cdot \frac{(20 - 5x) + x(5 - x)}{5 - x}$$

$$= 2k \cdot \frac{20 - x^2}{5 - x}$$

$$P'(x) = 2k \cdot \frac{(5 - x)(-2x) - (20 - x^2)(-1)}{(5 - x)^2}$$

$$= 2k \cdot \frac{x^2 - 10x + 20}{(5 - x)^2}$$

## 55. Continued

The solutions of  $P'(x) = 0$  are

$$x = \frac{10 \pm \sqrt{(-10)^2 - 4(1)(20)}}{2(1)} = 5 \pm \sqrt{5}, \text{ so the solution in the}$$

appropriate domain is  $x = 5 - \sqrt{5} \approx 2.76$ .

Check the profit for the critical point and endpoints:

Critical point:  $x \approx 2.76$   $P(x) \approx 11.06k$

End points:  $x = 0$   $P(x) = 8k$

$x = 4$   $P(x) = 8k$


The highest profit is obtained when  $x \approx 2.76$  and  $y \approx 5.53$ , which corresponds to 276 grade A tires and 553 grade B tires.

56. (a) The distance between the particles is  $|f(t)|$  where

$$f(t) = -\cos t + \cos\left(t + \frac{\pi}{4}\right). \text{ Then}$$

$$f'(t) = \sin t - \sin\left(t + \frac{\pi}{4}\right)$$

Solving  $f'(t) = 0$  graphically, we obtain  $t \approx 1.178$ ,  $t \approx 4.230$ , and so on.



$[0, 2\pi]$  by  $[-2, 2]$

Alternatively,  $f'(t) = 0$  may be solved analytically as follows.

$$\begin{aligned} f'(t) &= \sin\left[t + \frac{\pi}{8} - \frac{\pi}{8}\right] - \sin\left[t + \frac{\pi}{8} + \frac{\pi}{8}\right] \\ &= \left[\sin\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} - \cos\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8}\right] \\ &\quad - \left[\sin\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} + \cos\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8}\right] \\ &= -2\sin\frac{\pi}{8}\cos\left(t + \frac{\pi}{8}\right), \end{aligned}$$


so the critical points occur when

$$\cos\left(t + \frac{\pi}{8}\right) = 0, \text{ or } t = \frac{3\pi}{8} + k\pi. \text{ At each of these values,}$$

$$f(t) = \pm 2\cos\frac{3\pi}{8} \approx \pm 0.765 \text{ units, so the maximum distance between the particles is 0.765 units.}$$

(b) Solving  $\cos t = \cos\left(t + \frac{\pi}{4}\right)$  graphically, we obtain

$$t \approx 2.749, t \approx 5.890, \text{ and so on.}$$



$[0, 2\pi]$  by  $[-2, 2]$

Alternatively, this problem may be solved analytically as follows.

$$\begin{aligned} \cos t &= \cos\left(t + \frac{\pi}{4}\right) \\ \cos\left[t + \frac{\pi}{8} - \frac{\pi}{8}\right] &= \cos\left[t + \frac{\pi}{8} + \frac{\pi}{8}\right] \\ \cos\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} + \sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} &= \cos\left(t + \frac{\pi}{8}\right)\cos\frac{\pi}{8} \\ &\quad - \sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} \\ 2\sin\left(t + \frac{\pi}{8}\right)\sin\frac{\pi}{8} &= 0 \\ \sin\left(t + \frac{\pi}{8}\right) &= 0 \\ t &= \frac{7\pi}{8} + k\pi \end{aligned}$$

The particles collide when  $t = \frac{7\pi}{8} \approx 2.749$  (plus multiples of  $\pi$  if they keep going.)

57. The dimensions will be  $x$  in. by  $10 - 2x$  in. by  $16 - 2x$  in.,


so  $V(x) = x(10 - 2x)(16 - 2x) = 4x^3 - 52x^2 + 160x$  for  $0 < x < 5$ .

Then  $V'(x) = 12x^2 - 104x + 160 = 4(x - 2)(3x - 20)$ , so the critical point in the correct domain is  $x = 2$ .

This critical point corresponds to the maximum possible volume because  $V'(x) > 0$  for  $0 < x < 2$  and  $V'(x) < 0$  for  $2 < x < 5$ . The box of largest volume has a height of 2 in. and a base measuring 6 in. by 12 in., and its volume is

$$144 \text{ in}^3.$$

Graphical support:



$[0, 5]$  by  $[-40, 160]$

## 58. Step 1:

 $r$  = radius of circle $A$  = area of circle

## Step 2:

At the instant in question,  $\frac{dr}{dt} = -\frac{2}{\pi}$  m/sec and  $r = 10$  m.

## Step 3:

We want to find  $\frac{dA}{dt}$ .

## Step 4:

$$A = \pi r^2$$

## Step 5:

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

## Step 6:

$$\frac{dA}{dt} = 2\pi(10)\left(-\frac{2}{\pi}\right) = -40$$

The area is changing at the rate of  $-40$  m<sup>2</sup>/sec.

## 59. Step 1:

 $x$  =  $x$ -coordinate of particle $y$  =  $y$ -coordinate of particle $D$  = distance from origin to particle

## Step 2:

At the instant in question,  $x = 5$  m,  $y = 12$  m,

$$\frac{dx}{dt} = -1 \text{ m/sec, and } \frac{dy}{dt} = -5 \text{ m/sec.}$$

## Step 3:

We want to find  $\frac{dD}{dt}$ .

## Step 4:

$$D = \sqrt{x^2 + y^2}$$

## Step 5:

$$\frac{dD}{dt} = \frac{1}{2\sqrt{x^2 + y^2}} \left( 2x \frac{dx}{dt} + 2y \frac{dy}{dt} \right) = \frac{x \frac{dx}{dt} + y \frac{dy}{dt}}{\sqrt{x^2 + y^2}}$$

## Step 6:

$$\frac{dD}{dt} = \frac{(5)(-1) + (12)(-5)}{\sqrt{5^2 + 12^2}} = -5 \text{ m/sec}$$

Since  $\frac{dD}{dt}$  is negative, the particle is *approaching* the origin at the *positive* rate of 5 m/sec.

## 60. Step 1:

 $x$  = edge of length of cube $V$  = volume of cube

## Step 2:

At the instant in question,

$$\frac{dV}{dt} = 1200 \text{ cm}^3/\text{min and } x = 20 \text{ cm.}$$

## Step 3:

We want to find  $\frac{dx}{dt}$ .

## Step 4:

$$V = x^3$$

## Step 5:

$$\frac{dV}{dt} = 3x^2 \frac{dx}{dt}$$

## Step 6:

$$1200 = 3(20)^2 \frac{dx}{dt}$$

$$\frac{dx}{dt} = 1 \text{ cm/min}$$

The edge length is increasing at the rate of 1 cm/min.

## 61. Step 1:

 $x$  =  $x$ -coordinate of point $y$  =  $y$ -coordinate of point $D$  = distance from origin to point

## Step 2:

At the instant in question,  $x = 3$  and  $\frac{dD}{dt} = 11$  units per sec.

## Step 3:

We want to find  $\frac{dx}{dt}$ .

## Step 4:

Since  $D^2 = x^2 + y^2$  and  $y = x^{3/2}$ , we have

$$D = \sqrt{x^2 + x^3} \text{ for } x \geq 0.$$

## Step 5:

$$\begin{aligned} \frac{dD}{dt} &= \frac{1}{2\sqrt{x^2 + x^3}} (2x + 3x^2) \frac{dx}{dt} \\ &= \frac{2x + 3x^2}{2x\sqrt{1+x}} \frac{dx}{dt} = \frac{3x+2}{2\sqrt{1+x}} \frac{dx}{dt} \end{aligned}$$

## 61. Continued

Step 6:

$$11 = \frac{3(3) + 2 \frac{dx}{dt}}{2\sqrt{4}}$$

$$\frac{dx}{dt} = 4 \text{ units per sec}$$

62. (a) Since  $\frac{h}{r} = \frac{10}{4}$ , we may write  $h = \frac{5r}{2}$  or  $r = \frac{2h}{5}$ .

(b) Step 1:

$h$  = depth of water in tank  
 $r$  = radius of surface of water  
 $V$  = volume of water in tank

Step 2:

At the instant in question,

$$\frac{dV}{dt} = -5 \text{ ft}^3/\text{min} \text{ and } h = 6 \text{ ft.}$$

Step 3:

We want to find  $-\frac{dh}{dt}$ .

Step 4:

$$V = \frac{1}{3}\pi r^2 h = \frac{4}{75}\pi h^3$$

Step 5:

$$\frac{dV}{dt} = \frac{4}{25}\pi h^2 \frac{dh}{dt}$$

Step 6:

$$-5 = \frac{4}{25}\pi(6)^2 \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{125}{144\pi} \approx -0.276 \text{ ft/min}$$

Since  $\frac{dh}{dt}$  is negative, the water level is *dropping* at the positive rate of  $\approx 0.276$  ft/min.

63. Step 1:

$r$  = radius of outer layer of cable on the spool  
 $\theta$  = clockwise angle turned by spool  
 $s$  = length of cable that has been unwound

Step 2:

At the instant in question,  $\frac{ds}{dt} = 6$  ft/sec and  $r = 1.2$  ft

Step 3:

We want to find  $\frac{d\theta}{dt}$ .

Step 4:

$$s = r\theta$$

Step 5:

Since  $r$  is essentially constant,  $\frac{ds}{dt} = r \frac{d\theta}{dt}$

Step 6:

$$6 = 1.2 \frac{d\theta}{dt}$$

$$\frac{d\theta}{dt} = 5 \text{ radians/sec}$$

The spool is turning at the rate of 5 radians per second.

64.  $a(t) = v'(t) = -g = -32 \text{ ft/sec}^2$

Since  $v(0) = 32 \text{ ft/sec}$ ,  $v(t) = s'(t) = -32t + 32$ .

Since  $s(0) = -17$ ,  $s(t) = -16t^2 + 32t - 17$ .

The shovelful of dirt reaches its maximum height when  $v(t) = 0$ , at  $t = 1$  sec. Since  $s(1) = -1$ , the shovelful of dirt is still below ground level at this time. There was not enough speed to get the dirt out of the hole. Duck!

65. We have  $V = \frac{1}{3}\pi r^2 h$ , so  $\frac{dV}{dr} = \frac{2}{3}\pi r h$  and  $dV = \frac{2}{3}\pi r h dr$ .

When the radius changes from  $a$  to  $a + dr$ , the volume change is approximately  $dV = \frac{2}{3}\pi a h dr$ .

66. (a) Let  $x$  = edge of length of cube and  $S$  = surface area of

cube. Then  $S = 6x^2$ , which means  $\frac{dS}{dx} = 12x$  and

$$dS = 12x dx. \text{ We want } |dS| \leq 0.02S, \text{ which gives}$$

$$|12x dx| \leq 0.02(6x^2) \text{ or } |dx| \leq 0.01x. \text{ The edge should be measured with an error of no more than 1\%.}$$

(b) Let  $V$  = volume of cube. Then  $V = x^3$ , which means

$$\frac{dV}{dx} = 3x^2 \text{ and } dV = 3x^2 dx. \text{ We have } |dx| \leq 0.01x,$$

$$\text{which means } |3x^2 dx| \leq 3x^2(0.01x) = 0.03V,$$

so  $|dV| \leq 0.03V$ . The volume calculation will be accurate to within approximately 3% of the correct volume.

67. Let  $C$  = circumference,  $r$  = radius,  $S$  = surface area, and  $V$  = volume.

(a) Since  $C = 2\pi r$ , we have  $\frac{dC}{dr} = 2\pi$  and so  $dC = 2\pi dr$ .

$$\text{Therefore, } \left| \frac{dC}{C} \right| = \left| \frac{2\pi dr}{2\pi r} \right| = \left| \frac{dr}{r} \right| < \frac{0.4 \text{ cm}}{10 \text{ cm}} = 0.04$$

The calculated radius will be within approximately 4% of the correct radius.

(b) Since  $S = 4\pi r^2$ , we have  $\frac{dS}{dr} = 8\pi r$  and so

$$dS = 8\pi r dr. \text{ Therefore,}$$

$$\left| \frac{dS}{S} \right| = \left| \frac{8\pi r dr}{4\pi r^2} \right| = \left| \frac{2 dr}{r} \right| \leq 2(0.04) = 0.08. \text{ The}$$

calculated surface area will be within approximately 8% of the correct surface area.

(c) Since  $V = \frac{4}{3}\pi r^3$ , we have  $\frac{dV}{dr} = 4\pi r^2$  and so

$$dV = 4\pi r^2 dr. \text{ Therefore}$$

$$\left| \frac{dV}{V} \right| = \left| \frac{4\pi r^2 dr}{\frac{4}{3}\pi r^3} \right| = \left| \frac{3 dr}{r} \right| \leq 3(0.04) = 0.12.$$

The calculated volume will be within approximately 12% of the correct volume.

68. By similar triangles, we have  $\frac{a}{6} = \frac{a+20}{h}$ , which gives

$ah = 6a + 120$ , or  $h = 6 + 120a^{-1}$ . The height of the lamp post is approximately  $6 + 120(15)^{-1} = 14$  ft. The estimated error in measuring  $a$  was

$$|da| \leq 1 \text{ in.} = \frac{1}{12} \text{ ft. Since } \frac{dh}{da} = -120a^{-2}, \text{ we have}$$

$$|dh| = |-120a^{-2} da| \leq 120(15)^{-2} \left( \frac{1}{12} \right) = \frac{2}{45} \text{ ft, so the}$$

estimated possible error is  $\pm \frac{2}{45}$  ft or  $\pm \frac{8}{15}$  in.


69.  $\frac{dy}{dx} = 2 \sin x \cos x - 3$ . Since  $\sin x$  and  $\cos x$  are both between 1 and  $-1$ , the value of  $2 \sin x \cos x$  is never greater than 2. Therefore,  $\frac{dy}{dx} \leq 2 - 3 = -1$  for all values of  $x$ .

Since  $\frac{dy}{dx}$  is always negative, the function decreases on every interval.

70. (a)  $f$  has a relative maximum at  $x = -2$ . This is where  $f'(x) = 0$ , causing  $f'$  to go from positive to negative.

(b)  $f$  has a relative minimum at  $x = 0$ . This is where  $f'(x) = 0$ , causing  $f'$  to go from negative to positive.

(c) The graph of  $f$  is concave up on  $(-1, 1)$  and on  $(2, 3)$ . These are the intervals on which the derivatives of  $f$  are increasing.



71. (a)  $A = \pi r^2$

$$\frac{dA}{dt} = 2\pi r dr$$

$$\frac{dA}{dt} = 2\pi(2) \left( \frac{1}{3} \right) = \frac{4}{3} \pi \frac{\text{in.}^2}{\text{sec}}$$

(b)  $dA = dV$

$$\frac{4}{3} \pi = \frac{1}{3} \pi r^2 dh$$

$$\frac{4}{3} \pi = \frac{1}{3} \pi (2)^2 dh$$

$$\frac{dh}{dt} = 1 \frac{\text{in.}}{\text{sec}}$$

(c)  $\frac{dA}{dh} = \frac{\frac{4}{3} \pi}{1} = \frac{4}{3} \pi \frac{\text{in.}^2}{\text{in.}}$

72. (a)  $2a + 4b = 60$

$$b = 15 - 2a$$

$$V = \pi a^2 b = \pi a^2 (15 - 2a)$$

$$\frac{dV}{da} = 30\pi a - \frac{3\pi a^2}{2}$$


$$30\pi a = \frac{3\pi a^2}{2}$$

$$a = 20$$

$$2(20) + 4b = 60$$

$$b = 5$$

(b) The sign graph for the derivative  $\frac{dV}{da} = \frac{3\pi a}{2}(20 - a)$  on the interval  $(0, 30)$  is as follows:



By the First Derivative Test, there is a maximum at  $x = 20$ .



## Chapter 5

### The Definite Integral



#### Section 5.1 Estimating with Finite Sums

(pp. 263-273)

#### Exploration 1 Which RAM is the Biggest?



LRAM > MRAM > RRAM



MRAM > RRAM > LRAM

3. RRAM > MRAM > LRAM, because the heights of the rectangles increase as you move toward the right under an increasing function.

4. LRAM > MRAM > RRAM, because the heights of the rectangles decrease as you move toward the right under a decreasing function.

#### Quick Review 5.1

- $80 \text{ mph} \cdot 5 \text{ hr} = 400 \text{ mi}$
- $48 \text{ mph} \cdot 3 \text{ hr} = 144 \text{ mi}$
- $10 \text{ ft/sec}^2 \cdot 10 \text{ sec} = 100 \text{ ft/sec}$   
 $100 \text{ ft/sec} \cdot \frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{3600 \text{ sec}}{1 \text{ h}} \approx 68.18 \text{ mph}$
- $300,000 \text{ km/sec} \cdot \frac{3600 \text{ sec}}{1 \text{ hr}} \cdot \frac{24 \text{ hr}}{1 \text{ day}} \cdot \frac{365 \text{ days}}{1 \text{ yr}} \cdot 1 \text{ yr}$   
 $\approx 9.46 \times 10^{12} \text{ km}$
- $(6 \text{ mph})(3 \text{ h}) + (5 \text{ mph})(2 \text{ h}) = 18 \text{ mi} + 10 \text{ mi} = 28 \text{ mi}$
- $20 \text{ gal/min} \cdot 1 \text{ h} \cdot \frac{60 \text{ min}}{1 \text{ h}} = 1200 \text{ gal}$
- $(-1^\circ\text{C/h})(12 \text{ h}) + (1.5^\circ\text{C})(6 \text{ h}) = -3^\circ\text{C}$
- $300 \text{ ft}^3/\text{sec} \cdot \frac{3600 \text{ sec}}{1 \text{ h}} \cdot \frac{24 \text{ h}}{1 \text{ day}} \cdot 1 \text{ day} = 25,920,000 \text{ ft}^3$
- $350 \text{ people/mi}^2 \cdot 50 \text{ mi}^2 = 17,500 \text{ people}$
- $70 \text{ times/sec} \cdot \frac{3600 \text{ sec}}{1 \text{ h}} \cdot 1 \text{ h} \cdot 0.7 = 176,400 \text{ times}$

#### Section 5.1 Exercises

- Since  $v(t) = 5$  is a straight line, compute the area under the curve.  
 $x = (t) v(t) = (4)(5) = 20$
- Since  $v(t) = 2t + 1$  creates a trapezoid with the  $x$ -axis, compute the area of the curve under the trapezoid.  
 $A = \frac{h}{2}(a + b)$   
 $a = t = 0 = v(0) = 2(0) + 1 = 1$   
 $b = t = 4 = v(4) = 2(4) + 1 = 9$   
 $h = 4$   
 $A = \frac{4}{2}(9 + 1) = 20$
- Each rectangle has base 1. The height of each rectangle is found by using the points  $t = (0.5, 1.5, 2.5, 3.5)$  in the equation  $v(t) = t^2 + 1$ . The area under the curve is approximately  $1\left(\frac{5}{4} + \frac{13}{4} + \frac{29}{4} + \frac{53}{4}\right) = 25$ , so the particle is close to  $x = 25$ .
- Each rectangle has base 1. The height of each rectangle is found by using the points  $y = (0.5, 1.5, 2.5, 3.5, 4.5)$  in the equation  $v(t) = t^2 + 1$ . The area under the curve is approximately  $1\left(\frac{5}{4} + \frac{13}{4} + \frac{29}{4} + \frac{53}{4} + \frac{85}{4}\right) = 46.25$ , so the particle is close to  $x = 46.25$ .