

Extra Practice Chapter 4

Let f be a twice-differentiable function with $f(0) = 4$. The derivative of f is given by $f'(x) = \sin(x^2 - 2x + 1)$ for $-2 \leq x \leq 2$.

- (a) Find all values of x in the interval $-2 < x < 2$ at which f has a critical point. Classify each as the location of a relative minimum, a relative maximum, or neither. Justify your answers.
- (b) Use the line tangent to the graph of f at $x = 0$ to approximate $f(0.25)$.
- (c) On the interval $0 \leq x \leq 0.25$, $f'(x) > 0$ and $f''(x) < 0$. Is the approximation found in part (b) an overestimate or an underestimate for $f(0.25)$? Give a reason for your answer.
- (d) Using the Mean Value Theorem, explain why the average rate of change of f over the interval $-2 \leq x \leq 2$ cannot equal 1.25.

$\lim_{x \rightarrow 0} \frac{x^2}{1 - \cos x}$ is

- A -2
- B 0
- C 1
- D 2
- E nonexistent

Let f be a twice-differentiable function with $f(1.5) = 3$. The derivative of f is given by $f'(x) = (7x - 19) \sin(x^2 - 4x + 4)$ for $1 \leq x \leq 4$.

(a) Find all values of x in the interval $1 < x < 4$ at which f has a critical point. Classify each as the location of a relative minimum, a relative maximum, or neither. Justify your answers.

(b) Use the line tangent to the graph of f at $x = 1.5$ to approximate $f(1.8)$.

On the interval $1.5 \leq x \leq 1.8$, $f'(x) < 0$ and $f''(x) > 0$. Is the approximation found in part (b) an overestimate or an underestimate for $f(1.8)$? Give a reason for your answer.

Using the Mean Value Theorem, explain why the average rate of change of f over the interval $1 \leq x \leq 4$ cannot equal 6.5.

$\lim_{x \rightarrow \infty} \frac{x^3}{e^{3x}}$ is

A 0

B $\frac{2}{9}$

C $\frac{2}{3}$

D 1

E infinite

Functions f , g , and h are twice-differentiable functions with $g(2) = h(2) = 4$. The line $y = 4 + \frac{2}{3}(x - 2)$ is tangent to both the graph of g at $x = 2$ and the graph of h at $x = 2$.

a) Find $h'(2)$.

b) Let a be the function given by $a(x) = 3x^3h(x)$. Write an expression for $a'(x)$. Find $a'(2)$.

c) The function h satisfies $h(x) = \frac{x^2 - 4}{1 - (f(x))^4}$ for $x \neq 2$. It is known that $\lim_{x \rightarrow 2} h(x)$ can be evaluated using L'Hospital's Rule. Use $\lim_{x \rightarrow 2} h(x)$ to find $f'(2)$ and $f''(2)$. Show the work that leads to your answers.

d) It is known that $g(x) \leq h(x)$ for $1 < x < 3$. Let k be a function satisfying $g(x) \leq k(x) \leq h(x)$ for $1 < x < 3$. Is k continuous at $x = 2$? Justify your answer.

$$\lim_{h \rightarrow 0} \frac{e^{(2+h)} - e^2}{h} =$$

A 0

B 1

C $2e$

D e^2

E $2e^2$

x	$f(x)$	$f'(x)$	$f''(x)$	$f'''(x)$
2	0	0	5	7

The third derivative of the function f is continuous on the interval $(0,4)$. Values for f and its first three derivatives at $x=2$ are given in the table above. What is

$$\lim_{x \rightarrow 2} \frac{f(x)}{(x-2)^2} ?$$

- (A) 0
- (B) $\frac{5}{2}$
- (C) 5
- (D) 7
- (E) The limit does not exist.

x	2	3	5	8	13
$f(x)$	1	4	-2	3	6

Let f be a function that is twice differentiable for all real numbers. The table above gives values of f for selected points in the closed interval $2 \leq x \leq 13$.

Suppose $f(5)=3$ and $f'(x)<0$ for all x in the closed interval $5 \leq x \leq 8$. Use the line tangent to the graph of f at $x=5$ to show that $f(7) \leq 4$. Use the secant line for the graph of f on $5 \leq x \leq 8$ to show that $f(7) \geq 4/3$.

A differentiable function f has the property that $f(5) = 3$ and $f'(5) = 4$. What is the estimate for $f(4.8)$ using the local linear approximation for f at $x = 5$?

- (A) 2.2
- (B) 2.8
- (C) 3.4
- (D) 3.8
- (E) 4.6


x	-1.5	-1.0	-0.5	0	0.5	1.0	1.5
$f(x)$	-1	-4	-6	-7	-6	-4	-1
$f'(x)$	-7	-5	-3	0	3	5	7

Let f be a function that is differentiable for all real numbers. The table above gives the values of f and its derivative f' for selected points x in the closed interval $-1.5 \leq x \leq 1.5$. The second derivative of f has the property that $f''(x) > 0$ for $-1.5 \leq x \leq 1.5$.

Write an equation of the line tangent to the graph of f at the point where $x = 1$. Use this line to approximate the value of $f(1.2)$. Is this approximation greater than or less than the actual value of $f(1.2)$? Give a reason for your answer.

The function f is continuous for $-2 \leq x \leq 2$ and $f(-2) = f(2) = 0$. If there is no c , where $-2 < c < 2$ for which $f'(c) = 0$, which of the following statements must be true?

- (A) For $-2 < k < 2$, $f(k) > 0$
- (B) For $-2 < k < 2$, $f(k) < 0$
- (C) For $-2 < k < 2$, $f(k)$ exists.
- (D) For $-2 < k < 2$, $f(k)$ exists, but f is not continuous.
- (E) For some k , where $-2 < k < 2$, $f(k)$ does not exist.



The figure above shows the graph of the differentiable function f for $1 \leq x \leq 5$. Which of the following could be the x -coordinate of a point at which the line tangent to the graph of f is parallel to the secant line through the points $(1, f(1))$ and $(5, f(5))$?

- (A) r
- (B) s
- (C) u
- (D) w
- (E) There is no such point.

t (sec)	0	15	25	30	35	50	60
$v(t)$ (ft/sec)	-20	-30	-20	-14	-10	0	10
$a(t)$ (ft/sec ²)	1	5	2	1	2	4	2


A car travels on a straight track. During the time interval $0 \leq t \leq 60$ seconds, the car's velocity v , measured in feet per second, and acceleration a , measured in feet per second per second, are continuous functions. The table above shows selected values of these functions.

For $0 < t < 60$, must there be a time t when $a(t) = 0$? Justify your answer.

t (minutes)	0	1	2	3	4	5	6
$C(t)$ (ounces)	0	5.3	8.8	11.2	12.8	13.8	14.5


Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time t , $0 \leq t \leq 6$, is given by a differentiable function C , where t is measured in minutes. Selected values of $C(t)$, measured in ounces, are given in the table above.

Is there a time t , $2 \leq t \leq 4$, at which $C'(t) = 2$? Justify your answer.



Let g be a continuous function with $g(2) = 5$. The graph of the piecewise-linear function g' , the derivative of g , is shown above for $-3 \leq x \leq 7$.

Find the average rate of change of $g(x)$ on the interval $-3 \leq x \leq 7$. Does the Mean Value Theorem applied on the interval $-3 \leq x \leq 7$ guarantee a value of c , for $-3 < c < 7$, such that $g'(c)$ is equal to this average rate of change? Why or why not?



The function f shown in the figure above is continuous on the closed interval $[0, 12]$ and differentiable on the open interval $(0, 12)$. Based on the graph, what are all values of x that satisfy the conclusion of the Mean Value Theorem applied to f on the closed interval $[0, 12]$?

- (A) 4.5 only because this is the value where $f(x)$ equals the average rate of change of f on $[0, 12]$.
- (B) 3 and 8 because these are the values where $f'(x) = 0$ on $[0, 12]$.
- (C) 2 and 9 only because these are the values where the instantaneous rate of change of f at those values is equal to the average rate of change of f on $[0, 12]$.
- (D) 2, 4.5, and 9 because these are the values where either the instantaneous rate of change of f at the value is equal to the average rate of change of f on $[0, 12]$ or the value of $f(x)$ is equal to the average rate of change of f on $[0, 12]$.