

Step 2: Definite integral. The amount consumed from $t = 2$ to $t = 4$ is the limit of these sums as the norms of the partitions go to zero.

$$\int_2^4 C(t) dt = \int_2^4 (2.2 + 1.1t) dt \text{ million bushels}$$

Step 3: Evaluate. Evaluating numerically, we obtain

$$\text{NINT}(2.2 + 1.1t, t, 2, 4) \approx 7.066 \text{ million bushels.}$$

Now try Exercise 21

Net Change from Data

Many real applications begin with data, not a fully modeled function. In the next example, we are given data on the rate at which a pump operates in consecutive 5-minute intervals and asked to find the total amount pumped.

Table 7.1 Pumping Rates

Time (min)	Rate (gal/min)
0	58
5	60
10	65
15	64
20	58
25	57
30	55
35	55
40	59
45	60
50	60
55	63
60	63

EXAMPLE 6 Finding Gallons Pumped from Rate Data

A pump connected to a generator operates at a varying rate, depending on how much power is being drawn from the generator to operate other machinery. The rate (gallons per minute) at which the pump operates is recorded at 5-minute intervals for one hour as shown in Table 7.1. How many gallons were pumped during that hour?

SOLUTION

Let $R(t)$, $0 \leq t \leq 60$, be the pumping rate as a continuous function of time for the hour. We can partition the hour into short subintervals of length Δt on which the rate is nearly constant and form the sum $\sum R(t_k) \Delta t$ as an approximation to the amount pumped during the hour. This reveals the integral formula for the number of gallons pumped to be

$$\text{Gallons pumped} = \int_0^{60} R(t) dt.$$

We have no formula for R in this instance, but the 13 equally spaced values in Table 7.1 enable us to estimate the integral with the Trapezoidal Rule:

$$\begin{aligned} \int_0^{60} R(t) dt &\approx \frac{60}{2 \cdot 12} [58 + 2(60) + 2(65) + \cdots + 2(63) + 63] \\ &= 3582.5. \end{aligned}$$

The total amount pumped during the hour is about 3580 gal.

Now try Exercise 21

Work

In everyday life, *work* means an activity that requires muscular or mental effort. In science, the term refers specifically to a force acting on a body and the body's subsequent displacement. When a body moves a distance d along a straight line as a result of the action of a force of constant magnitude F in the direction of motion, the **work** done by the force is

$$W = Fd.$$

The equation $W = Fd$ is the **constant-force formula** for work.

The units of work are force \times distance. In the metric system, the unit is the newton-meter, which, for historical reasons, is called a joule (see margin note). In the U.S. customary system, the most common unit of work is the **foot-pound**.

Joules

The joule, abbreviated J and pronounced "jewel," is named after the English physicist James Prescott Joule (1818–1889). The defining equation is

$$1 \text{ joule} = (1 \text{ newton})(1 \text{ meter}).$$

In symbols, $1 \text{ J} = 1 \text{ N} \cdot \text{m}$.

It takes a force of about 1 N to lift an apple from a table. If you lift it 1 m you have done about 1 J of work on the apple. If you eat the apple, you will have consumed about 80 food calories, the heat equivalent of nearly 335,000 joules. If this energy were directly useful for mechanical work (it's not), it would enable you to lift 335,000 more apples up 1 m.

Hooke's Law for springs says that the force it takes to stretch or compress a spring x units from its natural (unstressed) length is a constant times x . In symbols,

$$F = kx,$$

where k , measured in force units per unit length, is a characteristic of the spring called the **force constant**.

EXAMPLE 7 A Bit of Work

It takes a force of 10 N to stretch a spring 2 m beyond its natural length. How much work is done in stretching the spring 4 m from its natural length?

SOLUTION

We let $F(x)$ represent the force in newtons required to stretch the spring x meters from its natural length. By Hooke's Law, $F(x) = kx$ for some constant k . We are told that

$$F(2) = 10 = k \cdot 2, \quad \text{The force required to stretch the spring 2 m is 10 newtons.}$$

so $k = 5 \text{ N/m}$ and $F(x) = 5x$ for this particular spring.

We construct an integral for the work done in applying F over the interval from $x = 0$ to $x = 4$.

Step 1: Riemann sum. We partition the interval into subintervals on each of which F is so nearly constant that we can apply the constant-force formula for work. If x_k is any point in the k th subinterval, the value of F throughout the interval is approximately $F(x_k) = 5x_k$. The work done by F across the interval is approximately $5x_k \Delta x$, where Δx is the length of the interval. The sum

$$\sum F(x_k) \Delta x = \sum 5x_k \Delta x$$

approximates the work done by F from $x = 0$ to $x = 4$.

Steps 2 and 3: Integrate. The limit of these sums as the norms of the partitions go to zero is

$$\int_0^4 F(x) dx = \int_0^4 5x dx = 5 \left[\frac{x^2}{2} \right]_0^4 = 40 \text{ N} \cdot \text{m}.$$


Now try Exercise 29.

We will revisit work in Section 7.5.

Quick Review 7.1 (For help, go to Section 1.2.)

In Exercises 1–10, find all values of x (if any) at which the function changes sign on the given interval. Sketch a number line graph of the interval, and indicate the sign of the function on each subinterval.

Example: $f(x) = x^2 - 1$ on $[-2, 3]$



Changes sign at $x = \pm 1$.

1. $\sin 2x$ on $[-3, 2]$

2. $x^2 - 3x + 2$ on $[-2, 4]$

3. $x^2 - 2x + 3$ on $[-4, 2]$

4. $2x^3 - 3x^2 + 1$ on $[-2, 2]$

5. $x \cos 2x$ on $[0, 4]$

6. xe^{-x} on $[0, \infty)$

7. $\frac{x}{x^2 + 1}$ on $[-5, 30]$

8. $\frac{x^2 - 2}{x^2 - 4}$ on $[-3, 3]$

9. $\sec(1 + \sqrt{1 - \sin^2 x})$ on $(-\infty, \infty)$

10. $\sin(1/x)$ on $[0.1, 0.2]$

Section 7.1 Exercises

In Exercises 1–8, the function $v(t)$ is the velocity in m/sec of a particle moving along the x -axis. Use analytic methods to do each of the following:

- (a) Determine when the particle is moving to the right, to the left, and stopped.
- (b) Find the particle's displacement for the given time interval. If $x(0) = 3$, what is the particle's final position?
- (c) Find the total distance traveled by the particle.

1. $v(t) = 5 \cos t, \quad 0 \leq t \leq 2\pi$

2. $v(t) = 6 \sin 3t, \quad 0 \leq t \leq \pi/2$

3. $v(t) = 49 - 9.8t, \quad 0 \leq t \leq 10$

4. $v(t) = 6t^2 - 18t + 12, \quad 0 \leq t \leq 2$

5. $v(t) = 5 \sin^2 t \cos t, \quad 0 \leq t \leq 2\pi$

6. $v(t) = \sqrt{4-t}, \quad 0 \leq t \leq 4$

7. $v(t) = e^{\sin t} \cos t, \quad 0 \leq t \leq 2\pi$

8. $v(t) = \frac{t}{1+t^2}, \quad 0 \leq t \leq 3$

9. An automobile accelerates from rest at $1 + 3\sqrt{t}$ mph/sec for 9 seconds.

- (a) What is its velocity after 9 seconds?
- (b) How far does it travel in those 9 seconds?


10. A particle travels with velocity $v(t) = (t - 2) \sin t$ m/sec for $0 \leq t \leq 4$ sec.

- (a) What is the particle's displacement?
- (b) What is the total distance traveled?

11. **Projectile** Recall that the acceleration due to Earth's gravity is 32 ft/sec^2 . From ground level, a projectile is fired straight upward with velocity 90 feet per second.

- (a) What is its velocity after 3 seconds?
- (b) When does it hit the ground?
- (c) When it hits the ground, what is the net distance it has traveled?
- (d) When it hits the ground, what is the total distance it has traveled?





In Exercises 12–16, a particle moves along the x -axis (units in cm). Its initial position at $t = 0$ sec is $x(0) = 15$. The figure shows the graph of the particle's velocity $v(t)$. The numbers are the areas of the enclosed regions.



- 12. What is the particle's displacement between $t = 0$ and $t = c$?
- 13. What is the total distance traveled by the particle in the same time period?
- 14. Give the positions of the particle at times $a, b,$ and c .
- 15. Approximately where does the particle achieve its greatest positive acceleration on the interval $[0, b]$?
- 16. Approximately where does the particle achieve its greatest positive acceleration on the interval $[0, c]$?

In Exercises 17–20, the graph of the velocity of a particle moving on the x -axis is given. The particle starts at $x = 2$ when $t = 0$.

- (a) Find where the particle is at the end of the trip.
- (b) Find the total distance traveled by the particle.



- 21. **U.S. Oil Consumption** The rate of consumption of oil in the United States during the 1980s (in billions of barrels per year) is modeled by the function $C = 27.08 \cdot e^{t/25}$, where t is the number of years after January 1, 1980. Find the total consumption of oil in the United States from January 1, 1980 to January 1, 1990.
- 22. **Home Electricity Use** The rate at which your home consumes electricity is measured in kilowatts. If your home consumes electricity at the rate of 1 kilowatt for 1 hour, you will be charged

for 1 "kilowatt-hour" of electricity. Suppose that the average consumption rate for a certain home is modeled by the function $C(t) = 3.9 - 2.4 \sin(\pi t/12)$, where $C(t)$ is measured in kilowatts and t is the number of hours past midnight. Find the average daily consumption for this home, measured in kilowatt-hours.

Population Density Population density measures the number of people per square mile inhabiting a given living area. Washerton's population density, which decreases as you move away from the city center, can be approximated by the function $10,000(2 - r)$ at a distance r miles from the city center.

- (a) If the population density approaches zero at the edge of the city, what is the city's radius?
- (b) A thin ring around the center of the city has thickness Δr and radius r . If you straighten it out, it suggests a rectangular strip. Approximately what is its area?
- (c) **Writing to Learn** Explain why the population of the ring in part (b) is approximately

$$10,000(2 - r)(2\pi r) \Delta r.$$

(d) Estimate the total population of Washerton by setting up and evaluating a definite integral.

Oil Flow Oil flows through a cylindrical pipe of radius 3 inches, but friction from the pipe slows the flow toward the outer edge. The speed at which the oil flows at a distance r inches from the center is $8(10 - r^2)$ inches per second.

(a) In a plane cross section of the pipe, a thin ring with thickness Δr at a distance r inches from the center approximates a rectangular strip when you straighten it out. What is the area of the strip (and hence the approximate area of the ring)?

(b) Explain why we know that oil passes through this ring at approximately $8(10 - r^2)(2\pi r) \Delta r$ cubic inches per second.

(c) Set up and evaluate a definite integral that will give the rate (in cubic inches per second) at which oil is flowing through the pipe.

Group Activity Bagel Sales From 1995 to 2005, the Königsberg Bakery noticed a consistent increase in annual sales of its bagels. The annual sales (in thousands of bagels) are shown below.

Year	Sales (thousands)
1995	5
1996	8.9
1997	16
1998	26.3
1999	39.8
2000	56.5
2001	76.4
2002	99.5
2003	125.8
2004	155.3
2005	188

- (a) What was the total number of bagels sold over the 11-year period? (This is not a calculus question!)
- (b) Use quadratic regression to model the annual bagel sales (in thousands) as a function $B(x)$, where x is the number of years after 1995.
- (c) Integrate $B(x)$ over the interval $[0, 11]$ to find total bagel sales for the 11-year period.
- (d) Explain graphically why the answer in part (a) is smaller than the answer in part (c).

26. **Group Activity** (Continuation of Exercise 25)

(a) Integrate $B(x)$ over the interval $[-0.5, 10.5]$ to find total bagel sales for the 11-year period.

(b) Explain graphically why the answer in part (a) is better than the answer in 25(c).

27. **Filling Milk Cartons** A machine fills milk cartons with milk at an approximately constant rate, but backups along the assembly line cause some variation. The rates (in cases per hour) are recorded at hourly intervals during a 10-hour period, from 8:00 A.M. to 6:00 P.M.

Time	Rate (cases/h)
8	120
9	110
10	115
11	115
12	119
1	120
2	120
3	115
4	112
5	110
6	121

Use the Trapezoidal Rule with $n = 10$ to determine approximately how many cases of milk were filled by the machine over the 10-hour period.

28. **Writing to Learn** As a school project, Anna accompanies her mother on a trip to the grocery store and keeps a log of the car's speed at 10-second intervals. Explain how she can use the data to estimate the distance from her home to the store. What is the connection between this process and the definite integral?

29. **Hooke's Law** A certain spring requires a force of 6 N to stretch it 3 cm beyond its natural length.

- (a) What force would be required to stretch the string 9 cm beyond its natural length?
- (b) What would be the work done in stretching the string 9 cm beyond its natural length?


30. **Hooke's Law** Hooke's Law also applies to compressing springs; that is, it requires a force of kx to compress a spring a distance x from its natural length. Suppose a 10,000-lb force compressed a spring from its natural length of 12 inches to a length of 11 inches. How much work was done in compressing the spring

- (a) the first half-inch?
- (b) the second half-inch?


Standardized Test Questions

Calculator You may use a graphing calculator to solve the following problems.

- 31. True or False** The figure below shows the velocity for a particle moving along the x -axis. The displacement for this particle is negative. Justify your answer.



- 32. True or False** If the velocity of a particle moving along the x -axis is always positive, then the displacement is equal to the total distance traveled. Justify your answer.
- 33. Multiple Choice** The graph below shows the rate at which water is pumped from a storage tank. Approximate the total gallons of water pumped from the tank in 24 hours.
- (A) 600 (B) 2400 (C) 3600 (D) 4200 (E) 4800



- 34. Multiple Choice** The data for the acceleration $a(t)$ of a car from 0 to 15 seconds are given in the table below. If the velocity at $t = 0$ is 5 ft/sec, which of the following gives the approximate velocity at $t = 15$ using the Trapezoidal Rule?
- (A) 47 ft/sec (B) 52 ft/sec (C) 120 ft/sec
(D) 125 ft/sec (E) 141 ft/sec

t (sec)	0	3	6	9	12	15
$a(t)$ (ft/sec ²)	4	8	6	9	10	10

- 35. Multiple Choice** The rate at which customers arrive at a counter to be served is modeled by the function F defined by $F(t) = 12 + 6 \cos\left(\frac{t}{\pi}\right)$ for $0 \leq t \leq 60$, where $F(t)$ is measured in customers per minute and t is measured in minutes. To the nearest whole number, how many customers arrive at the counter over the 60-minute period?
- (A) 720 (B) 725 (C) 732 (D) 744 (E) 756

- 36. Multiple Choice** Pollution is being removed from a lake at a rate modeled by the function $y = 20e^{-0.5t}$ tons/yr, where t is the number of years since 1995. Estimate the amount of pollution moved from the lake between 1995 and 2005. Round your answer to the nearest ton.
- (A) 40 (B) 47 (C) 56 (D) 61 (E) 71

Extending the Ideas


- 37. Inflation** Although the economy is continuously changing, we analyze it with discrete measurements. The following table records the *annual* inflation rate as measured each month for 13 consecutive months. Use the Trapezoidal Rule with $n = 12$ to find the overall inflation rate for the year.

Month	Annual Rate
January	0.04
February	0.04
March	0.05
April	0.06
May	0.05
June	0.04
July	0.04
August	0.05
September	0.04
October	0.06
November	0.06
December	0.05
January	0.05

- 38. Inflation Rate** The table below shows the *monthly* inflation rate (in *thousandths*) for energy prices for thirteen consecutive months. Use the Trapezoidal Rule with $n = 12$ to approximate the *annual* inflation rate for the 12-month period running from the middle of the first month to the middle of the last month.

Month	Monthly Rate (in thousandths)
January	3.6
February	4.0
March	3.1
April	2.8
May	2.8
June	3.2
July	3.3
August	3.1
September	3.2
October	3.4
November	3.4
December	3.9
January	4.0

- Center of Mass** Suppose we have a finite collection of masses in the coordinate plane, the mass m_k located at the point (x_k, y_k) as shown in the figure.



Each mass m_k has **moment $m_k y_k$ with respect to the x -axis** and **moment $m_k x_k$ about the y -axis**. The moments of the entire system with respect to the two axes are

$$\text{Moment about } x\text{-axis: } M_x = \sum m_k y_k.$$

$$\text{Moment about } y\text{-axis: } M_y = \sum m_k x_k.$$

The **center of mass** is (\bar{x}, \bar{y}) where

$$\bar{x} = \frac{M_y}{M} = \frac{\sum m_k x_k}{\sum m_k} \quad \text{and} \quad \bar{y} = \frac{M_x}{M} = \frac{\sum m_k y_k}{\sum m_k}.$$

Suppose we have a thin, flat plate occupying a region in the plane.

- (a) Imagine the region cut into thin strips parallel to the y -axis. Show that

$$\bar{x} = \frac{\int x \, dm}{\int dm},$$

where $dm = \delta \, dA$, δ = density (mass per unit area), and A = area of the region.

- (b) Imagine the region cut into thin strips parallel to the x -axis. Show that

$$\bar{y} = \frac{\int y \, dm}{\int dm},$$

where $dm = \delta \, dA$, δ = density, and A = area of the region.

In Exercises 40 and 41, use Exercise 39 to find the center of mass of the region with given density.

- 40.** the region bounded by the parabola $y = x^2$ and the line $y = 4$ with constant density δ
- 41.** the region bounded by the lines $y = x$, $y = -x$, $x = 2$ with constant density δ




Figure 7.12 The region in Example 6.




Figure 7.13 If we integrate with respect to x in Example 6, we must split the region at $x = a$.




Figure 7.14 The area of the blue region is the area under the parabola $y = \sqrt{x}$ minus the area of the triangle. (Example 7)

We must still be careful to subtract the lower number from the higher number when forming the integrand. In this case, the higher numbers are the higher x -values, which are on the line $x = y + 2$ because the line lies to the right of the parabola. So,

$$\text{Area of } R = \int_0^2 (y + 2 - y^2) dy = \left[\frac{y^2}{2} + 2y - \frac{y^3}{3} \right]_0^2 = \frac{10}{3} \text{ units squared.}$$

Now try Exercise 1

EXAMPLE 6 Making the Choice

Find the area of the region enclosed by the graphs of $y = x^3$ and $x = y^2 - 2$.

SOLUTION

We can produce a graph of the region on a calculator by graphing the three curves $y = x^3, y = \sqrt{x+2}$, and $y = -\sqrt{x+2}$ (Figure 7.12).

This conveniently gives us all of our bounding curves as functions of x . If we integrate in terms of x , however, we need to split the region at $x = a$ (Figure 7.13).

On the other hand, we can integrate from $y = b$ to $y = d$ and handle the entire region at once. We solve the cubic for x in terms of y :

$$y = x^3 \text{ becomes } x = y^{1/3}.$$

To find the limits of integration, we solve $y^{1/3} = y^2 - 2$. It is easy to see that the lower limit is $b = -1$, but a calculator is needed to find that the upper limit $d = 1.793003715$. We store this value as D .

The cubic lies to the right of the parabola, so

$$\text{Area} = \text{NINT} (y^{1/3} - (y^2 - 2), y, -1, D) = 4.214939673.$$

The area is about 4.21.

Now try Exercise 2

Saving Time with Geometry Formulas

Here is another way to handle Example 4.

EXAMPLE 7 Using Geometry

Find the area of the region in Example 4 by subtracting the area of the triangular region from the area under the square root curve.

SOLUTION

Figure 7.14 illustrates the strategy, which enables us to integrate with respect to x without splitting the region.

$$\text{Area} = \int_0^4 \sqrt{x} dx - \frac{1}{2} (2)(2) = \left[\frac{2}{3} x^{3/2} \right]_0^4 - 2 = \frac{10}{3} \text{ units squared}$$

Now try Exercise 3

The moral behind Examples 4, 5, and 7 is that you often have options for finding the area of a region, some of which may be easier than others. You can integrate with respect to x or with respect to y , you can partition the region into subregions, and sometimes you can even use traditional geometry formulas. Sketch the region first and take a moment to determine the best way to proceed.

Quick Review 7.2 (For help, go to Sections 1.2 and 5.1.)

In Exercises 1–5, find the area between the x -axis and the graph of the given function over the given interval.

1. $y = \sin x$ over $[0, \pi]$

2. $y = e^{2x}$ over $[0, 1]$

3. $y = \sec^2 x$ over $[-\pi/4, \pi/4]$

4. $y = 4x - x^3$ over $[0, 2]$

5. $y = \sqrt{9 - x^2}$ over $[-3, 3]$

In Exercises 6–10, find the x - and y -coordinates of all points where the graphs of the given functions intersect. If the curves never intersect, write “NI.”

6. $y = x^2 - 4x$ and $y = x + 6$

7. $y = e^x$ and $y = x + 1$







8. $y = x^2 - \pi x$ and $y = \sin x$

9. $y = \frac{2x}{x^2 + 1}$ and $y = x^3$

10. $y = \sin x$ and $y = x^3$

Section 7.2 Exercises



In Exercises 1–6, find the area of the shaded region analytically.



In Exercises 7 and 8, use a calculator to find the area of the region enclosed by the graphs of the two functions.

7. $y = \sin x, y = 1 - x^2$ 8. $y = \cos(2x), y = x^2 - 2$



In Exercises 9 and 10, find the area of the shaded region analytically.



In Exercises 11 and 12, find the area enclosed by the graphs of the two curves by integrating with respect to y .

11. $y^2 = x + 1, y^2 = 3 - x$ 12. $y^2 = x + 3, y = 2x$

In Exercises 13 and 14, find the total shaded area.



In Exercises 15–34, find the area of the regions enclosed by the lines and curves.

15. $y = x^2 - 2$ and $y = 2$
 16. $y = 2x - x^2$ and $y = -3$
 17. $y = 7 - 2x^2$ and $y = x^2 + 4$


18. $y = x^4 - 4x^2 + 4$ and $y = x^2$
 19. $y = x\sqrt{a^2 - x^2}$, $a > 0$, and $y = 0$
 20. $y = \sqrt{|x|}$ and $5y = x + 6$
 (How many intersection points are there?)

21. $y = |x^2 - 4|$ and $y = (x^2/2) + 4$
 22. $x = y^2$ and $x = y + 2$
 23. $y^2 - 4x = 4$ and $4x - y = 16$
 24. $x - y^2 = 0$ and $x + 2y^2 = 3$
 25. $x + y^2 = 0$ and $x + 3y^2 = 2$
 26. $4x^2 + y = 4$ and $x^4 - y = 1$
 27. $x + y^2 = 3$ and $4x + y^2 = 0$
 28. $y = 2 \sin x$ and $y = \sin 2x$, $0 \leq x \leq \pi$
 29. $y = 8 \cos x$ and $y = \sec^2 x$, $-\pi/3 \leq x \leq \pi/3$
 30. $y = \cos(\pi x/2)$ and $y = 1 - x^2$
 31. $y = \sin(\pi x/2)$ and $y = x$
 32. $y = \sec^2 x$, $y = \tan^2 x$, $x = -\pi/4$, $x = \pi/4$
 33. $x = \tan^2 y$ and $x = -\tan^2 y$, $-\pi/4 \leq y \leq \pi/4$
 34. $x = 3 \sin y \sqrt{\cos y}$ and $x = 0$, $0 \leq y \leq \pi/2$

In Exercises 35 and 36, find the area of the region by subtracting the area of a triangular region from the area of a larger region.

35. The region on or above the x -axis bounded by the curves $y^2 = x + 3$ and $y = 2x$.
36. The region on or above the x -axis bounded by the curves $y = 4 - x^2$ and $y = 3x$.
37. Find the area of the propeller-shaped region enclosed by the curve $x - y^3 = 0$ and the line $x - y = 0$.
38. Find the area of the region in the first quadrant bounded by the line $y = x$, the line $x = 2$, the curve $y = 1/x^2$, and the x -axis.
39. Find the area of the “triangular” region in the first quadrant bounded on the left by the y -axis and on the right by the curves $y = \sin x$ and $y = \cos x$.
40. Find the area of the region between the curve $y = 3 - x^2$ and the line $y = -1$ by integrating with respect to (a) x , (b) y .
41. The region bounded below by the parabola $y = x^2$ and above by the line $y = 4$ is to be partitioned into two subsections of equal area by cutting across it with the horizontal line $y = c$.
- (a) Sketch the region and draw a line $y = c$ across it that looks about right. In terms of c , what are the coordinates of the points where the line and parabola intersect? Add them to your figure.
- (b) Find c by integrating with respect to y . (This puts c in the limits of integration.)
- (c) Find c by integrating with respect to x . (This puts c into the integrand as well.)
42. Find the area of the region in the first quadrant bounded on the left by the y -axis, below by the line $y = x/4$, above left by the curve $y = 1 + \sqrt{x}$, and above right by the curve $y = 2/\sqrt{x}$.

The figure here shows triangle AOC inscribed in the region cut from the parabola $y = x^2$ by the line $y = a^2$. Find the limit of the ratio of the area of the triangle to the area of the parabolic region as a approaches zero.




Suppose the area of the region between the graph of a positive continuous function f and the x -axis from $x = a$ to $x = b$ is 4 square units. Find the area between the curves $y = f(x)$ and $y = 2f(x)$ from $x = a$ to $x = b$.

Writing to Learn Which of the following integrals, if either, calculates the area of the shaded region shown here? Give reasons for your answer.

i. $\int_{-1}^1 (x - (-x)) dx = \int_{-1}^1 2x dx$

ii. $\int_{-1}^1 (-x - (x)) dx = \int_{-1}^1 -2x dx$



Writing to Learn Is the following statement true, sometimes true, or never true? The area of the region between the graphs of the continuous functions $y = f(x)$ and $y = g(x)$ and the vertical lines $x = a$ and $x = b$ ($a < b$) is

$$\int_a^b [f(x) - g(x)] dx.$$

Give reasons for your answer.

Find the area of the propeller-shaped region enclosed between the graphs of

$$y = \frac{2x}{x^2 + 1} \quad \text{and} \quad y = x^3.$$

Find the area of the propeller-shaped region enclosed between the graphs of $y = \sin x$ and $y = x^3$.

Find the positive value of k such that the area of the region enclosed between the graph of $y = k \cos x$ and the graph of $y = kx^2$ is 2.

Standardized Test Questions

You should solve the following problems without using a graphing calculator.

50. **True or False** The area of the region enclosed by the graph of $y = x^2 + 1$ and the line $y = 10$ is 36. Justify your answer.

51. **True or False** The area of the region in the first quadrant enclosed by the graphs of $y = \cos x$, $y = x$, and the y -axis is given by the definite integral $\int_0^{0.739} (x - \cos x) dx$. Justify your answer.

52. **Multiple Choice** Let R be the region in the first quadrant bounded by the x -axis, the graph of $x = y^2 + 2$, and the line $x = 4$. Which of the following integrals gives the area of R ?

(A) $\int_0^{\sqrt{2}} [4 - (y^2 + 2)] dy$ (B) $\int_0^{\sqrt{2}} [(y^2 + 2) - 4] dy$

(C) $\int_{-\sqrt{2}}^{\sqrt{2}} [4 - (y^2 + 2)] dy$ (D) $\int_{-\sqrt{2}}^{\sqrt{2}} [(y^2 + 2) - 4] dy$


(E) $\int_2^4 [4 - (y^2 + 2)] dy$

53. **Multiple Choice** Which of the following gives the area of the region between the graphs of $y = x^2$ and $y = -x$ from $x = 0$ to $x = 3$?

(A) 2 (B) 9/2 (C) 13/2 (D) 13 (E) 27/2

54. **Multiple Choice** Let R be the shaded region enclosed by the graphs of $y = e^{-x^2}$, $y = -\sin(3x)$, and the y -axis as shown in the figure below. Which of the following gives the approximate area of the region R ?

(A) 1.139 (B) 1.445 (C) 1.869 (D) 2.114 (E) 2.340



55. **Multiple Choice** Let f and g be the functions given by $f(x) = e^x$ and $g(x) = 1/x$. Which of the following gives the area of the region enclosed by the graphs of f and g between $x = 1$ and $x = 2$?

(A) $e^2 - e - \ln 2$

(B) $\ln 2 - e^2 + e$

(C) $e^2 - \frac{1}{2}$

(D) $e^2 - e - \frac{1}{2}$

(E) $\frac{1}{e} - \ln 2$

Exploration

56. Group Activity Area of Ellipse


An ellipse with major axis of length $2a$ and minor axis of length $2b$ can be coordinatized with its center at the origin and its major axis horizontal, in which case it is defined by the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

- Find the equations that define the upper and lower semiellipses as functions of x .
- Write an integral expression that gives the area of the ellipse.
- With your group, use NINT to find the areas of ellipses for various lengths of a and b .
- There is a simple formula for the area of an ellipse with major axis of length $2a$ and minor axis of length $2b$. Can you tell what it is from the areas you and your group have found?
- Work with your group to write a *proof* of this area formula by showing that it is the exact value of the integral expression in part (b).

Extending the Ideas

57. Cavalieri's Theorem Bonaventura Cavalieri (1598–1647) discovered that if two plane regions can be arranged to lie over the same interval of the x -axis in such a way that they have identical vertical cross sections at every point (see figure), then the regions have the same area. Show that this theorem is true.



58. Find the area of the region enclosed by the curves

$$y = \frac{x}{x^2 + 1} \quad \text{and} \quad y = mx, \quad 0 < m < 1.$$

7.3 Volumes

What you'll learn about

- Volume As an Integral
- Square Cross Sections
- Circular Cross Sections
- Cylindrical Shells
- Other Cross Sections

and why

The techniques of this section allow us to compute volumes of certain solids in three dimensions.




Figure 7.15 The cross section of an arbitrary solid at point x .

Volume As an Integral

In Section 5.1, Example 3, we estimated the volume of a sphere by partitioning it into thin slices that were nearly cylindrical and summing the cylinders' volumes using MRAM. MRAM sums are Riemann sums, and had we known how at the time, we could have continued on to express the volume of the sphere as a definite integral.

Starting the same way, we can now find the volumes of a great many solids by integration. Suppose we want to find the volume of a solid like the one in Figure 7.15. The cross section of the solid at each point x in the interval $[a, b]$ is a region $R(x)$ of area $A(x)$. If A is a continuous function of x , we can use it to define and calculate the volume of the solid as an integral in the following way.

We partition $[a, b]$ into subintervals of length Δx and slice the solid, as we would a loaf of bread, by planes perpendicular to the x -axis at the partition points. The k th slice, the one between the planes at x_{k-1} and x_k , has approximately the same volume as the cylinder between the two planes based on the region $R(x_k)$ (Figure 7.16).




Figure 7.16 Enlarged view of the slice of the solid between the planes at x_{k-1} and x_k .

The volume of the cylinder is

$$V_k = \text{base area} \times \text{height} = A(x_k) \times \Delta x.$$

The sum

$$\sum V_k = \sum A(x_k) \times \Delta x$$

approximates the volume of the solid.

This is a Riemann sum for $A(x)$ on $[a, b]$. We expect the approximations to improve as the norms of the partitions go to zero, so we define their limiting integral to be the *volume of the solid*.

DEFINITION Volume of a Solid

The **volume of a solid** of known integrable cross section area $A(x)$ from $x = a$ to $x = b$ is the integral of A from a to b ,

$$V = \int_a^b A(x) \, dx.$$

The volume of the paperweight is

$$\begin{aligned} V &= \int_0^\pi A(x) \, dx \\ &= \frac{\pi}{2} \int_0^\pi (\sin x)^2 \, dx \\ &\approx \frac{\pi}{2} \text{NINT}((\sin x)^2, x, 0, \pi) \\ &\approx \frac{\pi}{2}(1.570796327). \end{aligned}$$

The number in parentheses looks like half of π , an observation that can be confirmed analytically, and which we support numerically by dividing by π to get 0.5. The volume of the paperweight is


$$\frac{\pi}{2} \cdot \frac{\pi}{2} = \frac{\pi^2}{4} \approx 2.47 \text{ in}^3.$$

Now try Exercise 39(a)

Bonaventura Cavalieri
(1598–1647)



Cavalieri, a student of Galileo, discovered that if two plane regions can be arranged to lie over the same interval of the x -axis in such a way that they have identical vertical cross sections at every point, then the regions have the same area. This theorem and a letter of recommendation from Galileo were enough to win Cavalieri a chair at the University of Bologna in 1629. The solid geometry version in Example 7, which Cavalieri never proved, was named after him by later geometers.



EXAMPLE 7 Cavalieri's Volume Theorem

Cavalieri's volume theorem says that solids with equal altitudes and identical cross section areas at each height have the same volume (Figure 7.31). This follows immediately from the definition of volume, because the cross section area function $A(x)$ and the interval $[a, b]$ are the same for both solids.





Figure 7.31 Cavalieri's volume theorem: These solids have the same volume. You can illustrate this yourself with stacks of coins. (Example 7)

Now try Exercise 41

EXPLORATION 2 Surface Area

We know how to find the volume of a solid of revolution, but how would we find the *surface area*? As before, we partition the solid into thin slices, but now we wish to form a Riemann sum of approximations to *surface areas of slices* (rather than of volumes of slices).



A typical slice has a surface area that can be approximated by $2\pi \cdot f(x) \cdot \Delta s$, where Δs is the tiny *slant height* of the slice. We will see in Section 7.4, when we study arc length, that $\Delta s = \sqrt{\Delta x^2 + \Delta y^2}$, and that this can be written as $\Delta s = \sqrt{1 + (f'(x_k))^2} \Delta x$.

Thus, the surface area is approximated by the Riemann sum

$$\sum_{k=1}^n 2\pi f(x_k) \sqrt{1 + (f'(x_k))^2} \Delta x.$$

1. Write the limit of the Riemann sums as a definite integral from a to b . When will the limit exist?
2. Use the formula from part 1 to find the surface area of the solid generated by revolving a single arch of the curve $y = \sin x$ about the x -axis.
3. The region enclosed by the graphs of $y^2 = x$ and $x = 4$ is revolved about the x -axis to form a solid. Find the surface area of the solid.


Quick Review 7.3 (For help, go to Section 1.2.)

- Exercises 1–10, give a formula for the area of the plane region in terms of the single variable x .
1. a square with sides of length x
 2. a square with diagonals of length x
 3. a semicircle of radius x
 4. a semicircle of diameter x
 5. an equilateral triangle with sides of length x
 6. an isosceles right triangle with legs of length x
 7. an isosceles right triangle with hypotenuse x
 8. an isosceles triangle with two sides of length $2x$ and one side of length x
 9. a triangle with sides $3x$, $4x$, and $5x$
 10. a regular hexagon with sides of length x


Section 7.3 Exercises

In Exercises 1 and 2, find a formula for the area $A(x)$ of the cross sections of the solid that are perpendicular to the x -axis.


1. The solid lies between planes perpendicular to the x -axis at $x = -1$ and $x = 1$. The cross sections perpendicular to the x -axis between these planes run from the semicircle $y = -\sqrt{1-x^2}$ to the semicircle $y = \sqrt{1-x^2}$.
- (a) The cross sections are circular disks with diameters in the xy -plane.




- (b) The cross sections are squares with bases in the xy -plane.




- (c) The cross sections are squares with diagonals in the xy -plane. (The length of a square's diagonal is $\sqrt{2}$ times the length of its sides.)




- (d) The cross sections are equilateral triangles with bases in the xy -plane.



2. The solid lies between planes perpendicular to the x -axis at $x = 0$ and $x = 4$. The cross sections perpendicular to the x -axis between these planes run from $y = -\sqrt{x}$ to $y = \sqrt{x}$.
- (a) The cross sections are circular disks with diameters in the xy -plane.




- (b) The cross sections are squares with bases in the xy -plane.



- (c) The cross sections are squares with diagonals in the xy -plane.
- (d) The cross sections are equilateral triangles with bases in the xy -plane.

In Exercises 3–6, find the volume of the solid analytically.

3. The solid lies between planes perpendicular to the x -axis at $x = 0$ and $x = 4$. The cross sections perpendicular to the x -axis on the interval $0 \leq x \leq 4$ are squares whose diagonals run from $y = -\sqrt{x}$ to $y = \sqrt{x}$.
4. The solid lies between planes perpendicular to the x -axis at $x = -1$ and $x = 1$. The cross sections perpendicular to the x -axis between these planes are circular disks whose diameters run from the parabola $y = x^2$ to the parabola $y = 2 - x^2$.




5. The solid lies between planes perpendicular to the x -axis at $x = -1$ and $x = 1$. The cross sections perpendicular to the x -axis between these planes are squares whose bases run from the semicircle $y = -\sqrt{1-x^2}$ to the semicircle $y = \sqrt{1-x^2}$.


The solid lies between planes perpendicular to the x -axis at $x = -1$ and $x = 1$. The cross sections perpendicular to the x -axis between these planes are squares whose diagonals run from the semicircle $y = -\sqrt{1-x^2}$ to the semicircle $y = \sqrt{1-x^2}$.

Exercises 7–10, find the volume of the solid generated by revolving the shaded region about the given axis.


7. about the x -axis




8. about the y -axis



9. about the y -axis



10. about the x -axis



Exercises 11–20, find the volume of the solid generated by revolving the region bounded by the lines and curves about the indicated axis.

11. $y = x^2$, $y = 0$, $x = 2$ 12. $y = x^3$, $y = 0$, $x = 2$
 13. $y = \sqrt{9-x^2}$, $y = 0$ 14. $y = x - x^2$, $y = 0$
 15. $y = x$, $y = 1$, $x = 0$ 16. $y = 2x$, $y = x$, $x = 1$
 17. $y = x^2 + 1$, $y = x + 3$ 18. $y = 4 - x^2$, $y = 2 - x$
 19. $y = \sec x$, $y = \sqrt{2}$, $-\pi/4 \leq x \leq \pi/4$
 20. $y = -\sqrt{x}$, $y = -2$, $x = 0$

Exercises 21 and 22, find the volume of the solid generated by revolving the region about the given line.

21. the region in the first quadrant bounded above by the line $y = \sqrt{2}$, below by the curve $y = \sec x \tan x$, and on the left by the y -axis, about the line $y = \sqrt{2}$

22. the region in the first quadrant bounded above by the line $y = 2$, below by the curve $y = 2 \sin x$, $0 \leq x \leq \pi/2$, and on the left by the y -axis, about the line $y = 2$

Exercises 23–28, find the volume of the solid generated by revolving the region about the y -axis.

23. the region enclosed by $x = \sqrt{5}y^2$, $x = 0$, $y = -1$, $y = 1$

24. the region enclosed by $x = y^{3/2}$, $x = 0$, $y = 2$

25. the region enclosed by the triangle with vertices $(1, 0)$, $(2, 1)$, and $(1, 1)$

26. the region enclosed by the triangle with vertices $(0, 1)$, $(1, 0)$, and $(1, 1)$


27. the region in the first quadrant bounded above by the parabola $y = x^2$, below by the x -axis, and on the right by the line $x = 2$
28. the region bounded above by the curve $y = \sqrt{x}$ and below by the line $y = x$

Group Activity In Exercises 29–32, find the volume of the solid described.


29. Find the volume of the solid generated by revolving the region bounded by $y = \sqrt{x}$ and the lines $y = 2$ and $x = 0$ about
 (a) the x -axis. (b) the y -axis.
 (c) the line $y = 2$. (d) the line $x = 4$.
30. Find the volume of the solid generated by revolving the triangular region bounded by the lines $y = 2x$, $y = 0$, and $x = 1$ about
 (a) the line $x = 1$. (b) the line $x = 2$.
31. Find the volume of the solid generated by revolving the region bounded by the parabola $y = x^2$ and the line $y = 1$ about
 (a) the line $y = 1$. (b) the line $y = 2$.
 (c) the line $y = -1$.
32. By integration, find the volume of the solid generated by revolving the triangular region with vertices $(0, 0)$, $(b, 0)$, $(0, h)$ about
 (a) the x -axis. (b) the y -axis.

In Exercises 33 and 34, use the cylindrical shell method to find the volume of the solid generated by revolving the shaded region about the indicated axis.

33. (a) the x -axis (b) the line $y = 1$
 (c) the line $y = 8/5$ (d) the line $y = -2/5$



34. (a) the x -axis (b) the line $y = 2$
 (c) the line $y = 5$ (d) the line $y = -5/8$




In Exercises 35–38, use the cylindrical shell method to find the volume of the solid generated by revolving the region bounded by the curves about the y -axis.


- 35. $y = x, y = -x/2, x = 2$
- 36. $y = x^2, y = 2 - x, x = 0, \text{ for } x \geq 0$
- 37. $y = \sqrt{x}, y = 0, x = 4$
- 38. $y = 2x - 1, y = \sqrt{x}, x = 0$

In Exercises 39–42, find the volume of the solid analytically.


- 39. The base of a solid is the region between the curve $y = 2\sqrt{\sin x}$ and the interval $[0, \pi]$ on the x -axis. The cross sections perpendicular to the x -axis are
 - (a) equilateral triangles with bases running from the x -axis to the curve as shown in the figure.



- (b) squares with bases running from the x -axis to the curve.
- 40. The solid lies between planes perpendicular to the x -axis at $x = -\pi/3$ and $x = \pi/3$. The cross sections perpendicular to the x -axis are
 - (a) circular disks with diameters running from the curve $y = \tan x$ to the curve $y = \sec x$.
 - (b) squares whose bases run from the curve $y = \tan x$ to the curve $y = \sec x$.
- 41. The solid lies between planes perpendicular to the y -axis at $y = 0$ and $y = 2$. The cross sections perpendicular to the y -axis are circular disks with diameters running from the y -axis to the parabola $x = \sqrt{5}y^2$.
- 42. The base of the solid is the disk $x^2 + y^2 \leq 1$. The cross sections by planes perpendicular to the y -axis between $y = -1$ and $y = 1$ are isosceles right triangles with one leg in the disk.



- 43. **Writing to Learn** A solid lies between planes perpendicular to the x -axis at $x = 0$ and $x = 12$. The cross sections by planes perpendicular to the x -axis are circular disks whose diameters run from the line $y = x/2$ to the line $y = x$ as shown in the figure. Explain why the solid has the same volume as a right circular cone with base radius 3 and height 12.




- 44. **A Twisted Solid** A square of side length s lies in a plane perpendicular to a line L . One vertex of the square lies on L . As this square moves a distance h along L , the square turns one revolution about L to generate a corkscrew-like column with square cross sections.


- (a) Find the volume of the column.
- (b) **Writing to Learn** What will the volume be if the square turns twice instead of once? Give reasons for your answer.

- 45. Find the volume of the solid generated by revolving the region in the first quadrant bounded by $y = x^3$ and $y = 4x$ about
 - (a) the x -axis,
 - (b) the line $y = 8$.
- 46. Find the volume of the solid generated by revolving the region bounded by $y = 2x - x^2$ and $y = x$ about
 - (a) the y -axis,
 - (b) the line $x = 1$.
- 47. The region in the first quadrant that is bounded above by the curve $y = 1/\sqrt{x}$, on the left by the line $x = 1/4$, and below by the line $y = 1$ is revolved about the y -axis to generate a solid. Find the volume of the solid by (a) the washer method and (b) the cylindrical shell method.

- 48. Let $f(x) = \begin{cases} (\sin x)/x, & 0 < x \leq \pi \\ 1, & x = 0. \end{cases}$
 - (a) Show that $xf(x) = \sin x, 0 \leq x \leq \pi$.
 - (b) Find the volume of the solid generated by revolving the shaded region about the y -axis.



Designing a Plumb Bob Having been asked to design a brass plumb bob that will weigh in the neighborhood of 190 g, you decide to shape it like the solid of revolution shown here.



- (a) Find the plumb bob's volume.
- (b) If you specify a brass that weighs 8.5 g/cm^3 , how much will the plumb bob weigh to the nearest gram?

Volume of a Bowl A bowl has a shape that can be generated by revolving the graph of $y = x^2/2$ between $y = 0$ and $y = 5$ about the y -axis.

- (a) Find the volume of the bowl.
- (b) If we fill the bowl with water at a constant rate of 3 cubic units per second, how fast will the water level in the bowl be rising when the water is 4 units deep?

The Classical Bead Problem A round hole is drilled through the center of a spherical solid of radius r . The resulting cylindrical hole has height 4 cm.

- (a) What is the volume of the solid that remains?
- (b) What is unusual about the answer?

Writing to Learn Explain how you could estimate the volume of a solid of revolution by measuring the shadow cast on a table parallel to its axis of revolution by a light shining directly above it.

Same Volume about Each Axis The region in the first quadrant enclosed between the graph of $y = ax - x^2$ and the x -axis generates the same volume whether it is revolved about the x -axis or the y -axis. Find the value of a .

(Continuation of Exploration 2) Let $x = g(y) > 0$ have a continuous first derivative on $[c, d]$. Show that the area of the surface generated by revolving the curve $x = g(y)$ about the y -axis is

$$S = \int_c^d 2\pi g(y) \sqrt{1 + (g'(y))^2} dy.$$

Exercises 55–62, find the area of the surface generated by revolving the curve about the indicated axis.

- 55. $x = \sqrt{y}, 0 \leq y \leq 2; y$ -axis
- 56. $x = y^3/3, 0 \leq y \leq 1; y$ -axis
- 57. $x = y^{1/2} - (1/3)^{3/2}, 1 \leq y \leq 3; y$ -axis
- 58. $x = \sqrt{2y - 1}, (5/8) \leq y \leq 1; y$ -axis

- 59. $y = x^2, 0 \leq x \leq 2; x$ -axis
- 60. $y = 3x - x^2, 0 \leq x \leq 3; x$ -axis
- 61. $y = \sqrt{2x - x^2}, 0.5 \leq x \leq 1.5; x$ -axis
- 62. $y = \sqrt{x + 1}, -1 \leq x \leq 5; x$ -axis

Standardized Test Questions


You may use a graphing calculator to solve the following problems.

- 63. **True or False** The volume of a solid of a known integrable cross section area $A(x)$ from $x = a$ to $x = b$ is $\int_a^b A(x) dx$. Justify your answer.
- 64. **True or False** If the region enclosed by the y -axis, the line $y = 2$, and the curve $y = \sqrt{x}$ is revolved about the y -axis, the volume of the solid is given by the definite integral $\int_0^2 \pi y^2 dy$. Justify your answer.
- 65. **Multiple Choice** The base of a solid S is the region enclosed by the graph of $y = \ln x$, the line $x = e$, and the x -axis. If the cross sections of S perpendicular to the x -axis are squares, which of the following gives the best approximation of the volume of S ?
 - (A) 0.718 (B) 1.718 (C) 2.718 (D) 3.171 (E) 7.388
- 66. **Multiple Choice** Let R be the region in the first quadrant bounded by the graph of $y = 8 - x^{3/2}$, the x -axis, and the y -axis. Which of the following gives the best approximation of the volume of the solid generated when R is revolved about the x -axis?
 - (A) 60.3 (B) 115.2 (C) 225.4 (D) 319.7 (E) 361.9
- 67. **Multiple Choice** Let R be the region enclosed by the graph of $y = x^2$, the line $x = 4$, and the x -axis. Which of the following gives the best approximation of the volume of the solid generated when R is revolved about the y -axis?
 - (A) 64π (B) 128π (C) 256π (D) 360 (E) 512
- 68. **Multiple Choice** Let R be the region enclosed by the graphs of $y = e^{-x}, y = e^x$, and $x = 1$. Which of the following gives the volume of the solid generated when R is revolved about the x -axis?
 - (A) $\int_0^1 (e^x - e^{-x}) dx$
 - (B) $\int_0^1 (e^{2x} - e^{-2x}) dx$
 - (C) $\int_0^1 (e^x - e^{-x})^2 dx$
 - (D) $\pi \int_0^1 (e^{2x} - e^{-2x}) dx$
 - (E) $\pi \int_0^1 (e^x - e^{-x})^2 dx$

Explorations

69. **Max-Min** The arch $y = \sin x$, $0 \leq x \leq \pi$, is revolved about the line $y = c$, $0 \leq c \leq 1$, to generate the solid in the figure.

- (a) Find the value of c that minimizes the volume of the solid. What is the minimum volume?
- (b) What value of c in $[0, 1]$ maximizes the volume of the solid?
- (c) **Writing to Learn** Graph the solid's volume as a function of c , first for $0 \leq c \leq 1$ and then on a larger domain. What happens to the volume of the solid as c moves away from $[0, 1]$? Does this make sense physically? Give reasons for your answers.



70. **A Vase** We wish to estimate the volume of a flower vase using only a calculator, a string, and a ruler. We measure the height of the vase to be 6 inches. We then use the string and the ruler to find circumferences of the vase (in inches) at half-inch intervals. (We list them from the top down to correspond with the picture of the vase.)




Circumferences	
5.4	10.8
4.5	11.6
4.4	11.6
5.1	10.8
6.3	9.0
7.8	6.3
9.4	

- (a) Find the areas of the cross sections that correspond to the given circumferences.
- (b) Express the volume of the vase as an integral with respect to y over the interval $[0, 6]$.
- (c) Approximate the integral using the Trapezoidal Rule with $n = 12$.

Extending the Ideas

71. **Volume of a Hemisphere** Derive the formula $V = (2/3)\pi R^3$ for the volume of a hemisphere of radius R by comparing its cross sections with the cross sections of a solid right circular cylinder of radius R and height R from which a solid right circular cone of base radius R and height R has been removed as suggested by the figure.



72. **Volume of a Torus** The disk $x^2 + y^2 \leq a^2$ is revolved about the line $x = b$ ($b > a$) to generate a solid shaped like a doughnut called a *torus*. Find its volume. (Hint: $\int_{-a}^a \sqrt{a^2 - y^2} dy = \pi a^2$ since it is the area of a semicircle of radius a .)

73. **Filling a Bowl**

- (a) **Volume** A hemispherical bowl of radius a contains water to a depth h . Find the volume of water in the bowl.
- (b) **Related Rates** Water runs into a sunken concrete hemispherical bowl of radius 5 m at a rate of $0.2 \text{ m}^3/\text{sec}$. How fast is the water level in the bowl rising when the water is 4 m deep?

74. **Consistency of Volume Definitions** The volume formulas in calculus are consistent with the standard formulas from geometry in the sense that they agree on objects to which both apply.

- (a) As a case in point, show that if you revolve the region enclosed by the semicircle $y = \sqrt{a^2 - x^2}$ and the x -axis about the x -axis to generate a solid sphere, the calculus formula for volume at the beginning of the section will give $(4/3)\pi a^3$ for the volume just as it should.
- (b) Use calculus to find the volume of a right circular cone of height h and base radius r .

Quick Quiz for AP* Preparation: Sections 7.1–7.3

You may use a graphing calculator to solve the following problems.

Multiple Choice The base of a solid is the region in the first quadrant bounded by the x -axis, the graph of $y = \sin^{-1} x$, and the vertical line $x = 1$. For this solid, each cross section perpendicular to the x -axis is a square. What is the volume?

- (A) 0.117 (B) 0.285 (C) 0.467 (D) 0.571 (E) 1.571

Multiple Choice Let R be the region in the first quadrant bounded by the graph of $y = 3x - x^2$ and the x -axis. A solid is generated when R is revolved about the vertical line $x = -1$. Set up, but do not evaluate, the definite integral that gives the volume of this solid.

- (A) $\int_0^3 2\pi(x+1)(3x-x^2) dx$
- (B) $\int_{-1}^3 2\pi(x+1)(3x-x^2) dx$
- (C) $\int_0^3 2\pi(x)(3x-x^2) dx$
- (D) $\int_0^3 2\pi(3x-x^2)^2 dx$
- (E) $\int_0^3 (3x-x^2) dx$

3. **Multiple Choice** A developing country consumes oil at a rate given by $r(t) = 20e^{0.2t}$ million barrels per year, where t is time measured in years, for $0 \leq t \leq 10$. Which of the following expressions gives the amount of oil consumed by the country during the time interval $0 \leq t \leq 10$?

- (A) $r(10)$
- (B) $r(10) - r(0)$
- (C) $\int_0^{10} r'(t) dt$
- (D) $\int_0^{10} r(t) dt$
- (E) $10 \cdot r(10)$

4. **Free Response** Let R be the region bounded by the graphs of $y = \sqrt{x}$, $y = e^{-x}$, and the y -axis.

- (a) Find the area of R .
- (b) Find the volume of the solid generated when R is revolved about the horizontal line $y = -1$.
- (c) The region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is a semicircle whose diameter runs from the graph of $y = \sqrt{x}$ to the graph of $y = e^{-x}$. Find the volume of this solid.

EXAMPLE 2 Applying the Definition

Find the exact length of the curve

$$y = \frac{4\sqrt{2}}{3}x^{3/2} - 1 \quad \text{for } 0 \leq x \leq 1.$$

SOLUTION

$$\frac{dy}{dx} = \frac{4\sqrt{2}}{3} \cdot \frac{3}{2}x^{1/2} = 2\sqrt{2}x^{1/2},$$

which is continuous on $[0, 1]$. Therefore,

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\ &= \int_0^1 \sqrt{1 + (2\sqrt{2}x^{1/2})^2} dx \\ &= \int_0^1 \sqrt{1 + 8x} dx \\ &= \frac{2}{3} \cdot \frac{1}{8} (1 + 8x)^{3/2} \Big|_0^1 \\ &= \frac{13}{6}. \end{aligned}$$

Now try Exercise 11

We asked for an exact length in Example 2 to take advantage of the rare opportunity afforded of taking the antiderivative of an arc length integrand. When you add 1 to the square of the derivative of an arbitrary smooth function and then take the square root of the sum, the result is rarely antiderivable by reasonable methods. We know a few nice functions that give "nice" integrands, but we are saving those for the exercises.

Vertical Tangents, Corners, and Cusps

Sometimes a curve has a vertical tangent, corner, or cusp where the derivative we need to work with is undefined. We can sometimes get around such a difficulty in ways illustrated in the following examples.

EXAMPLE 3 A Vertical Tangent

Find the length of the curve $y = x^{1/3}$ between $(-8, -2)$ and $(8, 2)$.

SOLUTION

The derivative

$$\frac{dy}{dx} = \frac{1}{3}x^{-2/3} = \frac{1}{3x^{2/3}}$$

is not defined at $x = 0$. Graphically, there is a vertical tangent at $x = 0$ where the derivative becomes infinite (Figure 7.36). If we change to x as a function of y , the tangent at the origin will be horizontal (Figure 7.37) and the derivative will be zero instead of undefined. Solving $y = x^{1/3}$ for x gives $x = y^3$, and we have

$$L = \int_{-2}^2 \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy = \int_{-2}^2 \sqrt{1 + (3y^2)^2} dy \approx 17.26. \quad \text{Using NINT}$$

Now try Exercise 12




Figure 7.36 The graph of $y = x^{1/3}$ has a vertical tangent line at the origin where dy/dx does not exist. (Example 3)




Figure 7.37 The curve in Figure 7.36 plotted with x as a function of y . The tangent at the origin is now horizontal. (Example 3)

What happens if you fail to notice that dy/dx is undefined at $x = 0$ and ask your calculator to compute

$$\text{NINT} \left(\sqrt{1 + \left((1/3)x^{-2/3} \right)^2}, x, -8, 8 \right)?$$

This actually depends on your calculator. If, in the process of its calculations, it tries to evaluate the function at $x = 0$, then some sort of domain error will result. If it tries to find convergent Riemann sums near $x = 0$, it might get into a long, futile loop of computations that you will have to interrupt. Or it might actually produce an answer—in which case you hope it would be sufficiently bizarre for you to realize that it should not be trusted.

EXAMPLE 4 Getting Around a Corner

Find the length of the curve $y = x^2 - 4|x| - x$ from $x = -4$ to $x = 4$.

SOLUTION

We should always be alert for abrupt slope changes when absolute value is involved. We graph the function to check (Figure 7.38).

There is clearly a corner at $x = 0$ where neither dy/dx nor dx/dy can exist. To find the length, we split the curve at $x = 0$ to write the function without absolute values:


$$x^2 - 4|x| - x = \begin{cases} x^2 + 3x & \text{if } x < 0, \\ x^2 - 5x & \text{if } x \geq 0. \end{cases}$$

Then,

$$\begin{aligned} L &= \int_{-4}^0 \sqrt{1 + (2x + 3)^2} dx + \int_0^4 \sqrt{1 + (2x - 5)^2} dx \\ &\approx 19.56. \quad \text{By NINT} \end{aligned}$$

Now try Exercise 27.

Finally, cusps are handled the same way corners are: split the curve into smooth pieces and add the lengths of those pieces.



[-5, 5] by [-7, 5]

Figure 7.38 The graph of

$$y = x^2 - 4|x| - x, \quad -4 \leq x \leq 4,$$

has a corner at $x = 0$ where neither dy/dx nor dx/dy exists. We find the lengths of the two smooth pieces and add them together. (Example 4)

Quick Review 7.4 (For help, go to Sections 1.3 and 3.2.)

In Exercises 1–5, simplify the function.

1. $\sqrt{1 + 2x + x^2}$ on $[1, 5]$

2. $\sqrt{1 - x + \frac{x^2}{4}}$ on $[-3, -1]$

3. $\sqrt{1 + (\tan x)^2}$ on $[0, \pi/3]$

4. $\sqrt{1 + (x/4 - 1/x)^2}$ on $[4, 12]$

5. $\sqrt{1 + \cos 2x}$ on $[0, \pi/2]$

In Exercises 6–10, identify all values of x for which the function fails to be differentiable.

6. $f(x) = |x - 4|$

7. $f(x) = 5x^{2/3}$

8. $f(x) = \sqrt[5]{x + 3}$

9. $f(x) = \sqrt{x^2 - 4x + 4}$

10. $f(x) = 1 + \sqrt[3]{\sin x}$

Section 7.4 Exercises

In Exercises 1–10,

- (a) set up an integral for the length of the curve;
- (b) graph the curve to see what it looks like;
- (c) use NINT to find the length of the curve.

1. $y = x^2, -1 \leq x \leq 2$
2. $y = \tan x, -\pi/3 \leq x \leq 0$
3. $x = \sin y, 0 \leq y \leq \pi$
4. $x = \sqrt{1 - y^2}, -1/2 \leq y \leq 1/2$
5. $y^2 + 2y = 2x + 1$, from $(-1, -1)$ to $(7, 3)$
6. $y = \sin x - x \cos x, 0 \leq x \leq \pi$
7. $y = \int_0^x \tan t \, dt, 0 \leq x \leq \pi/6$
8. $x = \int_0^y \sqrt{\sec^2 t - 1} \, dt, -\pi/3 \leq y \leq \pi/4$
9. $y = \sec x, -\pi/3 \leq x \leq \pi/3$
10. $y = (e^x + e^{-x})/2, -3 \leq x \leq 3$

In Exercises 11–18, find the exact length of the curve analytically by antidifferentiation. You will need to simplify the integrand algebraically before finding an antiderivative.

11. $y = (1/3)(x^2 + 2)^{3/2}$ from $x = 0$ to $x = 3$
12. $y = x^{3/2}$ from $x = 0$ to $x = 4$
13. $x = (y^3/3) + 1/(4y)$ from $y = 1$ to $y = 3$
[Hint: $1 + (dx/dy)^2$ is a perfect square.]
14. $x = (y^4/4) + 1/(8y^2)$ from $y = 1$ to $y = 2$
[Hint: $1 + (dx/dy)^2$ is a perfect square.]
15. $x = (y^3/6) + 1/(2y)$ from $y = 1$ to $y = 2$
[Hint: $1 + (dx/dy)^2$ is a perfect square.]
16. $y = (x^3/3) + x^2 + x + 1/(4x + 4), 0 \leq x \leq 2$
17. $x = \int_0^y \sqrt{\sec^4 t - 1} \, dt, -\pi/4 \leq y \leq \pi/4$
18. $y = \int_{-2}^x \sqrt{3t^4 - 1} \, dt, -2 \leq x \leq -1$
19. (a) **Group Activity** Find a curve through the point $(1, 1)$ whose length integral is

$$L = \int_1^4 \sqrt{1 + \frac{1}{4x}} \, dx.$$

(b) **Writing to Learn** How many such curves are there? Give reasons for your answer.

20. (a) **Group Activity** Find a curve through the point $(0, 1)$ whose length integral is

$$L = \int_1^2 \sqrt{1 + \frac{1}{y^4}} \, dy.$$


(b) **Writing to Learn** How many such curves are there? Give reasons for your answer.

21. Find the length of the curve

$$y = \int_0^x \sqrt{\cos 2t} \, dt$$

from $x = 0$ to $x = \pi/4$.


22. **The Length of an Astroid** The graph of the equation $x^{2/3} + y^{2/3} = 1$ is one of the family of curves called *astroids* (not “asteroids”) because of their starlike appearance (see figure). Find the length of this particular astroid by finding the length of half the first quadrant portion, $y = (1 - x^{2/3})^{3/2}, \sqrt{2}/4 \leq x \leq 1$ and multiplying by 8.




23. **Fabricating Metal Sheets** Your metal fabrication company is bidding for a contract to make sheets of corrugated steel roofing like the one shown here. The cross sections of the corrugated sheets are to conform to the curve

$$y = \sin\left(\frac{3\pi}{20}x\right), \quad 0 \leq x \leq 20 \text{ in.}$$

If the roofing is to be stamped from flat sheets by a process that does not stretch the material, how wide should the original material be? Give your answer to two decimal places.



24. **Tunnel Construction** Your engineering firm is bidding for the contract to construct the tunnel shown on the next page. The tunnel is 300 ft long and 50 ft wide at the base. The cross section is shaped like one arch of the curve $y = 25 \cos(\pi x/50)$. Upon completion, the tunnel’s inside surface (excluding the roadway) will be treated with a waterproof sealer that costs \$1.75 per square foot to apply. How much will it cost to apply the sealer?



Exercises 25 and 26, find the length of the curve.

25. $f(x) = x^{1/3} + x^{2/3}, 0 \leq x \leq 2$

26. $f(x) = \frac{x-1}{4x^2+1}, -\frac{1}{2} \leq x \leq 1$

Exercises 27–29, find the length of the nonsmooth curve.

27. $y = x^3 + 5|x|$ from $x = -2$ to $x = 1$

28. $\sqrt{x} + \sqrt{y} = 1$

29. $y = \sqrt[3]{x}$ from $x = 0$ to $x = 16$

Writing to Learn Explain geometrically why it does not work to use short horizontal line segments to approximate the lengths of small arcs when we search for a Riemann sum that leads to the formula for arc length.

Writing to Learn A curve is totally contained inside the square with vertices $(0, 0), (1, 0), (1, 1),$ and $(0, 1)$. Is there any limit to the possible length of the curve? Explain.

Standardized Test Questions

You should solve the following problems without using a graphing calculator.

True or False If a function $y = f(x)$ is continuous on an interval $[a, b]$, then the length of its curve is given by

$$\int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx.$$
 Justify your answer.

True or False If a function $y = f(x)$ is differentiable on an interval $[a, b]$, then the length of its curve is given by

$$\int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx.$$
 Justify your answer.

34. **Multiple Choice** Which of the following gives the best approximation of the length of the arc of $y = \cos(2x)$ from $x = 0$ to $x = \pi/4$?
(A) 0.785 (B) 0.955 (C) 1.0 (D) 1.318 (E) 1.977

35. **Multiple Choice** Which of the following expressions gives the length of the graph of $x = y^3$ from $y = -2$ to $y = 2$?

- (A) $\int_{-2}^2 (1 + y^6) \, dy$
- (B) $\int_{-2}^2 \sqrt{1 + y^6} \, dy$
- (C) $\int_{-2}^2 \sqrt{1 + 9y^4} \, dy$
- (D) $\int_{-2}^2 \sqrt{1 + x^2} \, dx$
- (E) $\int_{-2}^2 \sqrt{1 + x^4} \, dx$

36. **Multiple Choice** Find the length of the curve described by $y = \frac{2}{3}x^{3/2}$ from $x = 0$ to $x = 8$.


- (A) $\frac{26}{3}$
- (B) $\frac{52}{3}$
- (C) $\frac{512\sqrt{2}}{15}$
- (D) $\frac{512\sqrt{2}}{15} + 8$
- (E) 96

37. **Multiple Choice** Which of the following expressions should be used to find the length of the curve $y = x^{2/3}$ from $x = -1$ to $x = 1$?

- (A) $2 \int_0^1 \sqrt{1 + \frac{9}{4}y} \, dy$
- (B) $\int_{-1}^1 \sqrt{1 + \frac{9}{4}y} \, dy$
- (C) $\int_0^1 \sqrt{1 + y^3} \, dy$
- (D) $\int_0^1 \sqrt{1 + y^6} \, dy$
- (E) $\int_0^1 \sqrt{1 + y^{9/4}} \, dy$


Exploration

38. **Modeling Running Tracks** Two lanes of a running track are modeled by the semiellipses as shown. The equation for lane 1 is $y = \sqrt{100 - 0.2x^2}$, and the equation for lane 2 is $y = \sqrt{150 - 0.2x^2}$. The starting point for lane 1 is at the negative x -intercept $(-\sqrt{500}, 0)$. The finish points for both lanes are the positive x -intercepts. Where should the starting point be placed on lane 2 so that the two lane lengths will be equal (running clockwise)?



Extending the Ideas

39. Using Tangent Fins to Find Arc Length Assume f is smooth on $[a, b]$ and partition the interval $[a, b]$ in the usual way. In each subinterval $[x_{k-1}, x_k]$ construct the *tangent fin* at the point $(x_{k-1}, f(x_{k-1}))$ as shown in the figure.



(a) Show that the length of the k th tangent fin over the interval $[x_{k-1}, x_k]$ equals

$$\sqrt{(\Delta x_k)^2 + (f'(x_{k-1})\Delta x_k)^2}.$$

(b) Show that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n (\text{length of } k\text{th tangent fin}) = \int_a^b \sqrt{1 + (f'(x))^2} dx,$$

which is the length L of the curve $y = f(x)$ from $x = a$ to $x = b$.

40. Is there a smooth curve $y = f(x)$ whose length over the interval $0 \leq x \leq a$ is always $a\sqrt{2}$? Give reasons for your answer.

7.5

Applications from Science and Statistics

What you'll learn about

- Work Revisited
- Fluid Force and Fluid Pressure
- Normal Probabilities
- ... and why

It is important to see applications of integrals as various accumulation functions.

Our goal in this section is to hint at the diversity of ways in which the definite integral can be used. The contexts may be new to you, but we will explain what you need to know as we go along.

Work Revisited

Recall from Section 7.1 that *work* is defined as force (in the direction of motion) times displacement. A familiar example is to move against the force of gravity to lift an object. The object has to move, incidentally, before “work” is done, no matter how tired you get *trying*.

If the force $F(x)$ is not constant, then the work done in moving an object from $x = a$ to $x = b$ is the definite integral $W = \int_a^b F(x)dx$.

1 newton \approx 1 lb

1 newton(1 meter) = 1 N \cdot m = 1 Joule

EXAMPLE 1 Finding the Work Done by a Force

Find the work done by the force $F(x) = \cos(\pi x)$ newtons along the x -axis from $x = 0$ meters to $x = 1/2$ meter.

SOLUTION

$$\begin{aligned} W &= \int_0^{1/2} \cos(\pi x) dx \\ &= \frac{1}{\pi} \sin(\pi x) \Big|_0^{1/2} \\ &= \frac{1}{\pi} \left(\sin\left(\frac{\pi}{2}\right) - \sin(0) \right) \\ &= \frac{1}{\pi} \approx 0.318 \end{aligned}$$

Now try Exercise 1.

EXAMPLE 2 Work Done Lifting

A leaky bucket weighs 22 newtons (N) empty. It is lifted from the ground at a constant rate to a point 20 m above the ground by a rope weighing 0.4 N/m. The bucket starts with 70 N (approximately 7.1 liters) of water, but it leaks at a constant rate and just finishes draining as the bucket reaches the top. Find the amount of work done

- (a) lifting the bucket alone;
- (b) lifting the water alone;
- (c) lifting the rope alone;
- (d) lifting the bucket, water, and rope together.

SOLUTION

(a) *The bucket alone.* This is easy because the bucket’s weight is constant. To lift it, you must exert a force of 22 N through the entire 20-meter interval.

$$\text{Work} = (22 \text{ N}) \times (20 \text{ m}) = 440 \text{ N} \cdot \text{m} = 440 \text{ J}$$

Figure 7.39 shows the graph of force vs. distance applied. The work corresponds to the area under the force graph.




Figure 7.39 The work done by a constant 22-N force lifting a bucket 20 m is 440 N \cdot m. (Example 2)

continued

EXAMPLE 6 A Telephone Help Line

Suppose a telephone help line takes a mean of 2 minutes to answer calls. If the standard deviation is $\sigma = 0.5$, then 68% of the calls are answered in the range of 1.5 to 2.5 minutes and 99.7% of the calls are answered in the range of 0.5 to 3.5 minutes.

Now try Exercise 8

EXAMPLE 7 Weights of Spinach Boxes

Suppose that frozen spinach boxes marked as “10 ounces” of spinach have a mean weight of 10.3 ounces and a standard deviation of 0.2 ounce.

- (a) What percentage of *all* such spinach boxes can be expected to weigh between 10 and 11 ounces?
- (b) What percentage would we expect to weigh less than 10 ounces?
- (c) What is the probability that a box weighs *exactly* 10 ounces?

SOLUTION

Assuming that some person or machine is *trying* to pack 10 ounces of spinach into these boxes, we expect that most of the weights will be around 10, with probabilities tapering off for boxes being heavier or lighter. We expect, in other words, that a normal pdf will model these probabilities. First, we define $f(x)$ using the formula:

$$f(x) = \frac{1}{0.2\sqrt{2\pi}} e^{-(x-10.3)^2/(0.08)}$$

The graph (Figure 7.50) has the look we are expecting.

- (a) For an arbitrary box of this spinach, the probability that it weighs between 10 and 11 ounces is the area under the curve from 10 to 11, which is

$$\text{NINT}(f(x), x, 10, 11) \approx 0.933.$$

So without doing any more measuring, we can predict that about 93.3% of all such spinach boxes will weigh between 10 and 11 ounces.


- (b) For the probability that a box weighs less than 10 ounces, we use the entire area under the curve to the left of $x = 10$. The curve actually approaches the x -axis as an asymptote, but you can see from the graph (Figure 7.50) that $f(x)$ approaches zero quite quickly. Indeed, $f(9)$ is only slightly larger than a billionth. So getting the area from 9 to 10 should do it:

$$\text{NINT}(f(x), x, 9, 10) \approx 0.067.$$

We would expect only about 6.7% of the boxes to weigh less than 10 ounces.

- (c) This would be the integral from 10 to 10, which is zero. This zero probability might seem strange at first, but remember that we are assuming a continuous, unbroken interval of possible spinach weights, and 10 is but one of an infinite number of them.

Now try Exercise 9



[9, 11.5] by [-1, 2.5]

Figure 7.50 The normal pdf for the spinach weights in Example 7. The mean is at the center.

Quick Review 7.5 (For help, go to Section 5.2.)

In Exercises 1–5, find the definite integral by (a) antiderivatives and (b) using NINT.

1. $\int_0^1 e^{-x} dx$

2. $\int_0^1 e^x dx$

3. $\int_{\pi/4}^{\pi/2} \sin x dx$

4. $\int_0^3 (x^2 + 2) dx$

5. $\int_1^2 \frac{x^2}{x^3 + 1} dx$

Exercises 6–10 find, but do not evaluate, the definite integral that the limit as the norms of the partitions go to zero of the Riemann sums on the closed interval $[0, 7]$.

6. $\sum 2\pi(x_k + 2)(\sin x_k)\Delta x$

7. $\sum (1 - x_k^2)(2\pi x_k)\Delta x$

8. $\sum \pi(\cos x_k)^2 \Delta x$

9. $\sum \pi\left(\frac{y_k}{2}\right)^2 (10 - y_k)\Delta y$

10. $\sum \frac{\sqrt{3}}{4} (\sin^2 x_k)\Delta x$

Section 7.5 Exercises

Exercises 1–4, find the work done by the force of $F(x)$ newtons along the x -axis from $x = a$ meters to $x = b$ meters.

1. $F(x) = xe^{-x/3}$, $a = 0$, $b = 5$

2. $F(x) = x \sin(\pi x/4)$, $a = 0$, $b = 3$

3. $F(x) = x\sqrt{9 - x^2}$, $a = 0$, $b = 3$

4. $F(x) = e^{\sin x} \cos x + 2$, $a = 0$, $b = 10$

5. **Leaky Bucket** The workers in Example 2 changed to a larger bucket that held 50 L (490 N) of water, but the new bucket had an even larger leak so that it too was empty by the time it reached the top. Assuming the water leaked out at a steady rate, how much work was done lifting the water to a point 20 meters above the ground? (Do not include the rope and bucket.)

6. **Leaky Bucket** The bucket in Exercise 5 is hauled up more quickly so that there is still 10 L (98 N) of water left when the bucket reaches the top. How much work is done lifting the water this time? (Do not include the rope and bucket.)

7. **Leaky Sand Bag** A bag of sand originally weighing 144 lb was lifted at a constant rate. As it rose, sand leaked out at a constant rate. The sand was half gone by the time the bag had been lifted 18 ft. How much work was done lifting the sand this far? (Neglect the weights of the bag and lifting equipment.)

8. **Stretching a Spring** A spring has a natural length of 10 in. An 800-lb force stretches the spring to 14 in.

- (a) Find the force constant.
- (b) How much work is done in stretching the spring from 10 in. to 12 in.?
- (c) How far beyond its natural length will a 1600-lb force stretch the spring?

9. **Subway Car Springs** It takes a force of 21,714 lb to compress a coil spring assembly on a New York City Transit Authority subway car from its free height of 8 in. to its fully compressed height of 5 in.

- (a) What is the assembly’s force constant?

- (b) How much work does it take to compress the assembly the first half inch? the second half inch? Answer to the nearest inch-pound.


(Source: Data courtesy of Bombardier, Inc., Mass Transit Division, for spring assemblies in subway cars delivered to the New York City Transit Authority from 1985 to 1987.)

10. **Bathroom Scale** A bathroom scale is compressed 1/16 in. when a 150-lb person stands on it. Assuming the scale behaves like a spring that obeys Hooke’s Law,

- (a) how much does someone who compresses the scale 1/8 in. weigh?
- (b) how much work is done in compressing the scale 1/8 in.?

11. **Hauling a Rope** A mountain climber is about to haul up a 50-m length of hanging rope. How much work will it take if the rope weighs 0.624 N/m?

12. **Compressing Gas** Suppose that gas in a circular cylinder of cross section area A is being compressed by a piston (see figure).



- (a) If p is the pressure of the gas in pounds per square inch and V is the volume in cubic inches, show that the work done in compressing the gas from state (p_1, V_1) to state (p_2, V_2) is given by the equation

$$\text{Work} = \int_{(p_1, V_1)}^{(p_2, V_2)} p dV \text{ in} \cdot \text{lb},$$

where the force against the piston is pA .


- (b) Find the work done in compressing the gas from $V_1 = 243 \text{ in}^3$ to $V_2 = 32 \text{ in}^3$ if $p_1 = 50 \text{ lb/in}^2$ and p and V obey the gas law $pV^{1.4} = \text{constant}$ (for adiabatic processes).

Group Activity In Exercises 13–16, the vertical end of a tank containing water (blue shading) weighing 62.4 lb/ft^3 has the given shape.


(a) **Writing to Learn** Explain how to approximate the force against the end of the tank by a Riemann sum.

(b) Find the force as an integral and evaluate it.


13. semicircle




14. semiellipse




15. triangle



16. parabola



17. **Pumping Water** The rectangular tank shown here, with its top at ground level, is used to catch runoff water. Assume that the water weighs 62.4 lb/ft^3 .



(a) How much work does it take to empty the tank by pumping the water back to ground level once the tank is full?


(b) If the water is pumped to ground level with a $(5/11)$ -horsepower motor (work output $250 \text{ ft} \cdot \text{lb/sec}$), how long will it take to empty the full tank (to the nearest minute)?

(c) Show that the pump in part (b) will lower the water level 10 ft (halfway) during the first 25 min of pumping.


(d) **The Weight of Water** Because of differences in the strength of Earth's gravitational field, the weight of a cubic foot of water at sea level can vary from as little as 62.26 lb at the equator to as much as 62.59 lb near the poles, a variation of about 0.5%. A cubic foot of water that weighs 62.4 lb in Melbourne or New York City will weigh 62.5 lb in Juneau or Stockholm. What are the answers to parts (a) and (b) in a location where water weighs 62.26 lb/ft^3 ? 62.5 lb/ft^3 ?

18. **Emptying a Tank** A vertical right cylindrical tank measures 30 ft high and 20 ft in diameter. It is full of kerosene weighing 51.2 lb/ft^3 . How much work does it take to pump the kerosene to the level of the top of the tank?

19. **Writing to Learn** The cylindrical tank shown here is to be filled by pumping water from a lake 15 ft below the bottom of the tank. There are two ways to go about this. One is to pump the water through a hose attached to a valve in the bottom of the tank. The other is to attach the hose to the rim of the tank and let the water pour in. Which way will require less work? Give reasons for your answer.




20. **Drinking a Milkshake** The truncated conical container shown here is full of strawberry milkshake that weighs $(4/9) \text{ oz/in}^3$. As you can see, the container is 7 in. deep, 2.5 in. across at the base, and 3.5 in. across at the top (a standard size at Brigham's in Boston). The straw sticks up an inch above the top. About how much work does it take to drink the milkshake through the straw (neglecting friction)? Answer in inch-ounces.



21. **Revisiting Example 3** How much work will it take to pump the oil in Example 3 to a level 3 ft above the cone's rim?


Pumping Milk Suppose the conical tank in Example 3 contains milk weighing 64.5 lb/ft^3 instead of olive oil. How much work will it take to pump the contents to the rim?

Writing to Learn You are in charge of the evacuation and repair of the storage tank shown here. The tank is a hemisphere of radius 10 ft and is full of benzene weighing 56 lb/ft^3 .



A firm you contacted says it can empty the tank for $1/2$ cent per foot-pound of work. Find the work required to empty the tank by pumping the benzene to an outlet 2 ft above the tank. If you have budgeted \$5000 for the job, can you afford to hire the firm?

Water Tower Your town has decided to drill a well to increase its water supply. As the town engineer, you have determined that a water tower will be necessary to provide the pressure needed for distribution, and you have designed the system shown here. The water is to be pumped from a 300-ft well through a vertical 4-in. pipe into the base of a cylindrical tank 20 ft in diameter and 25 ft high. The base of the tank will be 60 ft above ground. The pump is a 3-hp pump, rated at $1650 \text{ ft} \cdot \text{lb/sec}$. To the nearest hour, how long will it take to fill the tank the first time? (Include the time it takes to fill the pipe.) Assume water weighs 62.4 lb/ft^3 .



Fish Tank A rectangular freshwater fish tank with base $2 \times 4 \text{ ft}$ and height 2 ft (interior dimensions) is filled to within 2 in. of the top.

(a) Find the fluid force against each end of the tank.

(b) Suppose the tank is sealed and stood on end (without spilling) so that one of the square ends is the base. What does that do to the fluid forces on the rectangular sides?

26. **Milk Carton** A rectangular milk carton measures 3.75 in. by 3.75 in. at the base and is 7.75 in. tall. Find the force of the milk (weighing 64.5 lb/ft^3) on one side when the carton is full.

27. Find the probability that a clock stopped between 1:00 and 5:00.

28. Find the probability that a clock stopped between 3:00 and 6:00.

29. Suppose a telephone help line takes a mean of 2 minutes to answer calls. If the standard deviation is $\sigma = 2$, what percentage of the calls are answered in the range of 0 to 4 minutes?

30. **Test Scores** The mean score on a national aptitude test is 498 with a standard deviation of 100 points.

(a) What percentage of the population has scores between 400 and 500?

(b) If we sample 300 test-takers at random, about how many should have scores above 700?

31. **Heights of Females** The mean height of an adult female in New York City is estimated to be 63.4 inches with a standard deviation of 3.2 inches. What proportion of the adult females in New York City are


(a) less than 63.4 inches tall?

(b) between 63 and 65 inches tall?

(c) taller than 6 feet?

(d) exactly 5 feet tall?

32. **Writing to Learn** Exercises 30 and 31 are subtly different, in that the heights in Exercise 31 are measured *continuously* and the scores in Exercise 30 are measured *discretely*. The discrete probabilities determine rectangles above the individual test scores, so that there actually is a nonzero probability of scoring, say, 560. The rectangles would look like the figure below, and would have total area 1.



Explain why integration gives a good estimate for the probability, even in the discrete case.

33. **Writing to Learn** Suppose that $f(t)$ is the probability density function for the lifetime of a certain type of lightbulb where t is in hours. What is the meaning of the integral

$$\int_{100}^{800} f(t) dt?$$

Standardized Test Questions

You may use a graphing calculator to solve the following problems.

34. **True or False** A force is applied to compress a spring several inches. Assume the spring obeys Hooke's Law. Twice as much work is required to compress the spring the second inch than is required to compress the spring the first inch. Justify your answer.

35. **True or False** An aquarium contains water weighing 62.4 lb/ft³. The aquarium is in the shape of a cube where the length of each edge is 3 ft. Each side of the aquarium is engineered to withstand 1000 pounds of force. This should be sufficient to withstand the force from water pressure. Justify your answer.
36. **Multiple Choice** A force of $F(x) = 350x$ newtons moves a particle along a line from $x = 0$ m to $x = 5$ m. Which of the following gives the best approximation of the work done by the force?
 (A) 1750 J (B) 2187.5 J (C) 2916.67 J
 (D) 3281.25 J (E) 4375 J
37. **Multiple Choice** A leaky bag of sand weighs 50 n. It is lifted from the ground at a constant rate, to a height of 20 m above the ground. The sand leaks at a constant rate and just finishes draining as the bag reaches the top. Which of the following gives the work done to lift the sand to the top? (Neglect the bag.)
 (A) 50 J (B) 100 J (C) 250 J (D) 500 J (E) 1000 J
38. **Multiple Choice** A spring has a natural length of 0.10 m. A 200-n force stretches the spring to a length of 0.15 m. Which of the following gives the work done in stretching the spring from 0.10 m to 0.15 m?
 (A) 0.05 J (B) 5 J (C) 10 J (D) 200 J (E) 4000 J
39. **Multiple Choice** A vertical right cylindrical tank measures 12 ft high and 16 ft in diameter. It is full of water weighing 62.4 lb/ft³. How much work does it take to pump the water to the level of the top of the tank? Round your answer to the nearest ft-lb.
 (A) 149,490 ft-lb
 (B) 285,696 ft-lb
 (C) 360,240 ft-lb
 (D) 448,776 ft-lb
 (E) 903,331 ft-lb

Extending the Ideas

40. **Putting a Satellite into Orbit** The strength of Earth's gravitational field varies with the distance r from Earth's center, and the magnitude of the gravitational force experienced by a satellite of mass m during and after launch is

$$F(r) = \frac{mMG}{r^2}$$

Here, $M = 5.975 \times 10^{24}$ kg is Earth's mass, $G = 6.6726 \times 10^{-11}$ N · m²kg⁻² is the *universal gravitational constant*, and r is measured in meters. The work it takes to lift a 1000-kg satellite from Earth's surface to a circular orbit 35,780 km above Earth's center is therefore given by the integral

$$\text{Work} = \int_{6,370,000}^{35,780,000} \frac{1000MG}{r^2} dr \text{ joules.}$$

The lower limit of integration is Earth's radius in meters at the launch site. Evaluate the integral. (This calculation does not take into account energy spent lifting the launch vehicle or energy spent bringing the satellite to orbit velocity.)

41. **Forcing Electrons Together** Two electrons r meters apart repel each other with a force of

$$F = \frac{23 \times 10^{-29}}{r^2} \text{ newton.}$$

- (a) Suppose one electron is held fixed at the point $(1, 0)$ on the x -axis (units in meters). How much work does it take to move a second electron along the x -axis from the point $(-1, 0)$ to the origin?
 (b) Suppose an electron is held fixed at each of the points $(-1, 0)$ and $(1, 0)$. How much work does it take to move a third electron along the x -axis from $(5, 0)$ to $(3, 0)$?

42. **Kinetic Energy** If a variable force of magnitude $F(x)$ moves a body of mass m along the x -axis from x_1 to x_2 , the body's velocity v can be written as dx/dt (where t represents time). Use Newton's second law of motion, $F = m(dv/dt)$, and the Chain Rule

$$\frac{dv}{dt} = \frac{dv}{dx} \frac{dx}{dt} = v \frac{dv}{dx}$$

to show that the net work done by the force in moving the body from x_1 to x_2 is

$$W = \int_{x_1}^{x_2} F(x) dx = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2, \quad (1)$$

where v_1 and v_2 are the body's velocities at x_1 and x_2 . In physics the expression $(1/2)mv^2$ is the *kinetic energy* of the body moving with velocity v . Therefore, *the work done by the force equals the change in the body's kinetic energy*, and we can find the work by calculating this change.

Weight vs. Mass

Weight is the force that results from gravity pulling on a mass. The two are related by the equation in Newton's second law,

$$\text{weight} = \text{mass} \times \text{acceleration.}$$

Thus,

$$\text{newtons} = \text{kilograms} \times \text{m/sec}^2,$$

$$\text{pounds} = \text{slugs} \times \text{ft/sec}^2.$$

To convert mass to weight, multiply by the acceleration of gravity. To convert weight to mass, divide by the acceleration of gravity.

In Exercises 43–49, use Equation 1 from Exercise 42.

43. **Tennis** A 2-oz tennis ball was served at 160 ft/sec (about 109 mph). How much work was done on the ball to make it go this fast?
 44. **Baseball** How many foot-pounds of work does it take to throw a baseball 90 mph? A baseball weighs 5 oz = 0.3125 lb.
 45. **Golf** A 1.6-oz golf ball is driven off the tee at a speed of 280 ft/sec (about 191 mph). How many foot-pounds of work are done getting the ball into the air?
 46. **Tennis** During the match in which Pete Sampras won the 1990 U.S. Open men's tennis championship, Sampras hit a serve that was clocked at a phenomenal 124 mph. How much work did Sampras have to do on the 2-oz ball to get it to that speed?

47. **Football** A quarterback threw a 14.5-oz football 88 ft/sec (60 mph). How many foot-pounds of work were done on the ball to get it to that speed?
 48. **Softball** How much work has to be performed on a 6.5-oz softball to pitch it at 132 ft/sec (90 mph)?

Quick Quiz for AP* Preparation: Sections 7.4 and 7.5


1. You should solve the following problems without using a graphing calculator.
 1. **Multiple Choice** The length of a curve from $x = 0$ to $x = 1$ is given by $\int_0^1 \sqrt{1 + 16x^6} dx$. If the curve contains the point $(1, 4)$, which of the following could be an equation for this curve?
 (A) $y = x^4 + 3$
 (B) $y = x^4 + 1$
 (C) $y = 1 + 16x^6$
 (D) $y = \sqrt{1 + 16x^6}$
 (E) $y = x + \frac{x^7}{7}$
 2. **Multiple Choice** Which of the following gives the length of the path described by the parametric equations $x = \frac{1}{4}t^4$ and $y = t^3$, where $0 \leq t \leq 2$?

- (A) $\int_0^2 t^6 + 9t^4 dt$
 (B) $\int_0^2 \sqrt{t^6 + 1} dt$
 (C) $\int_0^2 \sqrt{1 + 9t^4} dt$
 (D) $\int_0^2 \sqrt{t^6 + 9t^4} dt$
 (E) $\int_0^2 \sqrt{t^3 + 3t^2} dt$

3. **Multiple Choice** The base of a solid is a circle of radius 2 inches. Each cross section perpendicular to a certain diameter is a square with one side lying in the circle. The volume of the solid in cubic inches is

- (A) 16 (B) 16π (C) $\frac{128}{3}$ (D) $\frac{128\pi}{3}$ (E) 32π


49. **A Ball Bearing** A 2-oz steel ball bearing is placed on a vertical spring whose force constant is $k = 18$ lb/ft. The spring is compressed 3 in. and released. About how high does the ball bearing go? (Hint: The kinetic (compression) energy, mgh , of a spring is $\frac{1}{2}ks^2$, where s is the distance the spring is compressed, m is the mass, g is the acceleration of gravity, and h is the height.)



The front of the tank can be partitioned into narrow horizontal bands of height Δh . The force exerted by the water on a band at depth h_i is approximately

$$\text{pressure} \cdot \text{area} = 62.4h_i \cdot 2\Delta h.$$

- (a) Write the Riemann sum that approximates the force exerted on the entire front of the tank.
 (b) Use the Riemann sum from part (a) to write and evaluate a definite integral that gives the force exerted on the front of the tank. Include correct units.
 (c) Find the total force exerted on the front of the tank if the front (and back) are semicircles with diameter 2 ft. Include correct units.



Calculus at Work

I am working toward becoming an archaeoastronomer and ethnoastronomer of Africa. I have a Bachelor's degree in Physics, a Master's degree in Astronomy, and a Ph.D. in Astronomy and Astrophysics. From 1988 to 1990 I was a member of the Peace Corps, and I taught mathematics to high school students in the Fiji Islands. Calculus is a required course in high schools there.

For my Ph.D. dissertation, I investigated the possibility of the birthrate of stars being related to the composition of star formation clouds. I collected data on the absorption of electromagnetic emissions emanating from these regions. The intensity of emissions graphed versus wave-

length produces a flat curve with downward spikes at the characteristic wavelengths of the elements present. An estimate of the area between a spike and the flat curve results in a concentration in molecules/cm³ of an element. This area is the difference in the integrals of the flat and spike curves. In particular, I was looking for a large concentration of water-ice, which increases the probability of planets forming in a region.

Currently, I am applying for two research grants. One will allow me to use the NASA infrared telescope on Mauna Kea to search for C₃S₂ in comets. The other will help me study the history of astronomy in Tunisia.



Jarita Holbrook
Los Angeles, CA

Chapter 7 Key Terms

- arc length (p. 413)
- area between curves (p. 390)
- Cavalieri's theorems (p. 404)
- center of mass (p. 389)
- constant-force formula (p. 384)
- cylindrical shells (p. 402)
- displacement (p. 380)
- fluid force (p. 421)
- fluid pressure (p. 421)
- foot-pound (p. 384)
- force constant (p. 385)
- Gaussian curve (p. 423)
- Hooke's Law (p. 385)
- inflation rate (p. 388)
- joule (p. 384)
- length of a curve (p. 413)
- mean (p. 423)
- moment (p. 389)
- net change (p. 379)
- newton (p. 384)
- normal curve (p. 423)
- normal pdf (p. 423)
- probability density function (pdf) (p. 422)
- 68-95-99.7 rule (p. 423)

- smooth curve (p. 413)
- smooth function (p. 413)
- solid of revolution (p. 400)
- standard deviation (p. 423)
- surface area (p. 405)
- total distance traveled (p. 381)
- universal gravitational constant (p. 428)
- volume by cylindrical shells (p. 402)
- volume by slicing (p. 400)
- volume of a solid (p. 399)
- weight-density (p. 421)
- work (p. 384)

Chapter 7 Review Exercises

The collection of exercises marked in red could be used as a chapter test. In Exercises 1–5, the application involves the accumulation of small changes over an interval to give the net change over that entire interval. Set up an integral to model the accumulation and evaluate it to answer the question.

1. A toy car slides down a ramp and coasts to a stop after 5 sec. Its velocity from $t = 0$ to $t = 5$ is modeled by $v(t) = t^2 - 0.2t^3$ ft/sec. How far does it travel?

2. The fuel consumption of a diesel motor between weekly maintenance periods is modeled by the function $c(t) = 4 + 0.001t^4$ gal/day, $0 \leq t \leq 7$. How many gallons does it consume in a week?
3. The number of billboards per mile along a 100-mile stretch of interstate highway approaching a certain city is modeled by the function $B(x) = 21 - e^{0.03x}$, where x is the distance from the city in miles. About how many billboards are along that stretch of highway?

4. A 2-meter rod has a variable density modeled by the function $p(x) = 11 - 4x$ g/m, where x is the distance in meters from the base of the rod. What is the total mass of the rod?


5. The electrical power consumption (measured in kilowatts) at a factory t hours after midnight during a typical day is modeled by $E(t) = 300(2 - \cos(\pi t/12))$. How many kilowatt-hours of electrical energy does the company consume in a typical day?

- 6–19. Exercises 6–19, find the area of the region enclosed by the lines and curves.

6. $y = x, y = 1/x^2, x = 2$

7. $y = x + 1, y = 3 - x^2$

8. $\sqrt{x} + \sqrt{y} = 1, x = 0, y = 0$



9. $x = 2y^2, x = 0, y = 3$

10. $4x = y^2 - 4, 4x = y + 16$


11. $y = \sin x, y = x, x = \pi/4$

12. $y = 2 \sin x, y = \sin 2x, 0 \leq x \leq \pi$

13. $y = \cos x, y = 4 - x^2$

14. $y = \sec^2 x, y = 3 - |x|$

15. **The Necklace** one of the smaller bead-shaped regions enclosed by the graphs of $y = 1 + \cos x$ and $y = 2 - \cos x$




$[-4\pi, 4\pi]$ by $[-4, 8]$

16. one of the larger bead-shaped regions enclosed by the curves in Exercise 15

17. **The Bow Tie** the region enclosed by the graphs of

$$y = x^3 - x \quad \text{and} \quad y = \frac{x}{x^2 + 1}$$


(shown in the next column).



$[-2, 2]$ by $[-1.5, 1.5]$


18. **The Bell** the region enclosed by the graphs of

$$y = 3^{1-x^2} \quad \text{and} \quad y = \frac{x^2 - 3}{10}$$



$[-4, 4]$ by $[-2, 3.5]$


19. **The Kissing Fish** the region enclosed between the graphs of $y = x \sin x$ and $y = -x \sin x$ over the interval $[-\pi, \pi]$



$[-5, 5]$ by $[-3, 3]$

20. Find the volume of the solid generated by revolving the region bounded by the x -axis, the curve $y = 3x^4$, and the lines $x = -1$ and $x = 1$ about the x -axis.
21. Find the volume of the solid generated by revolving the region enclosed by the parabola $y^2 = 4x$ and the line $y = x$ about
 - (a) the x -axis.
 - (b) the y -axis.
 - (c) the line $x = 4$.
 - (d) the line $y = 4$.
22. The section of the parabola $y = x^2/2$ from $y = 0$ to $y = 2$ is revolved about the y -axis to form a bowl.
 - (a) Find the volume of the bowl.
 - (b) Find how much the bowl is holding when it is filled to a depth of k units ($0 < k < 2$).
 - (c) If the bowl is filled at a rate of 2 cubic units per second, how fast is the depth k increasing when $k = 1$?


23. The profile of a football resembles the ellipse shown here (all dimensions in inches). Find the volume of the football to the nearest cubic inch.



24. The base of a solid is the region enclosed between the graphs of $y = \sin x$ and $y = -\sin x$ from $x = 0$ to $x = \pi$. Each cross section perpendicular to the x -axis is a semicircle with diameter connecting the two graphs. Find the volume of the solid.
25. The region enclosed by the graphs of $y = e^{x/2}$, $y = 1$, and $x = \ln 3$ is revolved about the x -axis. Find the volume of the solid generated.
26. A round hole of radius $\sqrt{3}$ feet is bored through the center of a sphere of radius 2 feet. Find the volume of the piece cut out.
27. Find the length of the arch of the parabola $y = 9 - x^2$ that lies above the x -axis.
28. Find the perimeter of the bow-tie-shaped region enclosed between the graphs of $y = x^3 - x$ and $y = x - x^3$.
29. A particle travels at 2 units per second along the curve $y = x^3 - 3x^2 + 2$. How long does it take to travel from the local maximum to the local minimum?
30. **Group Activity** One of the following statements is true for all $k > 0$ and one is false. Which is which? Explain.
 (a) The graphs of $y = k \sin x$ and $y = \sin kx$ have the same length on the interval $[0, 2\pi]$.
 (b) The graph of $y = k \sin x$ is k times as long as the graph of $y = \sin x$ on the interval $[0, 2\pi]$.
31. Let $F(x) = \int_1^x \sqrt{t^4 - 1} dt$. Find the exact length of the graph of F from $x = 2$ to $x = 5$ without using a calculator.
32. **Rock Climbing** A rock climber is about to haul up 100 N (about 22.5 lb) of equipment that has been hanging beneath her on 40 m of rope weighing 0.8 N/m. How much work will it take to lift
 (a) the equipment? (b) the rope?
 (c) the rope and equipment together?

33. **Hauling Water** You drove an 800-gallon tank truck from the base of Mt. Washington to the summit and discovered on arrival that the tank was only half full. You had started out with a full tank of water, had climbed at a steady rate, and had taken 50 minutes to accomplish the 4750-ft elevation change. Assuming that the water leaked out at a steady rate, how much work was spent in carrying the water to the summit? Water weighs 8 lb/gal. (Do not count the work done getting you and the truck to the top.)


34. **Stretching a Spring** If a force of 80 N is required to hold a spring 0.3 m beyond its unstressed length, how much work does it take to stretch the spring this far? How much work does it take to stretch the spring an additional meter?
35. **Writing to Learn** It takes a lot more effort to roll a stone up a hill than to roll the stone down the hill, but the weight of the stone and the distance it covers are the same. Does this mean that the same amount of work is done? Explain.
36. **Emptying a Bowl** A hemispherical bowl with radius 8 inches is filled with punch (weighing 0.04 pound per cubic inch) to within 2 inches of the top. How much work is done emptying the bowl if the contents are pumped just high enough to get over the rim?
37. **Fluid Force** The vertical triangular plate shown below is the end plate of a feeding trough full of hog slop, weighing 80 pounds per cubic foot. What is the force against the plate?



38. **Fluid Force** A standard olive oil can measures 5.75 in. by 3.5 in. by 10 in. Find the fluid force against the base and each side of the can when it is full. (Olive oil has a weight-density of 57 pounds per cubic foot.)



39. **Volume** A solid lies between planes perpendicular to the x -axis at $x = 0$ and $x = 6$. The cross sections between the planes are squares whose bases run from the x -axis up to the curve $\sqrt{x} + \sqrt{y} = \sqrt{6}$. Find the volume of the solid.



40. **Yellow Perch** A researcher measures the lengths of 3-year-old yellow perch in a fish hatchery and finds that they have a mean length of 17.2 cm with a standard deviation of 3.4 cm. What proportion of 3-year-old yellow perch raised under similar conditions can be expected to reach a length of 20 cm or more?

41. **Group Activity** Using as large a sample of classmates as possible, measure the span of each person's fully stretched hand, from the tip of the pinky finger to the tip of the thumb. Based on the mean and standard deviation of your sample, what percentage of students your age would have a finger span of more than 10 inches?
42. **The 68-95-99.7 Rule** (a) Verify that for every normal pdf, the proportion of the population lying within one standard deviation of the mean is close to 68%. (Hint: Since it is the same for every pdf, you can simplify the function by assuming that $\mu = 0$ and $\sigma = 1$. Then integrate from -1 to 1 .)
 (b) Verify the two remaining parts of the rule.
43. **Writing to Learn** Explain why the area under the graph of a probability density function has to equal 1.

Exercises 44–48, use the cylindrical shell method to find the volume of the solid generated by revolving the region bounded by the curves about the y -axis.

44. $y = 2x$, $y = x/2$, $x = 1$
 45. $y = 1/x$, $y = 0$, $x = 1/2$, $x = 2$
 46. $y = \sin x$, $y = 0$, $0 \leq x \leq \pi$
 47. $y = x - 3$, $y = x^2 - 3x$
 48. the bell-shaped region in Exercise 18

49. **Bundt Cake** A bundt cake (see Exploration 1, Section 7.3) has a hole of radius 2 inches and an outer radius of 6 inches at the base. It is 5 inches high, and each cross-sectional slice is parabolic.

- (a) Model a typical slice by finding the equation of the parabola with y -intercept 5 and x -intercepts ± 2 .
 (b) Revolve the parabolic region about an appropriate line to generate the bundt cake and find its volume.

50. **Finding a Function** Find a function f that has a continuous derivative on $(0, \infty)$ and that has both of the following properties.
 i. The graph of f goes through the point $(1, 1)$.
 ii. The length L of the curve from $(1, 1)$ to any point $(x, f(x))$ is given by the formula $L = \ln x + f(x) - 1$.

Exercises 51 and 52, find the area of the surface generated by revolving the curve about the indicated axis.

51. $y = \tan x$, $0 \leq x \leq \pi/4$; x -axis
 52. $xy = 1$, $1 \leq y \leq 2$; y -axis

AP* Examination Preparation

You may use a graphing calculator to solve the following problems.

53. Let R be the region in the first quadrant enclosed by the y -axis and the graphs of $y = 2 + \sin x$ and $y = \sec x$.
 (a) Find the area of R .
 (b) Find the volume of the solid generated when R is revolved about the x -axis.
 (c) Find the volume of the solid whose base is R and whose cross sections cut by planes perpendicular to the x -axis are squares.
54. The temperature outside a house during a 24-hour period is given by

$$F(t) = 80 - 10 \cos\left(\frac{\pi t}{12}\right), 0 \leq t \leq 24,$$

where $F(t)$ is measured in degrees Fahrenheit and t is measured in hours.

- (a) Find the average temperature, to the nearest degree Fahrenheit, between $t = 6$ and $t = 14$.
 (b) An air conditioner cooled the house whenever the outside temperature was at or above 78 degrees Fahrenheit. For what values of t was the air conditioner cooling the house?
 (c) The cost of cooling the house accumulates at the rate of \$0.05 per hour for each degree the outside temperature exceeds 78 degrees Fahrenheit. What was the total cost, to the nearest cent, to cool the house for this 24-hour period?

55. The rate at which people enter an amusement park on a given day is modeled by the function E defined by

$$E(t) = \frac{15600}{t^2 - 24t + 160}.$$

The rate at which people leave the same amusement park on the same day is modeled by the function L defined by

$$L(t) = \frac{9890}{t^2 - 38t + 370}.$$

Both $E(t)$ and $L(t)$ are measured in people per hour, and time t is measured in hours after midnight. These functions are valid for $9 \leq t \leq 23$, which are the hours that the park is open. At time $t = 9$, there are no people in the park.

- (a) How many people have entered the park by 5:00 P.M. ($t = 17$)? Round your answer to the nearest whole number.
 (b) The price of admission to the park is \$15 until 5:00 P.M. ($t = 17$). After 5:00 P.M., the price of admission to the park is \$11. How many dollars are collected from admissions to the park on the given day? Round your answer to the nearest whole number.
 (c) Let $H(t) = \int_9^t (E(x) - L(x)) dx$ for $9 \leq t \leq 23$. The value of $H(17)$ to the nearest whole number is 3725. Find the value of $H'(17)$ and explain the meaning of $H(17)$ and $H'(17)$ in the context of the park.
 (d) At what time t , for $9 \leq t \leq 23$, does the model predict that the number of people in the park is a maximum?