

19. a) $240 + 250 + 260 + \dots + 300$
 b) $S_n = 235n + 5n^2$
 c) 1890
 d) Nathan will continue to remove an extra 10 bushels per hour.
20. $(-27) + (-22) + (-17)$
21. Jeanette and Pierre have used two different forms of the same formula. Jeanette has replaced t_n with $t_1 + (n - 1)d$.

22. a) 100
 b) $S_{\text{green}} = 1 + 2 + 3 + \dots + 10$
 $S_{\text{blue}} = 0 + 1 + 2 + 3 + \dots + 9$
 $S_{\text{total}} = S_{\text{green}} + S_{\text{blue}}$
 $S_{\text{total}} = \frac{10}{2}(1 + 10) + \frac{10}{2}(0 + 9)$
 $S_{\text{total}} = 5(11) + 5(9)$
 $S_{\text{total}} = 55 + 45$
 $S_{\text{total}} = 100$

23. a) 55
 b) The n th triangular number is represented by S_n .
 $S_n = \frac{n}{2}[2t_1 + (n - 1)d]$
 $S_n = \frac{n}{2}[2(1) + (n - 1)(1)]$
 $S_n = \frac{n}{2}[2 + (n - 1)]$
 $S_n = \frac{n}{2}(1 + n)$

1.3 Geometric Sequences, pages 39 to 45

1. a) geometric; $r = 2$; $t_n = 2^{n-1}$
 b) not geometric
 c) geometric; $r = -3$; $t_n = 3(-3)^{n-1}$
 d) not geometric
 e) geometric; $r = 1.5$; $t_n = 10(1.5)^{n-1}$
 f) geometric; $r = 5$; $t_n = -1(5)^{n-1}$

2.

	Geometric Sequence	Common Ratio	6th Term	10th Term
a)	6, 18, 54, ...	3	1458	118 098
b)	1.28, 0.64, 0.32, ...	0.5	0.04	0.0025
c)	$\frac{1}{5}, \frac{3}{5}, \frac{9}{5}, \dots$	3	$\frac{243}{5}$	$\frac{19\,683}{5}$

3. a) 2, 6, 18, 54 b) -3, 12, -48, 192
 c) 4, -12, 36, -108 d) 2, 1, $\frac{1}{2}, \frac{1}{4}$
4. 18.9, 44.1, 102.9
5. a) $t_n = 3(2)^{n-1}$ b) $t_n = 192\left(-\frac{1}{4}\right)^{n-1}$
 c) $t_n = \frac{5}{9}(3)^{n-1}$ d) $t_n = 4(2)^{n-1}$
6. a) 4 b) 7 c) 5
 d) 6 e) 9 f) 8

7. 37

8. 16, 12, 9; $t_n = 16\left(\frac{3}{4}\right)^{n-1}$

9. a) $t_1 = 3$; $r = 0.75$
 b) $t_n = 3(0.75)^{n-1}$
 c) approximately 53.39 cm
 d) 7
10. a) 95%
 b) 100, 95, 90.25, 85.7375
 c) 0.95
 d) about 59.87%
 e) After 27 washings, 25% of the original colour would remain in the jeans. Example: The geometric sequence continues for each washing.
11. 1.77
12. a) 1, 2, 4, 8, 16 b) $t_n = 1(2)^{n-1}$
 c) 2^{29} or 536 870 912
13. a) 1.031 b) 216.3 cm
 c) 56 jumps
14. a) 1, 2, 4, 8, 16, 32 b) $t_n = 1(2)^{n-1}$
 c) 2^{25} or 33 554 432
 d) All cells continue to double and all cells live.
15. 2.9%
16. 8 weeks
17. 65.2 m
18. 0.920
19. a) 76.0 mL b) 26 h

20. a)

Time, d (days)	Charge Level, C (%)
0	100
1	98
2	96.04
3	94.12

- b) $t_n = 100(0.98)^{n-1}$
 c) The formula in part b) includes the first term at $d = 0$ in the sequence. The formula $C = 100(0.98)^n$ does not consider the first term of the sequence.
 d) 81.7%
21. a) 24.14 mm b) 1107.77 mm
22. Example: If a, b, c are terms of an arithmetic sequence, then $b - a = c - b$. If $6^a, 6^b, 6^c$ are terms of a geometric series, then $\frac{6^b}{6^a} = \frac{6^c}{6^b}$ and $6^{b-a} = 6^{c-b}$. Therefore, $b - a = c - b$. So, when $6^a, 6^b, 6^c$ form a geometric sequence, then a, b, c form an arithmetic sequence.
23. $\frac{5}{3}$; 9, 15, 25
24. a) 23.96 cm b) 19.02 cm
 c) 2.13 cm d) 2.01 cm
 e) 2.01, 1.90, 1.79; arithmetic; $d = -0.11$ cm
25. Mala's solution is correct. Since the aquarium loses 8% of the water every day, it maintains 92% of the water every day.

26.


	$\frac{1}{500}$			$\frac{50}{3}$				
	$\frac{1}{100}$	$\frac{1}{10}$	1	10				
	$\frac{1}{20}$		2	6	18	54		
$\frac{1}{16}$	$\frac{1}{4}$	1	4			9		
	$\frac{5}{4}$		8			$\frac{3}{2}$		
	$\frac{25}{4}$		16	4	1	$\frac{1}{4}$	$\frac{1}{16}$	$\frac{1}{64}$
	$\frac{125}{4}$		32					
	$\frac{625}{4}$	100	64					

27. a) 0.86 cm^2 b) 1.72 cm^2
 c) 3.43 cm^2 d) 109.88 cm^2

1.4 Geometric Series, pages 53 to 57

1. a) geometric; $r = 6$ b) geometric; $r = -\frac{1}{2}$
 c) not geometric d) geometric; $r = 1.1$
2. a) $t_1 = 6, r = 1.5, S_{10} = \frac{174\,075}{256}, S_{10} \approx 679.98$
 b) $t_1 = 18, r = -0.5, S_{12} = \frac{12\,285}{1024}, S_{12} \approx 12.00$
 c) $t_1 = 2.1, r = 2, S_9 = \frac{10\,731}{10}, S_9 = 1073.10$
 d) $t_1 = 0.3, r = 0.01, S_{12} = \frac{10}{33}, S_{12} \approx 0.30$
3. a) 12 276 b) $\frac{3280}{81}$
 c) $-\frac{209\,715}{256}$ d) $\frac{36\,855}{256}$
4. a) 40.50 b) 0.96
 c) 109 225 d) 39 063
5. a) 3 b) 295.7
6. 7
7. a) 81 b) $81 + 27 + 9 + 3 + 1$
8. $t_2 = -\frac{81}{16}; S_6 = 7.8$
9. a) If the person in charge is included, the series is $1 + 4 + 16 + 64 + \dots$. If the person in charge is not included, the series is $4 + 16 + 64 + \dots$.
 b) If the person in charge is included, the sum is 349 525. If the person in charge is not included, the sum is 1 398 100.

10. 46.4 m





11. 794.3 km

12. b)

Stage Number	Length of Each Line Segment	Number of Line Segments	Perimeter of Snowflake
1	1	3	3
2	$\frac{1}{3}$	12	4
3	$\frac{1}{9}$	48	$\frac{16}{3}$
4	$\frac{1}{27}$	192	$\frac{64}{9}$
5	$\frac{1}{81}$	768	$\frac{256}{27}$

- c) length, $t_n = \left(\frac{1}{3}\right)^{n-1}$;
 number of line segments, $t_n = 3(4)^{n-1}$;
 perimeter, $t_n = 3\left(\frac{4}{3}\right)^{n-1}$
 d) $\frac{1024}{81} \approx 12.64$

13. 98 739
 14. 91 mm
 15. a) 226.9 mg b) 227.3 mg
 16. 8
 17. $\frac{58\,025}{48}$
 18. $a = 5, b = 10, c = 20$ or $a = 20, b = 10, c = 5$
 19. 15
 20. $\frac{341}{4}\pi$
 21.



22. Examples:
 a) All butterflies produce the same number of eggs and all eggs hatch.
 b) No. Tom determined the total number of butterflies from the first to fifth generations. He should have found the fifth term, which would determine the total number of butterflies in the fifth generation only.

- c) This is a reasonable estimate, but it does include all butterflies up to the fifth generation, which is 6.42×10^7 more butterflies than those produced in the fifth generation.
- d) Determine $t_5 = 1(400)^4$ or 2.56×10^{10} .

1.5 Infinite Geometric Series, pages 63 to 65

1. a) divergent b) convergent
 c) convergent d) divergent
 e) divergent
2. a) $\frac{32}{5}$ b) no sum
 c) no sum d) 2
 e) 2.5
3. a) $0.87 + 0.0087 + 0.000\ 087 + \dots$;
 $S_\infty = \frac{87}{99}$ or $\frac{29}{33}$
- b) $0.437 + 0.000\ 437 + \dots$; $S_\infty = \frac{437}{999}$
4. Yes. The sum of the infinite series representing 0.999... is equal to 1.
5. a) 15 b) $\frac{4}{5}$ or 0.8
 c) 14
6. $t_1 = 27$; $27 + 18 + 12 + \dots$
7. $r = \frac{2}{5}$; $-8 - \frac{16}{5} - \frac{32}{25} - \frac{64}{125} - \dots$
8. a) 400 000 barrels of oil
 b) Determining the lifetime production assumes the oil well continues to produce at the same rate for many months. This is an unreasonable assumption because 94% is a high rate to maintain.
9. $x = \frac{1}{4}$; $1 + \frac{3}{4} + \frac{9}{16} + \frac{27}{64} + \dots$
10. $r = \frac{1}{2}$
11. a) $-1 < x < 1$ b) $-3 < x < 3$
 c) $-\frac{1}{2} < x < \frac{1}{2}$
12. 6 cm
13. 250 cm
14. No sum, since $r = 1.1 > 1$. Therefore, the series is divergent.
15. 48 m
16. a) approximately 170.86 cm
 b) 300 cm
17. a) Rita
 b) $r = -\frac{4}{3}$; therefore, $r < -1$, and the series is divergent.
18. 125 m
19. 72 cm
20. a) Example: $\frac{4}{5} + \left(\frac{4}{5}\right)^2 + \left(\frac{4}{5}\right)^3 + \dots + \left(\frac{4}{5}\right)^n$
 and $\frac{1}{5} + \left(\frac{1}{5}\right)^2 + \left(\frac{1}{5}\right)^3 + \dots + \left(\frac{1}{5}\right)^n$

b) $S_\infty = \frac{t_1}{1-r} = \frac{\frac{4}{5}}{1-\frac{4}{5}} = \frac{\frac{4}{5}}{\frac{1}{5}} = 4$ and
 $S_\infty = \frac{t_1}{1-r} = \frac{\frac{1}{5}}{1-\frac{1}{5}} = \frac{\frac{1}{5}}{\frac{4}{5}} = \frac{1}{4}$

21. Geometric series converge only when $-1 < r < 1$.

22. a) $S_n = -\frac{3}{8}n^2 + \frac{11}{8}n$

b) $S_n = \frac{\left(\frac{1}{4}\right)^n - 1}{-\frac{3}{4}}$

$S_n = -\frac{4}{3}\left(\frac{1}{4}\right)^n + \frac{4}{3}$

c) $S_\infty = \frac{1}{1-\frac{1}{4}}$

$S_\infty = \frac{4}{3}$


23. Step 3

n	1	2	3	4
Fraction of Paper	$\frac{1}{4}$	$\frac{1}{16}$	$\frac{1}{64}$	$\frac{1}{256}$

Step 4 $\frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \frac{1}{256}$, Example: $S_\infty = \frac{1}{3}$


Chapter 1 Review, pages 66 to 68

1. a) arithmetic, $d = 4$ b) arithmetic, $d = -5$
 c) not arithmetic d) not arithmetic
2. a) C b) D
 c) E d) B
 e) A
3. a) term, $n = 14$ b) not a term
 c) term, $n = 54$ d) not a term
4. a) A
 b)




In the graph, sequence 1 has a larger positive slope than sequence 2. The value of term 17 is greater in sequence 1 than in sequence 2.

5. $t_{10} = 41$
 6. 306 cm
 7. a) $S_{10} = 195$ b) $S_{12} = 285$
 c) $S_{10} = -75$ d) $S_{20} = 3100$
 8. $S_{40} = 3420$
 9. a) 29 b) 225
 c) 25 days
 10. a) 61 b) 495
 11. 1170
 12. a) not geometric
 b) geometric, $r = -2$, $t_1 = 1$, $t_n = (-2)^{n-1}$
 c) geometric, $r = \frac{1}{2}$, $t_1 = 1$, $t_n = \left(\frac{1}{2}\right)^{n-1}$
 d) not geometric
 13. a) 7346 bacteria
 b) $t_n = 5000(1.08)^n$
 14. 2π cm or approximately 6.28 cm
 15.



16. a) arithmetic b) geometric
 c) geometric d) arithmetic
 e) arithmetic f) geometric
 17. a) $S_{10} = \frac{174\,075}{256}$, $S_{10} \approx 679.98$
 b) $S_{12} = \frac{36\,855}{1024}$, $S_{12} \approx 35.99$
 c) $S_{20} = \frac{20\,000}{3}$, $S_{20} \approx 6666.67$
 d) $S_9 = \frac{436\,905}{4096}$, $S_9 \approx 106.67$
 18. a) 19.1 mm b) 1.37 m
 19. a) $S_\infty = 15$ b) $S_\infty = \frac{3}{4}$
 20. a) convergent, $S_\infty = 16$
 b) divergent
 c) convergent, $S_\infty = -28$
 d) convergent, $S_\infty = \frac{3}{2}$
 21. a) $r = -0.4$
 b) $S_1 = 7$, $S_2 = 4.2$, $S_3 = 5.32$, $S_4 = 4.872$,
 $S_5 = 5.0512$
 c) 5
 d) $S_\infty = 5$



- a) $1, \frac{1}{4}, \frac{1}{16}, \frac{1}{64}$. Yes. The areas form a geometric sequence. The common ratio is $\frac{1}{4}$.
 b) $1\frac{21}{64}$ or 1.328 125 square units
 c) $\frac{4}{3}$ square units
 23. a) A series is geometric if there is a common ratio r such that $r \neq 1$.
 An infinite geometric series converges if $-1 < r < 1$.
 An infinite geometric series diverges if $r < -1$ or $r > 1$.
 b) Example:
 $4 + 2 + 1 + 0.5 + \dots$; $S_\infty = 8$
 $21 - 10.5 + 5.25 - 2.625 + \dots$; $S_\infty = 14$

Chapter 1 Practice Test, pages 69 to 70

1. D
 2. B
 3. B
 4. B
 5. C
 6. 11.62 cm
 7. Arithmetic sequences form straight-line graphs, where the slope is the common difference of the sequence. Geometric sequences form curved graphs.
 8. $A = 15$, $B = 9$
 9. 0.7 km
 10. a) 5, 36, 67, 98, 129, 160
 b) $t_n = 31n - 26$
 c) 5, 10, 20, 40, 80, 160
 d) $t_n = 5(2)^{n-1}$
 11. a) 17, 34, 51, 68, 85
 b) $t_n = 17n$
 c) 353 million years
 d) Assume that the continents continue to separate at the same rate every year.
 12. a) 30 s, 60 s, 90 s, 120 s, 150 s
 b) arithmetic
 c) 60 days
 d) 915 min